Beiträge zur Algebra und Geometrie
Contributions to Algebra and Geometry
Volume 45 (2004), No. 2, 415-427.

On the Genus of the Graph of Tilting Modules

Dedicated to Idun Reiten on the occasion of her 60th birthday

Luise Unger **Michael Ungruhe**

FernUniversität Hagen, Fachbereich Mathematik
D 58084 Hagen, Germany

MSC 2000: 05C10, 16G20, 16G70

Let Λ be a finite dimensional, connected, associative algebra with unit over a field k. Let n be the number of isomorphism classes of simple Λ-modules. By $\operatorname{mod}\Lambda$ we denote the category of finite dimensional left Λ-modules.

A module ${}_\Lambda T \in \operatorname{mod}\Lambda$ is called a *tilting module* if

(i) the projective dimension $\operatorname{pd}{}_\Lambda T$ of ${}_\Lambda T$ is finite, and

(ii) $\operatorname{Ext}^i_\Lambda(T,T) = 0$ for all $i > 0$, and

(iii) there is an exact sequence $0 \to {}_\Lambda\Lambda \to {}_\Lambda T^1 \to \cdots \to {}_\Lambda T^d \to 0$ with ${}_\Lambda T^i \in \operatorname{add}{}_\Lambda T$ for all $1 \le i \le d$.

Here $\operatorname{add}{}_\Lambda T$ denotes the category of direct sums of direct summands of ${}_\Lambda T$.

Tilting modules play an important role in many branches of mathematics such as representation theory of Artin algebras or the theory of algebraic groups.

Let $\bigoplus_{i=1}^{m} T_i$ be the decomposition of ${}_\Lambda T$ into indecomposable direct summands. We call ${}_\Lambda T$ *basic* if ${}_\Lambda T_i \not\cong {}_\Lambda T_j$ for all $i \neq j$. A basic tilting module has n indecomposable direct summands.

A direct summand ${}_\Lambda M$ of a basic tilting module ${}_\Lambda T$ is called an *almost complete tilting module* if ${}_\Lambda M$ has $n-1$ indecomposable direct summands.

Let $\mathcal{T}(\Lambda)$ be the set of all non isomorphic basic tilting modules over Λ. We associate with $\mathcal{T}(\Lambda)$ a quiver $\overrightarrow{\mathcal{K}(\Lambda)}$ as follows: The vertices of $\overrightarrow{\mathcal{K}(\Lambda)}$ are the tilting modules in $\mathcal{T}(\Lambda)$, and there is an arrow ${}_\Lambda T' \to {}_\Lambda T$ if ${}_\Lambda T$ and ${}_\Lambda T'$ have a common direct summand which is an

0138-4821/93 $ 2.50 © 2004 Heldermann Verlag

almost complete tilting module and if $\operatorname{Ext}^1_\Lambda(T,T') \neq 0$. We call $\overrightarrow{\mathcal{K}(\Lambda)}$ the *quiver of tilting modules* over Λ. With $\mathcal{K}(\Lambda)$ we denote the underlying graph of $\overrightarrow{\mathcal{K}(\Lambda)}$. It has been recently shown [7] that $\mathcal{K}(\Lambda)$ is the Hasse diagram of a partial order of tilting modules which was basically introduced in [10]. From this it follows, that $\overrightarrow{\mathcal{K}(\Lambda)}$ has no oriented cycles.

If $\overrightarrow{\mathcal{K}(\Lambda)}$ is finite, then it is connected. Examples show that $\overrightarrow{\mathcal{K}(\Lambda)}$ may be rather complicated. One measure for the complicatedness of a graph G is its genus $\gamma(G)$. This is the minimal genus of an orientable surface on which G can be embedded.

The aim of these notes is to show that there are finite quivers of tilting modules of arbitrary genus. To be precise, we prove:

Theorem 1. *For all integers $r \geq 0$ there is a representation finite, connected algebra Λ_r such that $\gamma(\mathcal{K}(\Lambda_r)) = r$.*

The proof of the theorem is constructive. For each $r \in \mathbb{N}$ we give an explicit example of an algebra Λ_r and embed $\mathcal{K}(\Lambda_r)$ in an orientable surface of genus r. This gives an upper bound for $\gamma(\mathcal{K}(\Lambda_r))$. Then we use general results from graph theory to show that the bound is sharp. This will be done in Section 3. In Section 1 we recall some basic facts about tilting modules and embeddings of graphs. In Section 2 we introduce the algebras Λ_r and derive some properties of $\overrightarrow{\mathcal{K}(\Lambda_r)}$. For unexplained terminology and results from representation theory we refer to [1], and from graph theory to [8].

Acknowledgement. Part of this work was done while the first author was visiting the Centre of Advanced Studies in Oslo. She wants to thank the staff of the Centre for the warm hospitality and the possibility to enjoy the stimulating atmosphere at the Centre.

1. Preliminaries

1.1. The construction of $\overrightarrow{\mathcal{K}(\Lambda)}$

Let ${}_\Lambda M$ be a direct summand of a tilting module. A basic Λ-module ${}_\Lambda X$ is called a *complement* to ${}_\Lambda M$ if ${}_\Lambda M \oplus {}_\Lambda X$ is a tilting module and if $\operatorname{add} M \cap \operatorname{add} X = 0$. It was proved in [5] that every direct summand of a tilting module has a distinguished complement ${}_\Lambda X$ which is characterized by the fact that there is no epimorphism ${}_\Lambda E \to {}_\Lambda X$ with ${}_\Lambda E \in \operatorname{add}{}_\Lambda M$. The module ${}_\Lambda X$ is unique up to isomorphism, and it is called the *source complement* to ${}_\Lambda M$. There is the dual concept of a source complement. A complement ${}_\Lambda Y$ to ${}_\Lambda M$ is called a *sink complement* to a direct summand ${}_\Lambda M$ of a tilting module, if there is no monomorphism ${}_\Lambda Y \to {}_\Lambda E$ with ${}_\Lambda E \in \operatorname{add}{}_\Lambda M$. In contrast to source complements, sink complements do not always exist. If ${}_\Lambda M$ has a sink complement then it is unique up to isomorphism [6]. The source and the sink complement to an almost complete tilting module ${}_\Lambda M$ coincide if and only if ${}_\Lambda M$ is not faithful [4]. The following result is basically contained in [4], compare [6].

Proposition 1. *Let ${}_\Lambda M$ be a faithful almost complete tilting module. Let ${}_\Lambda X$ be a complement to ${}_\Lambda M$ which is not the sink complement to ${}_\Lambda M$. Then*

(1) *there is a complement ${}_\Lambda Y$ to ${}_\Lambda M$ which is not isomorphic to ${}_\Lambda X$,*

(2) *there is an exact sequence $\eta : 0 \to {}_\Lambda X \to {}_\Lambda E \to {}_\Lambda Y \to 0$ with ${}_\Lambda E \in \operatorname{add}{}_\Lambda M$,*

(3) $\operatorname{Ext}^i_\Lambda(X,Y) = 0$ *for all* $i > 0$, *and* $\operatorname{Ext}^i_\Lambda(Y,X) = 0$ *for all* $i > 1$,
(4) *the module* ${}_\Lambda Y$ *is uniquely determined by the property* (2).

We call η the *s*equence connecting the complements ${}_\Lambda X$ and ${}_\Lambda Y$ to ${}_\Lambda M$. This result allows an alternative definition of the quiver $\overrightarrow{\mathcal{K}(\Lambda)}$ which is more useful for calculations. The vertices are the elements from $\mathcal{T}(\Lambda)$ as above. There is an arrow ${}_\Lambda T' \to {}_\Lambda T$ in $\overrightarrow{\mathcal{K}(\Lambda)}$ if ${}_\Lambda T' = {}_\Lambda M \oplus {}_\Lambda X$ and ${}_\Lambda T = {}_\Lambda M \oplus {}_\Lambda Y$ where ${}_\Lambda X$ and ${}_\Lambda Y$ are indecomposable, and if there is an exact sequence $0 \to {}_\Lambda X \to {}_\Lambda E \to {}_\Lambda Y \to 0$ with ${}_\Lambda E \in \operatorname{add}{}_\Lambda M$.

If $\overrightarrow{\mathcal{K}(\Lambda)}$ is finite, then it is connected. Then the definition of $\overrightarrow{\mathcal{K}(\Lambda)}$ yields an algorithm to construct $\overrightarrow{\mathcal{K}(\Lambda)}$. We write the tilting module ${}_\Lambda\Lambda$ as a direct sum of indecomposable modules ${}_\Lambda\Lambda = \bigoplus_{i=1}^{n} {}_\Lambda\Lambda_i$. Then ${}_\Lambda\Lambda_i$ is the source complement to ${}_\Lambda\Lambda[i] = \bigoplus_{j\neq i} {}_\Lambda\Lambda_j$. If ${}_\Lambda\Lambda_i$ is not the sink complement to ${}_\Lambda\Lambda[i]$ we construct the exact sequence $0 \to {}_\Lambda\Lambda_i \to {}_\Lambda E_i \to {}_\Lambda Y_i \to 0$ with ${}_\Lambda E_i \in \operatorname{add}{}_\Lambda\Lambda[i]$ connecting the complements ${}_\Lambda\Lambda_i$ and ${}_\Lambda Y_i$ to ${}_\Lambda\Lambda[i]$. In this way we construct all neighbors of ${}_\Lambda\Lambda$. We now proceed analogously with the neighbors of ${}_\Lambda\Lambda$ and all vertices we constructed. Since $\overrightarrow{\mathcal{K}(\Lambda)}$ is finite and connected and has no oriented cycles this algorithm stops when we constructed all basic tilting modules over Λ.

1.2. Embeddings of graphs

Let G be a connected, finite graph with p vertices and q edges. We think of G as embedded on a surface $\mathcal{S}$. Then G forms a polyhedron of genus $\gamma(G)$. From the Euler polyhedron formula Beinecke and Harary [3] deduce the following lower bound for $\gamma(G)$ which we shall use in Section 3.

Proposition 2. *If* G *is connected and has no triangles, then* $\gamma(G) \geq \frac{1}{4}q - \frac{1}{2}(p-2)$.

In general this bound is not sharp. As an example we consider the following graph G which will become important in Section 3.



This graph has 18 vertices and 29 edges, hence the formula yields $\gamma(G) \geq -\frac{3}{4}$.

But G is not even planar, namely it contains the subgraph



which is homeomorphic to



This graph is isomorphic to the complete bigraph $K_{3,3}$: 

. Kuratowski's theorem [9] implies $\gamma(G) \geq 1$. Conversely, we draw G differently and shade some of its faces:



We push a cylinder through the lower cube, close it under the upper square, adjust the vertices and edges accordingly and obtain an embedding of G on a torus. To be precise, the

following figure shows an embedding of G on a torus:



The parallel dotted lines have to be identified. Hence $\gamma(G) = 1$.

2. The algebras Λ_r and properties of $\overrightarrow{\mathcal{K}(\Lambda_r)}$

2.1. The algebras Λ_r

Let Λ_1 be the path algebra of the quiver $\overrightarrow{\Delta}_1$:



bound by the relation $\alpha\beta = \gamma\delta$.

For all $r > 1$ let Λ_r be the path algebra of the quiver $\overrightarrow{\Delta}_r$:



bound by the relations $\alpha\beta = \gamma\delta$ and $\mathrm{rad}^2 = 0$, i.e. the composition of two consecutive arrows in $\overrightarrow{\Delta}_r \setminus \{a\}$ is zero.

The Auslander-Reiten quivers $\overrightarrow{\Gamma}_{\Lambda_r}$ of Λ_r are as follows:

$\overrightarrow{\Gamma}_{\Lambda_1}$



and for $r > 1$

$\overrightarrow{\Gamma}_{\Lambda_r}$



Here S_x denotes the simple module corresponding to the vertex x, the module P_x is the projective cover of S_x and I_x denotes the injective hull of I_x. Moreover, X is the radical of $P_d = I_a$ and $Y = I_a/\operatorname{soc} I_a$, where $\operatorname{soc} I_a$ is the socle of I_a.

For all $1 \leq i \leq r$ we identify an indecomposable Λ_i-module ${}_{\Lambda_i}M$ with the corresponding Λ_j-module ${}_{\Lambda_j}M$, $j \geq i$, whose support is Λ_i. With this identification $\overrightarrow{\Gamma}_{\Lambda_i}$ is a full, convex subquiver of $\overrightarrow{\Gamma}_{\Lambda_j}$ for all $1 \leq i < j \leq r$.

We have $\operatorname{gl\,dim}\Lambda_i = i + 1$ for all $1 \leq i \leq r$, where $\operatorname{gl\,dim}\Lambda$ denotes the global dimension of an algebra Λ. The simple module S_d is the unique indecomposable module of projective dimension 2, the modules I_d, S_1, S_2 are the unique indecomposable modules of projective dimension 3, and for all $3 \leq j \leq r$ the module S_j is the unique indecomposable module of projective dimension $j + 1$. These observations show:

Remark 1. Let $1 \leq j \leq r - 1$. A non projective indecomposable module ${}_{\Lambda_j}X$ lies in $\operatorname{mod}\Lambda_j \setminus \operatorname{mod}\Lambda_{j-1}$ if and only if $\operatorname{pd}_{\Lambda_j} X = j + 1$.

2.2. Properties of the quiver $\overrightarrow{\mathcal{K}(\Lambda_r)}$

The following technical lemmas roughly describe the structure of the quiver $\overrightarrow{\mathcal{K}(\Lambda_r)}$. Let r be an integer, $r \geq 2$, and let $1 \leq i < j \leq r$. We decompose the projective module ${}_{\Lambda_j}\Lambda_j$ into ${}_{\Lambda_j}\Lambda_j = {}_{\Lambda_j}\Lambda_i \oplus {}_{\Lambda_j}P_{ij}$. Hence ${}_{\Lambda_j}P_{ij}$ is the maximal direct summand of ${}_{\Lambda_j}\Lambda_j$ with $\operatorname{add}{}_{\Lambda_j}P_{ij} \cap \operatorname{add}{}_{\Lambda_j}\Lambda_i = 0$.

Lemma 1. *Let $1 \leq i < j \leq r$, and let ${}_{\Lambda_i}T$ and ${}_{\Lambda_i}T'$ be tilting modules over Λ_i. Then*

(a) *${}_{\Lambda_j}T \oplus {}_{\Lambda_j}M$ is a tilting module over Λ_j if and only if ${}_{\Lambda_j}M = {}_{\Lambda_j}P_{ij}$.*

(b) *${}_{\Lambda_j}T' \oplus {}_{\Lambda_j}P_{ij} \to {}_{\Lambda_j}T \oplus {}_{\Lambda_j}P_{ij}$ is an arrow in $\overrightarrow{\mathcal{K}(\Lambda_j)}$ if and only if ${}_{\Lambda_i}T' \to {}_{\Lambda_i}T$ is an arrow in $\overrightarrow{\mathcal{K}(\Lambda_i)}$.*

Proof. (a) Since ${}_{\Lambda_j}P_{ij}$ is projective, $\operatorname{Ext}^k_{\Lambda_j}(P_{ij}, T) = 0$ for all $k > 0$. Since no indecomposable direct summand of ${}_{\Lambda_j}T$ is a successor of an indecomposable direct summand of ${}_{\Lambda_j}P_{ij}$ in the Auslander-Reiten quiver of Λ_j, it follows that $\operatorname{Ext}^k_{\Lambda_j}(T, P_{ij}) = 0$ for all $k > 0$. Hence ${}_{\Lambda_j}T \oplus {}_{\Lambda_j}P_{ij}$ is a tilting module. The module ${}_{\Lambda_j}P_{ij}$ is the source and the sink complement to ${}_{\Lambda_j}T$, hence the unique complement.

(b) There is an arrow ${}_{\Lambda_j}T' \oplus {}_{\Lambda_j}P_{ij} \to {}_{\Lambda_j}T \oplus {}_{\Lambda_j}P_{ij}$ if and only if $\operatorname{Ext}^1_{\Lambda_j}(T \oplus P_{ij}, T' \oplus P_{ij}) \neq 0$ and if ${}_{\Lambda_j}T \oplus {}_{\Lambda_j}P_{ij}$ and ${}_{\Lambda_j}T' \oplus {}_{\Lambda_j}P_{ij}$ have a common direct summand which is an almost

complete tilting module. Equivalently, $\operatorname{Ext}^1_{\Lambda_i}(T,T') \neq 0$ and ${}_{\Lambda_i}T$ and ${}_{\Lambda_i}T'$ have a common direct summand which is an almost complete tilting module, hence if and only if there is an arrow ${}_{\Lambda_i}T' \to {}_{\Lambda_i}T$ in $\overrightarrow{\mathcal{K}(\Lambda_i)}$. □

In particular, we may identify $\overrightarrow{\mathcal{K}(\Lambda_i)}$ with the full convex subquiver of $\overrightarrow{\mathcal{K}(\Lambda_r)}$, $1 \leq i < r$, with vertices ${}_{\Lambda_r}T \oplus {}_{\Lambda_r}P_{ir}$ where ${}_{\Lambda_i}T$ are the tilting modules over Λ_i. With this identification, the building blocks of $\overrightarrow{\mathcal{K}(\Lambda_r)}$ are the subquivers $\overrightarrow{\mathcal{K}(\Lambda_i)} \setminus \overrightarrow{\mathcal{K}(\Lambda_{i-1})}$ of $\overrightarrow{\mathcal{K}(\Lambda_r)}$ with $1 \leq i \leq r$. To simplify the notation we denote by $\overrightarrow{\mathcal{K}(\Lambda_1)} \setminus \overrightarrow{\mathcal{K}(\Lambda_0)}$ the subquiver $\overrightarrow{\mathcal{K}(\Lambda_1)}$ of $\overrightarrow{\mathcal{K}(\Lambda_r)}$. The next lemma gives an algebraic description of the vertices in $\overrightarrow{\mathcal{K}(\Lambda_i)} \setminus \overrightarrow{\mathcal{K}(\Lambda_{i-1})}$ of $\overrightarrow{\mathcal{K}(\Lambda_r)}$ with $1 < i \leq r$.

Lemma 2. *For all $1 < i \leq r$, the subquiver $\overrightarrow{\mathcal{K}(\Lambda_i)} \setminus \overrightarrow{\mathcal{K}(\Lambda_{i-1})}$ of $\overrightarrow{\mathcal{K}(\Lambda_r)}$ has as vertices all tilting modules of projective dimension $i+1$.*

Proof. With the previous lemma, ${}_{\Lambda_r}T \in \overrightarrow{\mathcal{K}(\Lambda_i)}$ if and only if ${}_{\Lambda_r}T = {}_{\Lambda_r}T' \oplus {}_{\Lambda_r}P_{ir}$ with ${}_{\Lambda_i}T'$ a tilting module over Λ_i. Using the lemma again, ${}_{\Lambda_i}T' \notin \overrightarrow{\mathcal{K}(\Lambda_{i-1})}$ if and only if there is an indecomposable, non projective direct summand ${}_{\Lambda_r}X$ of ${}_{\Lambda_r}T'$ with ${}_{\Lambda_r}X \in \operatorname{mod}\Lambda_i \setminus \operatorname{mod}\Lambda_{i-1}$. With the remark in 2.1, this holds if and only if $\operatorname{pd}{}_{\Lambda_i}X = i+1$. □

Next we study arrows in $\overrightarrow{\mathcal{K}(\Lambda_r)}$ between vertices in different building blocks of $\overrightarrow{\mathcal{K}(\Lambda_r)}$.

Lemma 3. *Let $1 \leq i < j \leq r$. Let ${}_{\Lambda_r}T' \in \overrightarrow{\mathcal{K}(\Lambda_i)} \setminus \overrightarrow{\mathcal{K}(\Lambda_{i-1})}$ and ${}_{\Lambda_r}T \in \overrightarrow{\mathcal{K}(\Lambda_j)} \setminus \overrightarrow{\mathcal{K}(\Lambda_{j-1})}$ be tilting modules over Λ_r.*

(a) *There are no arrows ${}_{\Lambda_r}T \to {}_{\Lambda_r}T'$ in $\overrightarrow{\mathcal{K}(\Lambda_r)}$.*

(b) *If there is an arrow ${}_{\Lambda_r}T' \to {}_{\Lambda_r}T$ in $\overrightarrow{\mathcal{K}(\Lambda_r)}$ then $\operatorname{pd}{}_{\Lambda_r}T' = i+1$ and $\operatorname{pd}{}_{\Lambda_r}T = i+2$. In particular, $j = i+1$.*

Proof. (a) Assume there is an arrow ${}_{\Lambda_r}T \to {}_{\Lambda_r}T'$ in $\overrightarrow{\mathcal{K}(\Lambda_r)}$ with ${}_{\Lambda_r}T' \in \overrightarrow{\mathcal{K}(\Lambda_i)} \setminus \overrightarrow{\mathcal{K}(\Lambda_{i-1})}$ and ${}_{\Lambda_r}T \in \overrightarrow{\mathcal{K}(\Lambda_j)} \setminus \overrightarrow{\mathcal{K}(\Lambda_{j-1})}$ and $i < j$. Then ${}_{\Lambda_r}T' = {}_{\Lambda_r}\overline{T'} \oplus {}_{\Lambda_r}P_{ir}$ and ${}_{\Lambda_r}T = {}_{\Lambda_r}\overline{T} \oplus {}_{\Lambda_r}P_{jr}$, where ${}_{\Lambda_i}\overline{T'}$ and ${}_{\Lambda_j}\overline{T}$ are tilting modules over Λ_i respectively Λ_j. Note that ${}_{\Lambda_r}P_{jr}$ is a direct summand of ${}_{\Lambda_r}P_{ir}$. Then $0 \neq \operatorname{Ext}^1_{\Lambda_r}(T',T) = \operatorname{Ext}^1_{\Lambda_r}(\overline{T'},\overline{T}) = \operatorname{Ext}^1_{\Lambda_j}(\overline{T'},\overline{T})$. Since ${}_{\Lambda_r}T \in \overrightarrow{\mathcal{K}(\Lambda_j)} \setminus \overrightarrow{\mathcal{K}(\Lambda_{j-1})}$ there is an indecomposable direct summand ${}_{\Lambda_j}X \in \operatorname{mod}\Lambda_j \setminus \Lambda_{j-1}$ of ${}_{\Lambda_j}\overline{T}$ with $\operatorname{Ext}^1_{\Lambda_j}(\overline{T'},X) \neq 0$. This is a contradiction since ${}_{\Lambda_j}X$ is not a predecessor of an indecomposable direct summand of ${}_{\Lambda_j}T'$ in $\overrightarrow{\Gamma}_{\Lambda_j}$.

(b) Let ${}_{\Lambda_r}T' \to {}_{\Lambda_r}T$ in $\overrightarrow{\mathcal{K}(\Lambda_r)}$ be an arrow in $\overrightarrow{\mathcal{K}(\Lambda_r)}$ with ${}_{\Lambda_r}T' \in \overrightarrow{\mathcal{K}(\Lambda_i)} \setminus \overrightarrow{\mathcal{K}(\Lambda_{i-1})}$ and ${}_{\Lambda_r}T \in \overrightarrow{\mathcal{K}(\Lambda_j)} \setminus \overrightarrow{\mathcal{K}(\Lambda_{j-1})}$. Let $\eta : 0 \to {}_{\Lambda_r}X \to {}_{\Lambda_r}E \to {}_{\Lambda_r}Y \to 0$ be the corresponding sequence connecting the complements ${}_{\Lambda_r}X$ and ${}_{\Lambda_r}Y$, where ${}_{\Lambda_r}T' = {}_{\Lambda_r}X \oplus {}_{\Lambda_r}M$ and ${}_{\Lambda_r}T = {}_{\Lambda_r}Y \oplus {}_{\Lambda_r}M$. Since ${}_{\Lambda_r}T \in \overrightarrow{\mathcal{K}(\Lambda_j)} \setminus \overrightarrow{\mathcal{K}(\Lambda_{j-1})}$ it follows that ${}_{\Lambda_r}Y \in \operatorname{mod}\Lambda_j \setminus \operatorname{mod}\Lambda_{j-1}$, hence $\operatorname{pd}{}_{\Lambda_r}Y = j+1$. Let ${}_{\Lambda_r}Z \in \operatorname{mod}\Lambda_r$ with $\operatorname{Ext}^{j+1}_{\Lambda_r}(Y,Z) \neq 0$. We apply $\operatorname{Hom}_{\Lambda_r}(-,Z)$ to η and obtain $\operatorname{pd}{}_{\Lambda_r}X = j$. Since ${}_{\Lambda_r}T' \in \overrightarrow{\mathcal{K}(\Lambda_i)} \setminus \overrightarrow{\mathcal{K}(\Lambda_{i-1})}$ and $i \neq j$ we get $j = i+1$, the assertion. □

As a consequence we obtain

Lemma 4. *Let $r > 1$.*

(a) *There is an arrow ${}_{\Lambda_r}T' \to {}_{\Lambda_r}T$ in $\overrightarrow{\mathcal{K}(\Lambda_r)}$ with ${}_{\Lambda_r}T' \in \overrightarrow{\mathcal{K}(\Lambda_1)}$ and ${}_{\Lambda_r}T \in \overrightarrow{\mathcal{K}(\Lambda_2)} \setminus \overrightarrow{\mathcal{K}(\Lambda_1)}$ if and only if ${}_{\Lambda_r}T' = {}_{\Lambda_r}S_d \oplus {}_{\Lambda_r}M$ and ${}_{\Lambda_r}T = {}_{\Lambda_r}I_d \oplus {}_{\Lambda_r}M$.*

(b) *Let $3 \leq i \leq r$. There is an arrow ${}_{\Lambda_r}T' \to {}_{\Lambda_r}T$ in $\overrightarrow{\mathcal{K}(\Lambda_r)}$ with ${}_{\Lambda_r}T' \in \overrightarrow{\mathcal{K}(\Lambda_{i-1})}$ and ${}_{\Lambda_r}T \in \overrightarrow{\mathcal{K}(\Lambda_i)} \setminus \overrightarrow{\mathcal{K}(\Lambda_{i-1})}$ if and only if ${}_{\Lambda_r}T' = {}_{\Lambda_r}S_{i-1} \oplus {}_{\Lambda_r}M$ and ${}_{\Lambda_r}T = {}_{\Lambda_r}S_i \oplus {}_{\Lambda_r}M$.*

Proof. (a) Let ${}_{\Lambda_r}T' \to {}_{\Lambda_r}T$ be an arrow in $\overrightarrow{\mathcal{K}(\Lambda_r)}$ with ${}_{\Lambda_r}T' \in \overrightarrow{\mathcal{K}(\Lambda_1)}$ and ${}_{\Lambda_r}T \in \overrightarrow{\mathcal{K}(\Lambda_2)} \setminus \overrightarrow{\mathcal{K}(\Lambda_1)}$. Then $\operatorname{pd}{}_{\Lambda_r}T = 3$ with Lemma 2 and $\operatorname{pd}{}_{\Lambda_r}T' = 2$. Then ${}_{\Lambda_r}S_d$ is a direct summand of ${}_{\Lambda_r}T$. Moreover, Lemma 1 shows that ${}_{\Lambda_r}P_1 \oplus {}_{\Lambda_r}P_2$ is a direct summand of ${}_{\Lambda_r}T$. Hence the sequence connecting the complements is the Auslander-Reiten sequence, which implies that ${}_{\Lambda_r}T' = {}_{\Lambda_r}S_d \oplus {}_{\Lambda_r}M$ and ${}_{\Lambda_r}T = {}_{\Lambda_r}I_d \oplus {}_{\Lambda_r}M$. Conversely, if ${}_{\Lambda_r}T' = {}_{\Lambda_r}S_d \oplus {}_{\Lambda_r}M$ and ${}_{\Lambda_r}T = {}_{\Lambda_r}I_d \oplus {}_{\Lambda_r}M$, the Auslander-Reiten sequence starting in ${}_{\Lambda_r}S_d$ lies in $\operatorname{add}({}_{\Lambda_r}T \oplus {}_{\Lambda_r}T')$. Hence we obtain an arrow ${}_{\Lambda_r}T' \to {}_{\Lambda_r}T$ in $\overrightarrow{\mathcal{K}(\Lambda_r)}$ with ${}_{\Lambda_r}T' \in \overrightarrow{\mathcal{K}(\Lambda_1)}$ and ${}_{\Lambda_r}T \in \overrightarrow{\mathcal{K}(\Lambda_2)} \setminus \overrightarrow{\mathcal{K}(\Lambda_1)}$.

(b) Let $3 \leq i \leq r$ and let ${}_{\Lambda_r}T' \to {}_{\Lambda_r}T$ be an arrow in $\overrightarrow{\mathcal{K}(\Lambda_r)}$ with ${}_{\Lambda_r}T' \in \overrightarrow{\mathcal{K}(\Lambda_{i-1})}$ and ${}_{\Lambda_r}T \in \overrightarrow{\mathcal{K}(\Lambda_i)} \setminus \overrightarrow{\mathcal{K}(\Lambda_{i-1})}$. Then $\operatorname{pd}{}_{\Lambda_r}T = i + 1$ and $\operatorname{pd}{}_{\Lambda_r}T' = i$. Since $2 < i$ it follows that ${}_{\Lambda_r}S_{i-1}$ is a direct summand of of ${}_{\Lambda_r}T'$ and ${}_{\Lambda_r}S_i$ is a direct summand of of ${}_{\Lambda_r}T$. Conversely, if ${}_{\Lambda_r}T' = {}_{\Lambda_r}S_{i-1} \oplus {}_{\Lambda_r}M$ and ${}_{\Lambda_r}T = {}_{\Lambda_r}S_i \oplus {}_{\Lambda_r}M$ then the Auslander-Reiten sequence starting in ${}_{\Lambda_r}S_{i-1}$ lies in $\operatorname{add}({}_{\Lambda_r}T \oplus {}_{\Lambda_r}T')$. This yields an arrow ${}_{\Lambda_r}T' \to {}_{\Lambda_r}T$ in $\overrightarrow{\mathcal{K}(\Lambda_r)}$ with ${}_{\Lambda_r}T' \in \overrightarrow{\mathcal{K}(\Lambda_{i-1})}$ and ${}_{\Lambda_r}T \in \overrightarrow{\mathcal{K}(\Lambda_i)} \setminus \overrightarrow{\mathcal{K}(\Lambda_{i-1})}$. □

To summarize our observations in this section we obtain the following structure of $\overrightarrow{\mathcal{K}(\Lambda_r)}$:



There are arrows from vertices in $\overrightarrow{\mathcal{K}(\Lambda_i)} \setminus \overrightarrow{\mathcal{K}(\Lambda_{i-1})}$ to vertices in $\overrightarrow{\mathcal{K}(\Lambda_j)} \setminus \overrightarrow{\mathcal{K}(\Lambda_{j-1})}$ if and only if $j = i + 1$.

3. The proof of the theorem

3.1. An embedding of $\overrightarrow{\mathcal{K}(\Lambda_r)}$

We use induction on r to embed $\overrightarrow{\mathcal{K}(\Lambda_r)}$ on a surface of genus r.

Let $r = 1$. Direct calculation shows that $\overrightarrow{\mathcal{K}(\Lambda_1)}$ equals



where the parallel dotted lines have to be identified. We saw in 1.2 that the underlying graph $\mathcal{K}(\Lambda_1))$ of $\overrightarrow{\mathcal{K}(\Lambda_1)}$ has genus 1, hence can be embedded on a torus T_1. The vertices of $\overrightarrow{\mathcal{K}(\Lambda_1)}$ are the tilting modules

$$
\begin{array}{lll}
X_1^1 = P_a \oplus P_b \oplus P_c \oplus P_d, & X_2^1 = P_a \oplus P_c \oplus I_c \oplus P_d, & X_3^1 = P_a \oplus P_b \oplus I_b \oplus P_d, \\
X_4^1 = P_a \oplus I_b \oplus I_c \oplus P_d, & X_5^1 = P_b \oplus P_c \oplus X \oplus P_d, & X_6^1 = P_c \oplus S_c \oplus X \oplus P_d, \\
X_7^1 = P_b \oplus S_b \oplus X \oplus P_d, & X_8^1 = S_b \oplus S_c \oplus X \oplus P_d, & X_9^1 = P_c \oplus I_c \oplus S_c \oplus P_d, \\
X_{10}^1 = S_b \oplus S_c \oplus Y \oplus P_d, & X_{11}^1 = S_c \oplus I_c \oplus Y \oplus P_d, & X_{12}^1 = S_b \oplus I_b \oplus Y \oplus P_d, \\
X_{13}^1 = I_b \oplus I_c \oplus Y \oplus P_d, & X_{14}^1 = P_b \oplus I_b \oplus S_b \oplus P_d, & X_{15}^1 = P_b \oplus P_c \oplus S_d \oplus P_d, \\
X_{16}^1 = P_c \oplus I_c \oplus S_d \oplus P_d, & X_{17}^1 = P_b \oplus I_b \oplus S_d \oplus P_d, & X_{18}^1 = I_b \oplus I_c \oplus S_d \oplus P_d.
\end{array}
$$

Let $r = 2$. The quiver $\overrightarrow{\mathcal{K}(\Lambda_1)}$ is the full convex subquiver of $\overrightarrow{\mathcal{K}(\Lambda_2)}$ with vertices ${}_{\Lambda_2}X_i^2 = {}_{\Lambda_2}X_i^1 \oplus_{\Lambda_2} P_{12}$, where ${}_{\Lambda_2}P_{12} = {}_{\Lambda_2}P_1 \oplus_{\Lambda_2} P_2$. The quiver $\overrightarrow{\mathcal{K}(\Lambda_2)} \backslash \overrightarrow{\mathcal{K}(\Lambda_1)}$ is the full convex subquiver of $\overrightarrow{\mathcal{K}(\Lambda_2)}$ with vertices the tilting modules of projective dimension 3. Direct calculations show that $\overrightarrow{\mathcal{K}(\Lambda_2)} \setminus \overrightarrow{\mathcal{K}(\Lambda_1)}$ is



where we identify along the parallel horizontal, respectively vertical lines. It follows that $\overrightarrow{\mathcal{K}(\Lambda_2)} \setminus \overrightarrow{\mathcal{K}(\Lambda_1)}$ can be embedded on a torus T_2. The vertices of $\overrightarrow{\mathcal{K}(\Lambda_2)} \setminus \overrightarrow{\mathcal{K}(\Lambda_1)}$ are the tilting modules

$$
\begin{array}{ll}
Y_1^2 = P_d \oplus I_d \oplus I_c \oplus I_b \oplus P_1 \oplus S_1, & Y_2^2 = P_d \oplus I_d \oplus I_c \oplus P_c \oplus P_1 \oplus S_1, \\
Y_3^2 = P_d \oplus I_d \oplus I_c \oplus P_c \oplus P_1 \oplus P_2, & Y_4^2 = P_d \oplus I_d \oplus I_b \oplus I_c \oplus P_1 \oplus P_2, \\
Y_5^2 = P_d \oplus I_d \oplus I_c \oplus I_b \oplus S_2 \oplus S_1, & Y_6^2 = P_d \oplus I_d \oplus I_c \oplus P_c \oplus S_2 \oplus S_1, \\
Y_7^2 = P_d \oplus I_d \oplus I_c \oplus P_c \oplus S_2 \oplus P_2, & Y_8^2 = P_d \oplus I_d \oplus I_c \oplus I_b \oplus S_2 \oplus P_2, \\
Y_9^2 = P_d \oplus I_d \oplus P_b \oplus I_b \oplus S_2 \oplus S_1, & Y_{10}^2 = P_d \oplus I_d \oplus P_b \oplus P_c \oplus S_2 \oplus S_1, \\
Y_{11}^2 = P_d \oplus I_d \oplus P_b \oplus P_c \oplus S_2 \oplus P_2, & Y_{12}^2 = P_d \oplus I_d \oplus P_b \oplus I_b \oplus S_2 \oplus P_2, \\
Y_{13}^2 = P_d \oplus I_d \oplus P_b \oplus I_b \oplus P_1 \oplus S_1, & Y_{14}^2 = P_d \oplus I_d \oplus P_b \oplus P_c \oplus P_1 \oplus S_1, \\
Y_{15}^2 = P_d \oplus I_d \oplus P_b \oplus P_c \oplus P_1 \oplus P_2, & Y_{16}^2 = P_d \oplus I_d \oplus P_b \oplus I_b \oplus P_1 \oplus P_2.
\end{array}
$$

The subquivers $\overrightarrow{Q}_1$: $\begin{array}{ccc} X_{17}^2 & \longleftarrow & X_{15}^2 \\ \downarrow & & \downarrow \\ X_{18}^2 & \longleftarrow & X_{16}^2 \end{array}$ and $\overrightarrow{Q}_2$: $\begin{array}{ccc} Y_{16}^2 & \longleftarrow & Y_{15}^2 \\ \downarrow & & \downarrow \\ Y_4^2 & \longleftarrow & Y_3^2 \end{array}$

bound squares on T_1 respectively T_2. In $\overrightarrow{\mathcal{K}(\Lambda_2)}$ they are joint as follows:



We cut out the interiors of $\overrightarrow{Q}_1$ on T_1 and $\overrightarrow{Q}_2$ on T_2 and insert a cylinder connecting T_1 and T_2. We obtain a surface of genus 2 on which $\overrightarrow{\mathcal{K}(\Lambda_2)}$ can be embedded:



Let $r > 2$. We abbreviate the injective Λ_r-module ${}_{\Lambda_r}P_d \oplus {}_{\Lambda_r}I_d$ by ${}_{\Lambda_r}I$ and the projective-injective Λ_r-module $\bigoplus_{i=3}^{r} {}_{\Lambda_r}P_i$ by ${}_{\Lambda_r}P_{2r}$. Direct calculations show that $\overrightarrow{\mathcal{K}(\Lambda_r)} \setminus \overrightarrow{\mathcal{K}(\Lambda_{r-1})}$ is



with

$$Z_1 = I \oplus P_{2r} \oplus S_r \oplus I_c \oplus P_c \oplus S_1, \quad Z_2 = I \oplus P_{2r} \oplus S_r \oplus P_b \oplus I_c \oplus S_1,$$
$$Z_3 = I \oplus P_{2r} \oplus S_r \oplus P_b \oplus P_c \oplus P_2, \quad Z_4 = I \oplus P_{2r} \oplus S_r \oplus I_c \oplus P_c \oplus P_2,$$
$$Z_5 = I \oplus P_{2r} \oplus S_r \oplus I_c \oplus I_b \oplus S_1, \quad Z_6 = I \oplus P_{2r} \oplus S_r \oplus P_b \oplus I_b \oplus P_2$$
$$Z_7 = I \oplus P_{2r} \oplus S_r \oplus P_b \oplus I_b \oplus P_2, \quad Z_8 = I \oplus P_{2r} \oplus S_r \oplus I_c \oplus I_b \oplus P_2.$$

We assume by induction that $\overrightarrow{\mathcal{K}(\Lambda_{r-1})}$ is embedded on a surface $\mathcal{S}_{r-1}$ of genus $r-1$ such that

a)

$$\overrightarrow{Q}_1: \begin{array}{ccc} Z'_5 & \longleftarrow & Z'_1 \\ \uparrow & & \uparrow \\ Z'_6 & \longleftarrow & Z'_2 \end{array} \quad \text{and} \quad \overrightarrow{Q}_2: \begin{array}{ccc} Z'_4 & \longrightarrow & Z'_8 \\ \uparrow & & \uparrow \\ Z'_3 & \longrightarrow & Z'_7 \end{array}$$

or

b)

$$\overrightarrow{Q}_3: \begin{array}{ccc} Z'_6 & \longleftarrow & Z'_7 \\ \uparrow & & \uparrow \\ Z'_2 & \longleftarrow & Z'_3 \end{array} \quad \text{and} \quad \overrightarrow{Q}_4: \begin{array}{ccc} Z'_5 & \longleftarrow & Z'_8 \\ \uparrow & & \uparrow \\ Z'_1 & \longleftarrow & Z'_4 \end{array}$$

bound squares on $\mathcal{S}_{r-1}$. Here Z'_i, $1 \le i \le 4$, denotes the Λ_{r-1}-module which we obtain when we replace the direct summand S_r of Z_i by S_{r-1} and the direct summand P_{2r} by $P_{2,r-1}$. For $r-1=2$, let $P_{2,r-1}=0$.

Note that this assumption is satisfied for $r-1=2$. We embedded $\overrightarrow{\mathcal{K}(\Lambda_2)}$ on a surface $\mathcal{S}_2$ of genus 2 and the subquivers

$$\begin{array}{ccc} Z'_5 = Y_5^2 & \longleftarrow & Z'_1 = Y_6^2 \\ \uparrow & & \uparrow \\ Z'_6 = Y_9^2 & \longleftarrow & Z'_2 = Y_{10}^2 \end{array} \quad \text{and} \quad \begin{array}{ccc} Z'_4 = Y_7^2 & \longrightarrow & Z'_8 = Y_8^2 \\ \uparrow & & \uparrow \\ Z'_3 = Y_{11}^2 & \longrightarrow & Z'_7 = Y_{12}^2 \end{array}$$

bound squares on $\mathcal{S}_2$.

The quiver $\overrightarrow{\mathcal{K}(\Lambda_{r-1})}$ is the full convex subquiver of $\overrightarrow{\mathcal{K}(\Lambda_r)}$ with vertices the tilting modules over Λ_r of the form ${}_{\Lambda_r}T \oplus {}_{\Lambda_r}P_r$, where ${}_{\Lambda_r}T$ is a tilting module over Λ_{r-1}. Note that there is an arrow ${}_{\Lambda_r}Z''_i = {}_{\Lambda_r}Z'_i \oplus {}_{\Lambda_r}P_r \to {}_{\Lambda_r}Z_i$ in $\overrightarrow{\mathcal{K}(\Lambda_r)}$.

Let us assume first that we are in the situation (a). We cut out the interiors of the squares $\overrightarrow{Q}_1$ and $\overrightarrow{Q}_2$ and insert a handle



On this handle we embed $\overrightarrow{\mathcal{K}(\Lambda_r)} \setminus \overrightarrow{\mathcal{K}(\Lambda_{r-1})}$ and the arrows joining Z''_i and Z'_i:



This yields an embedding of $\overrightarrow{\mathcal{K}(\Lambda_r)}$ on a surface $\mathcal{S}_r$ of genus r and the squares

$$\begin{array}{ccc} Z'_6 & \longleftarrow & Z'_7 \\ \uparrow & & \uparrow \\ Z'_2 & \longleftarrow & Z'_3 \end{array} \quad \text{and} \quad \begin{array}{ccc} Z'_5 & \longrightarrow & Z'_8 \\ \uparrow & & \uparrow \\ Z'_1 & \longrightarrow & Z'_4 \end{array}$$

bound squares on $\mathcal{S}_r$.

We proceed analogously in case (b), and it follows that $\gamma(\mathcal{K}(\Lambda_r)) \leq r$.

3.2. A lower bound for $\gamma(\mathcal{K}(\Lambda_r))$

If $r = 1$, then $\gamma(\mathcal{K}(\Lambda_1)) = 1$ as it was shown in 1.2. Hence we may assume that $r > 1$.

Consider $\overrightarrow{\mathcal{K}(\Lambda_r)} \setminus \overrightarrow{\mathcal{K}(\Lambda_1)}$. We embedded this quiver on a surface of genus $r-1$, hence $\gamma(\mathcal{K}(\Lambda_r) \setminus \mathcal{K}(\Lambda_1)) \leq r-1$. The graph $\mathcal{K}(\Lambda_2) \setminus \mathcal{K}(\Lambda_1)$ has 16 vertices and 32 edges. For all $2 \leq i \leq r$, the graphs $\mathcal{K}(\Lambda_i) \setminus \mathcal{K}(\Lambda_{i-1})$ have 8 vertices and 12 edges. Moreover, there are 8 edges joining vertices in $\mathcal{K}(\Lambda_i) \setminus \mathcal{K}(\Lambda_{i-1})$ with vertices in $\mathcal{K}(\Lambda_{i+1}) \setminus \mathcal{K}(\Lambda_i)$, $2 < i < r$. Hence $\mathcal{K}(\Lambda_r) \setminus \mathcal{K}(\Lambda_1)$ has $p = 16+8(r-2)$ vertices and $q = 32+20(r-2)$ edges. Since $\mathcal{K}(\Lambda_r) \setminus \mathcal{K}(\Lambda_1)$ has no triangles we may use the formula in 1.2 which gives $\gamma(\mathcal{K}(\Lambda_r) \setminus \mathcal{K}(\Lambda_1) \geq \frac{1}{4}q - \frac{1}{2}(p-2) = r-1$, hence $\gamma(\mathcal{K}(\Lambda_r) \setminus \mathcal{K}(\Lambda_1)) = r-1$.

We saw above that there are 4 arrows $\alpha_1, \alpha_2, \alpha_3, \alpha_4$ joining vertices in $\overrightarrow{\mathcal{K}(\Lambda_1)}$ with vertices in $\overrightarrow{\mathcal{K}(\Lambda_2)} \setminus \overrightarrow{\mathcal{K}(\Lambda_1)}$. Let $\overrightarrow{\mathcal{K}(\Lambda_r)}'$ be the subquiver of $\overrightarrow{\mathcal{K}(\Lambda_r)}$ which we obtain by deleting three of these arrows, say $\alpha_2, \alpha_3, \alpha_4$. Then $\gamma(\mathcal{K}(\Lambda_r)) \geq \gamma(\mathcal{K}(\Lambda_r)')$. The blocks of $\mathcal{K}(\Lambda_r)'$, i.e. the maximal connected subgraphs of $\mathcal{K}(\Lambda_r)'$ which are connected, non trivial and have no cutpoints are $\mathcal{K}(\Lambda_1)$, $\circ \overset{\alpha_1}{—} \circ$ and $\mathcal{K}(\Lambda_r) \setminus \mathcal{K}(\Lambda_1)$. Since the genus of a graph is the sum of the genera of its blocks [2], we obtain that

$$\gamma(\mathcal{K}(\Lambda_r)) \geq \gamma(\mathcal{K}(\Lambda_r)') = \gamma(\mathcal{K}(\Lambda_1)) + \gamma(\circ — \circ) + \gamma(\mathcal{K}(\Lambda_r) \setminus \mathcal{K}(\Lambda_1)) = 1 + 0 + r - 1.$$

Hence $\gamma(\mathcal{K}(\Lambda_r)) = r$. To finish the proof of the theorem we have to show that there is an algebra Λ_0 with $\gamma(\mathcal{K}(\Lambda_0)) = 0$. If Λ_0 is the ground field, then $\mathcal{K}(\Lambda_0)$ consists of a single vertex, hence it has genus 0.

References

[1] Auslander, M.; Reiten, I.; Smalø, S.: *Representation Theory of Artin Algebras*. Cambridge University Press 1995. Zbl 0834.16001

[2] Battle, J.; Harary, F.; Kodama, Y.; Youngs, J. W. T.: *Additivity of the genus of a graph*. Bull. Amer. Math. Soc. **68** (1962), 565–568. Zbl 0142.41501

[3] Beinecke, L. W.; Harary, F.: *Inequalities involving the genus of a graph and its thickness*. Proc. Glasgow Math. Assoc. **7** (1965), 19–21.

[4] Coelho, F.; Happel, D.; Unger, L.: *Complements to partial tilting modules*. J. Algebra **170** (1994), 184–205. Zbl 0834.16012

[5] Happel, D.; Unger, L.: *Partial tilting modules and covariantly finite subcategories*. Comm. Algebra **22**(5) (1994), 1723–1727. Zbl 0804.16015

[6] Happel, D.; Unger, L.: *Complements and the generalized Nakayama Conjecture*. Proc. ICRA VIII, CMS Conference Proc. **24** (1998), 293–310. Zbl 0944.16010

[7] Happel, D.; Unger, L.: *On a partial order of tilting modules*. To appear in Algebras and Representation Theory.

[8] Harary, F.: *Graph Theory*. Addison Wesley Publishing Company 1969. Zbl 0182.57702

[9] Kuratowski, K.: *Sur le problème de courbes gauches en topologie*. Fund. Math. **15** (1930), 271–283.

[10] Riedtmann, C.; Schofield, A.: *On a simplicial complex associated with tilting modules*. Comment. Math. Helv. **66** (1991), 70–78. Zbl 0790.16013

Received December 10, 2002