Beiträge zur Algebra und Geometrie
Contributions to Algebra and Geometry
Volume 51 (2010), No. 1, 209-227.

Projectivity and Flatness over the Colour Endomorphism Ring of a Finitely Generated Graded Comodule

T. Guédénon

110 Penworth Drive S.E., Calgary, AB, T2A 5H4, Canada
e-mail: guedenth@yahoo.ca

Abstract. Let k be a field, G an abelian group with a bicharacter, A a colour algebra; i.e., an associative graded k-algebra with identity, $\mathcal{C}$ a graded A-coring that is projective as a right A-module, $\mathcal{C}^*$ the graded dual ring of $\mathcal{C}$ and Λ a left graded $\mathcal{C}$-comodule that is finitely generated as a graded right $\mathcal{C}^*$-module. We give necessary and sufficient conditions for projectivity and flatness of a graded module over the colour endomorphism ring ${}^{\mathcal{C}}END(\Lambda)$.

0. Introduction

The notion of graded corings (except graded algebras and graded coalgebras) rarely appears in the literature on corings. The only paper we know where this notion appears is [8]. In the present paper we will give some conditions to test projectivity or flatness over the colour endomorphism ring of a finitely generated graded $\mathcal{C}$-comodule, where $\mathcal{C}$ is a graded coring. Let k be a field, A a k-algebra, $\mathcal{C}$ an A-coring, ${}^*\mathcal{C}$ the left dual ring of $\mathcal{C}$ and Λ a right $\mathcal{C}$-comodule that is finitely generated as a left ${}^*\mathcal{C}$-module. In [11], we gave necessary and sufficient conditions for projectivity and flatness over the endomorphism ring $End^{\mathcal{C}}(\Lambda)$ of Λ. In the present paper, we will extend these results to a G-graded A-coring $\mathcal{C}$, where G is an abelian group with a bicharacter and A is a colour algebra; i.e., a graded associative k-algebra with identity. More precisely, let us denote by $\mathcal{C}^* = HOM_A(\mathcal{C}_A, A_A)$ the largest graded vector space contained in $Hom_A(\mathcal{C}_A, A_A)$. It has a colour algebra structure. Let Λ be a graded left $\mathcal{C}$-comodule that is finitely generated as a

0138-4821/93 $ 2.50 © 2010 Heldermann Verlag

graded right $\mathcal{C}^*$-module. We give necessary and sufficient conditions for projectivity and flatness over the colour endomorphism ring ${}^{\mathcal{C}}END(\Lambda)$ of Λ. The presence of the bicharacter makes the difference with the classical gradation. These results are interesting when $\mathcal{C} = A$, or when $\mathcal{C}$ contains a grouplike element, or when $\mathcal{C}$ comes from a graded entwining structure with respect to a bicharacter. If $\mathcal{C} = A$ then ${}^{gr-A}\mathcal{M}$ is the category of graded left A-modules and A^* is isomorphic to the opposite algebra A^{op} of A. If $\mathcal{C}$ contains a grouplike element X, then A is a graded left $\mathcal{C}$-comodule that is finitely generated as a graded right $\mathcal{C}^*$-module. In this case, ${}^{\mathcal{C}}END(A)$ is the colour subring of $(\mathcal{C}, X)$-coinvariants of A. Our techniques and methods are inspired from [10], [7] and [11].

1. Preliminary results

Throughout the paper, k is a field, G is an abelian group and $(./.)$ is a bicharacter on G; i.e., a map from $G \times G$ into $k^\times$ satisfying:

$$(x/y) = (y/x)^{-1} \quad \text{and} \quad (x/y+z) = (x/y)(x/z).$$

These two relations imply that $(x+y/z) = (x/z)(y/z)$. If M and N are vector spaces $Hom(M,N)$ is the vector space of k-linear maps from M to N.

A vector space A is G-graded or graded if $A = \oplus_{x\in G} A_x$, where the A_x are vector subspaces of A. An algebra A (not necessarily associative with identity) is said to be graded if A is a graded vector space as above and the A_x satisfy $A_x A_y \subseteq A_{x+y}$. According to [8, Section 1], a colour algebra is an associative graded algebra. In what follows we assume that all colour algebras are unital. We will consider k as a colour algebra with the trivial gradation. Given colour algebras A and B, a morphism of colour algebras $A \to B$ is a morphism of algebras which is homogeneous of degree 0. Let m be an element of a graded vector space M. If m is homogeneous, we denote by $|m|$ its degree. If $|m|$ occurs in some expression, this means that we regard m as a homogeneous element and that the expression extends to the other elements by linearity. Let M and N be graded vector spaces. An element of $Hom(M,N)$ is homogeneous of degree x if $f(M_y) \subseteq N_{x+y}$ for all $y \in G$. We denote by $HOM(M,N)_x$ the vector subspace of $Hom(M,N)$ whose elements are homogeneous of degree x and we will set $HOM(M,N) = \oplus_{x\in G} HOM(M,N)_x$. Clearly, $HOM(M,N)$ is the largest graded vector space contained in $Hom(M,N)$. The space $HOM(M,N)$ is denoted $Hom_k(M,N)_G$ in [9]. By [13, Corollary 1.2.11], $HOM(M,N) = Hom(M,N)$ if G is finite or if M is finite-dimensional. By [9], $HOM(M,M)$ is a colour algebra. If M, N, M' and N' are graded vector spaces and if $f : M \to M'$ and $g : N \to N'$ are homogeneous linear maps then $(f \otimes g)(m \otimes n) = (|g|/|m|) f(m) \otimes g(n)$. We will denote by ${}_{gr-k}\mathcal{M}$ the category of graded k-vector spaces. The morphisms of ${}_{gr-k}\mathcal{M}$ are the homogeneous k-linear maps of degree 0; we call them the graded k-linear maps. Let N be a graded vector space. For every x in G, the x-suspension of N is the graded vector space $N(x)$ obtained from N by a shift of the gradation by x. As vector spaces, N and $N(x)$ coincide but the gradations are related by $N(x)_y = N_{x+y}$ for all $y \in G$.

Let A be a colour algebra. A left A-module M is called a graded left A-module if M admits a decomposition as a direct sum of vector spaces $M = \oplus_{x\in G} M_x$ such that $A_x M_y \subseteq M_{x+y}$; $\forall x,\, y \in G$.

Definition 1.1. *Let M, N be graded left A-modules. A homogeneous element f of $Hom(M,N)$ is colour left A-linear if $f(am) = (|f|/|a|)af(m)$ for all $a \in A$.*

If M, N are graded left A-modules, we let ${}_AHOM(M,N)_x$ denote the vector subspace of $Hom(M,N)$ whose elements are colour A-linear of degree x. So the colour left A-linear maps of degree 0 are exactly the left A-linear maps of degree 0; i.e., ${}_AHOM(M,N)_0 = {}_AHom(M,N) \cap HOM(M,N)_0$. We define ${}_AHOM(M,N)$ to be the sum of these subspaces; the sum is direct: ${}_AHOM(M,N) = \oplus_{x\in G}\, {}_AHOM(M, N)_x$. We call ${}_AHOM(M,N)$ the subspace of colour left A-linear maps of $Hom(M, N)$. Contrary to the classical gradation, if $A \neq k$ and if the bicharacter is not trivial, there is no comparison relation between ${}_AHOM(M,N)$ and ${}_AHom(M,N)$ even if M is finitely generated as an A-module or if G is finite. If $G = \mathbb{Z}/2\mathbb{Z}$, colour A-linear maps are called A-superlinear in [16]. We will denote by ${}_{gr-A}\mathcal{M}$ the category of graded left A-modules. The morphisms of ${}_{gr-A}\mathcal{M}$ are the colour left A-linear maps of degree 0; we call them the graded left A-linear maps. It is well known that ${}_{gr-A}\mathcal{M}$ is a Grothendieck category. We can define in a similar way a graded right A-module and a graded A-bimodule. A colour right A-linear map of degree x is just a homogeneous right A-linear map of degree x. To establish our main results we will need the following well-known results of graded ring theory.

- If N is a graded left (right) A-module, $N(x)$ is a graded left (right) A-module which coincides with N as a graded left (right) A-module.

- An object of ${}_{gr-A}\mathcal{M}$ is projective (resp. flat) in ${}_{gr-A}\mathcal{M}$ if and only if it is projective (resp. flat) in ${}_A\mathcal{M}$, the category of left A-modules.

- An object of ${}_{gr-A}\mathcal{M}$ is free in ${}_{gr-A}\mathcal{M}$ if it has an A-basis consisting of homogeneous elements or equivalently, if it is isomorphic to some $\oplus_{i\in I} A(x_i)$, where $(x_i, i \in I)$ is a family of elements of G.

- An object of ${}_{gr-A}\mathcal{M}$ is called finitely generated if it is a quotient of a free graded module of finite rank $\oplus_{i\leq m} A(x_i)$, where the $x_i \in G$ and m is a natural integer.

- Any object of ${}_{gr-A}\mathcal{M}$ is a quotient of a free object in ${}_{gr-A}\mathcal{M}$, and any projective object in ${}_{gr-A}\mathcal{M}$ is isomorphic in ${}_{gr-A}\mathcal{M}$ to a direct summand of a free object.

- An object of ${}_{gr-A}\mathcal{M}$ is flat in ${}_{gr-A}\mathcal{M}$ if and only if it is the inductive limit of finitely generated free objects in ${}_{gr-A}\mathcal{M}$.

- An object Λ of ${}_{gr-A}\mathcal{M}$ is called finitely presented if there is an exact sequence $\oplus_{i\leq m} A(x_i) \to \oplus_{j\leq n} A(y_j) \to \Lambda \to 0$ for $x_i,\, y_j \in G$ and some natural integers m and n. A finitely presented graded module is finitely generated.

Lemma 1.2. *Let A be a colour algebra and M a graded left A-module which is generated as A-module by a homogeneous element m of degree 0. Then M is finitely generated as a graded left A-module.*

Proof. We have $M = Am$. The k-linear map $f : A \to M$; $a \mapsto am$ is surjective, homogeneous of degree 0 and left A-linear. So f is an epimorphism in ${}_{gr-A}\mathcal{M}$. □

An A-coring $\mathcal{C}$ is an A-bimodule together with two A-bimodule maps $\Delta_{\mathcal{C}} : \mathcal{C} \to \mathcal{C} \otimes_A \mathcal{C}$ and $\epsilon_{\mathcal{C}} : \mathcal{C} \to A$ such that the usual coassociativity and counit properties hold. Let $\mathcal{C}$ be an A-coring. A left $\mathcal{C}$-comodule is a left A-module M together with a left A-linear map $\rho_{M,\mathcal{C}} : M \to \mathcal{C} \otimes_A M$ such that

$$(\epsilon_{\mathcal{C}} \otimes_A id_M) \circ \rho_{M,\mathcal{C}} = id_M, \quad \text{and} \quad (\Delta_{\mathcal{C}} \otimes_A id_M) \circ \rho_{M,\mathcal{C}} = (id_{\mathcal{C}} \otimes_A \rho_{M,\mathcal{C}}) \circ \rho_{M,\mathcal{C}}.$$

For more details on corings, we refer to [1], [2], [3], [4] and [5].

An A-coring $\mathcal{C}$ is called a graded A-coring if $\mathcal{C}$ admits a decomposition as a direct sum of vector spaces $\mathcal{C} = \oplus_x \mathcal{C}_x$ such that $\mathcal{C}$ is a graded A-bimodule, and $\Delta_{\mathcal{C}}$ and $\epsilon_{\mathcal{C}}$ are graded left and right A-linear maps. Note that $\epsilon_{\mathcal{C}}(c) = 0$ if c is homogeneous of degree $|c| \neq 0$. We use the notation-type of Sweedler-Heyneman for $\Delta_{\mathcal{C}}$ but we will omit the parentheses on subscripts. So for every homogeneous element $c \in \mathcal{C}$ we will write $\Delta_{\mathcal{C}}(c) = \sum_{|c|} c_1 \otimes_A c_2$; where $\sum_{|c|} = \sum_{|c_1|+|c_2|=|c|}$. We have $\sum_{|c|} \sum_{|c_1|} c_{11} \otimes_A c_{12} \otimes_A c_2 = \sum_{|c|} \sum_{|c_2|} c_1 \otimes_A c_{21} \otimes_A c_{22}$. Note that $\epsilon_{\mathcal{C}}(c) = 0$ if $|c| \neq 0$. A left $\mathcal{C}$-comodule M is called a graded left $\mathcal{C}$-comodule if M admits a decomposition as a direct sum of vector spaces $M = \oplus_x M_x$ such that $\rho_{M,\mathcal{C}}$ is homogeneous of degree 0; i.e., $\rho_{M,\mathcal{C}}$ is a graded left A-linear map. We will write $\rho_{M,\mathcal{C}}(m) = \sum_{|m|} m_{(-1)} \otimes_A m_{(0)}$, where $\sum_{|m|} = \sum_{|m_{(-1)}|+|m_{(0)}|=|m|}$.

Any colour algebra A is a graded A-coring called the trivial A-coring, and a graded k-coalgebra is a graded k-coring. A morphism of graded left $\mathcal{C}$-comodules $f : M \to N$ is a morphism in ${}_{gr-A}\mathcal{M}$ such that

$$\rho_{N,\mathcal{C}} \circ f = (id_{\mathcal{C}} \otimes_A f) \circ \rho_{M,\mathcal{C}}, \quad \text{that is}$$

$$\sum_{|m|} f(m)_{(-1)} \otimes_A f(m)_{(0)} = \sum_{|m|} m_{(-1)} \otimes_A f(m_{(0)}) \quad \forall m \in M.$$

A morphism of graded left $\mathcal{C}$-comodule will be called a graded left $\mathcal{C}$-colinear map. We denote by ${}^{gr-\mathcal{C}}\mathcal{M}$ the category of graded left $\mathcal{C}$-comodules. The morphisms of ${}^{gr-\mathcal{C}}\mathcal{M}$ are the graded left $\mathcal{C}$-colinear maps. The category ${}^{gr-\mathcal{C}}\mathcal{M}$ has direct sums. If $\mathcal{C}$ is projective as a right A-module, then ${}^{gr-\mathcal{C}}\mathcal{M}$ is a Grothendieck category ([4] for the ungraded case).

Definition 1.3. *Let $\mathcal{C}$ be a graded A-coring and M, N be objects of ${}^{gr-\mathcal{C}}\mathcal{M}$. A homogeneous element $f \in Hom(M, N)$ is colour left $\mathcal{C}$-colinear if f is colour left A-linear and $\rho_{N,\mathcal{C}} \circ f = (id_{\mathcal{C}} \otimes_A f) \circ \rho_{M,\mathcal{C}}$.*

It follows from Definition 1.3 that a graded left $\mathcal{C}$-colinear map is a colour left $\mathcal{C}$-colinear map of degree 0. If M and N are objects of ${}^{gr-\mathcal{C}}\mathcal{M}$ and $x \in G$, we will denote by ${}^{\mathcal{C}}HOM(M, N)_x$ the vector subspace of $Hom(M, N)$ whose elements are colour left $\mathcal{C}$-colinear of degree x. So we have

$${}^{\mathcal{C}}HOM(M, N)_x = \{f \in {}_AHOM(M, N)_x, \sum_{|m|} f(m)_{(-1)} \otimes_A f(m)_{(0)} =$$

$$\sum_{|m|}(|f|,|m_{(-1)}|)m_{(-1)} \otimes_A f(m_{(0)})\}.$$

We will set ${}^{\mathcal{C}}HOM(M,N) = \oplus_{x\in G}{}^{\mathcal{C}}HOM(M,N)_x$. We call ${}^{\mathcal{C}}HOM(M,N)$ the subspace of colour left $\mathcal{C}$-colinear maps of $Hom(M,N)$. We can define in a similar way a graded right $\mathcal{C}$-comodule. A homogeneous colour right $\mathcal{C}$-colinear map is just a homogeneous right $\mathcal{C}$-colinear.

If N is a graded left $\mathcal{C}$-comodule, then for every x in G, the x-suspension $N(x)$ is a graded left $\mathcal{C}$-comodule which coincides with N as a $\mathcal{C}$-comodule. By [17], the linear map $i_{-x} : N \to N(x)$ defined by $i_{-x}(n) = (-x/|n|)n$ is bijective and homogeneous of degree $-x$. It is obviously colour left $\mathcal{C}$-colinear.

Lemma 1.4. *Let $\mathcal{C}$ be a graded A-coring, and M, N be graded left $\mathcal{C}$-comodules. For every $x \in G$, the linear map ${}^{\mathcal{C}}HOM(M,N)_x \to {}^{\mathcal{C}}HOM(M,N(x))_0; f \mapsto i_{-x} \circ f$, where i_{-x} is defined above is an isomorphism of vector spaces.*

Lemma 1.5. *Let P be an object of ${}^{gr-\mathcal{C}}\mathcal{M}$. Then the functor ${}^{\mathcal{C}}HOM(P,-) : {}^{gr-\mathcal{C}}\mathcal{M} \to_{gr-k} \mathcal{M}$ is left exact.*

Proof. Let $0 \to L \to M \to N \to 0$ be an exact sequence in ${}^{gr-\mathcal{C}}\mathcal{M}$; so $0 \to L(x) \to M(x) \to N(x) \to 0$ is exact in ${}^{gr-\mathcal{C}}\mathcal{M}$ for every $x \in G$. By [13, Corollary 1.2.2], P is projective in ${}_{gr-k}\mathcal{M}$. So the sequence

$$0 \to HOM(P,L(x))_0 \to HOM(P,M(x))_0 \to HOM(P,N(x))_0 \to 0$$

is exact for every $x \in G$. It follows from Lemma 1.4 that

$$0 \to HOM(P,L)_x \to HOM(P,M)_x \to HOM(P,N)_x \to 0$$

is an exact sequence for every $x \in G$. We know that $i \circ f \in {}^{\mathcal{C}}HOM(P,M)_x$ for all $f \in {}^{\mathcal{C}}HOM(P,L)_x$. This means that the sequence

$$0 \to {}^{\mathcal{C}}HOM(P,L)_x \to {}^{\mathcal{C}}HOM(P,M)_x \to {}^{\mathcal{C}}HOM(P,N)_x$$

is exact for every $x \in G$; i.e.,

$$0 \to {}^{\mathcal{C}}HOM(P,L) \to {}^{\mathcal{C}}HOM(P,M) \to {}^{\mathcal{C}}HOM(P,N)$$

is an exact sequence. So the functor ${}^{\mathcal{C}}HOM(P,-)$ is left exact. □

We say that an object P of ${}^{gr-\mathcal{C}}\mathcal{M}$ is projective if the functor ${}^{\mathcal{C}}HOM(P,-)_0$ is exact.

Lemma 1.6. *Let $\mathcal{C}$ be a graded A-coring. An object P of ${}^{gr-\mathcal{C}}\mathcal{M}$ is projective in ${}^{gr-\mathcal{C}}\mathcal{M}$ if and only if the functor ${}^{\mathcal{C}}HOM(P,-)$ is exact.*

Proof. Assume that ${}^{\mathcal{C}}HOM(P,-)$ is exact in ${}^{gr-\mathcal{C}}\mathcal{M}$. Let $0 \to L \to M \to N \to 0$ be an exact sequence in ${}^{gr-\mathcal{C}}\mathcal{M}$. So the sequence $0 \to {}^{\mathcal{C}}HOM(P,L) \to {}^{\mathcal{C}}HOM(P,M) \to {}^{\mathcal{C}}HOM(P,N) \to 0$ is exact. It follows that the sequence $0 \to {}^{\mathcal{C}}HOM(P,L)_0 \to {}^{\mathcal{C}}HOM(P,M)_0 \to {}^{\mathcal{C}}HOM(P,N)_0 \to 0$ is exact. This means

that P is a projective object in ${}^{gr-\mathcal{C}}\mathcal{M}$. Assume that P is projective in ${}^{gr-\mathcal{C}}\mathcal{M}$. Let $0 \to L \to M \to N \to 0$ be an exact sequence in ${}^{gr-\mathcal{C}}\mathcal{M}$. Clearly, $0 \to L(x) \to M(x) \to N(x) \to 0$ is an exact sequence in ${}^{gr-\mathcal{C}}\mathcal{M}$ for every $x \in G$. By the projectivity of P, the sequence

$$0 \to {}^{\mathcal{C}}HOM(P, L(x))_0 \to {}^{\mathcal{C}}HOM(P, M(x))_0 \to {}^{\mathcal{C}}HOM(P, N(x))_0 \to 0$$

is exact for every $x \in G$. Using Lemma 1.4, we get that the sequence $0 \to {}^{\mathcal{C}}HOM(P, L) \to {}^{\mathcal{C}}HOM(P, M) \to {}^{\mathcal{C}}HOM(P, N) \to 0$ is exact. □

Let us consider A as a graded right A-module. By [13], $HOM_A(\mathcal{C}_A, A_A) = \oplus_x HOM_A(\mathcal{C}_A, A_A)_x$ is a graded vector space: it is the largest graded vector space contained in $Hom_A(\mathcal{C}_A, A_A)$. We write $\mathcal{C}^*_x = HOM_A(\mathcal{C}_A, A_A)_x$ and $\mathcal{C}^* = HOM_A(\mathcal{C}_A, A_A)$. Then $\mathcal{C}^*$ is a colour algebra called the graded right dual ring of $\mathcal{C}$ (see [5, 17.8] for the ungraded case): the multiplication is defined by $f\#g = (|f|, |g|)g \circ (f \otimes_A id_{\mathcal{C}}) \circ \Delta_{\mathcal{C}}$; i.e., $f\#g(c) = \sum_{|c|}(|f|, |g|)g(f(c_1)c_2)$ for all colour right A-linear maps f, g: $\mathcal{C} \to A$ and homogeneous element $c \in \mathcal{C}$; where $\Delta_{\mathcal{C}}(c) = \sum_{|c|} c_1 \otimes_A c_2$. The unit of $\mathcal{C}^*$ is $\epsilon_{\mathcal{C}}$ and there is a morphism of colour algebras $i : A^{op} \to \mathcal{C}^*$ defined by $i(a)(c) = (|a|, |c|)\epsilon_{\mathcal{C}}(c)a$. We will denote by $\mathcal{M}_{gr-\mathcal{C}^*}$ the category of graded right $\mathcal{C}^*$-modules. Any graded left $\mathcal{C}$-comodule M is a graded right $\mathcal{C}^*$-module: the action is defined by $m.f = \sum_{|m|}(|m_{(-1)}|, |f|)f(m_{(-1)})m_{(0)}$. If $\mathcal{C}$ is projective as a right A-module, then ${}^{gr-\mathcal{C}}\mathcal{M}$ is a full subcategory of $\mathcal{M}_{gr-\mathcal{C}^*}$; i.e., ${}^{\mathcal{C}}HOM(M, N) = HOM_{\mathcal{C}^*}(M, N)$ for any $M, N \in {}^{gr-\mathcal{C}}\mathcal{M}$. As a consequence, an object of ${}^{gr-\mathcal{C}}\mathcal{M}$ that is projective in $\mathcal{M}_{gr-\mathcal{C}^*}$ is projective in ${}^{gr-\mathcal{C}}\mathcal{M}$. Another consequence is that if M and N are objects of ${}^{gr-\mathcal{C}}\mathcal{M}$ with M finitely generated as a right $\mathcal{C}^*$-module, then ${}^{\mathcal{C}}HOM(M, N) = HOM_{\mathcal{C}^*}(M, N) = Hom_{\mathcal{C}^*}(M, N)$.
Given two graded left $\mathcal{C}$-comodules Λ and N, the graded vector space ${}^{\mathcal{C}}HOM(\Lambda, N)$ is a graded left module over the colour endomorphism ring $B = {}^{\mathcal{C}}END(\Lambda)$ of Λ: the action is given by $bf = (|b|, |f|)(f \circ b); \forall f \in {}^{\mathcal{C}}HOM(\Lambda, N), b \in B$. This defines a functor $F' = {}^{\mathcal{C}}HOM(\Lambda, -) : {}^{gr-\mathcal{C}}\mathcal{M} \to {}_{gr-B}\mathcal{M}$. Let us consider Λ as a graded right B-module by $\lambda.b = (|\lambda|/|b|)b(\lambda)$. So Λ is a graded (A, B)-bimodule. For any $P \in {}_{gr-B}\mathcal{M}$, $\Lambda \otimes_B P$ is a graded left $\mathcal{C}$-comodule with the coaction $\rho_{P,\mathcal{C}} = \rho_{\Lambda,\mathcal{C}} \otimes_B id_P$.

Lemma 1.7. *Let Λ and N be graded left $\mathcal{C}$-comodules and P be a graded left B-module. For every $x \in G$, the canonical linear map*

$$\phi : {}^{\mathcal{C}}HOM(\Lambda \otimes_B P, N)_x \to {}_BHOM(P, {}^{\mathcal{C}}HOM(\Lambda, N))_x$$

defined by $\phi(f)(p)(\lambda) = (|p|, |\lambda|)f(\lambda \otimes_B p)$ is an isomorphism.

Proof. The inverse of ϕ is defined by $\psi(g)(\lambda \otimes_B p) = (|\lambda|, |p|)g(p)(\lambda)$. □

We deduce from Lemma 1.7 that ${}^{\mathcal{C}}HOM(\Lambda \otimes_B P, N)_0 \simeq {}_BHOM(P, {}^{\mathcal{C}}HOM(\Lambda, N))_0$, and this means that the functor F' has the left adjoint $F = \Lambda \otimes_B - :_{gr-B} \mathcal{M} \to^{gr-\mathcal{C}} \mathcal{M}$. The unit of the adjunction is given by the graded k-linear map

$$u_N : N \to {}^{\mathcal{C}}HOM(\Lambda, \Lambda \otimes_B N), n \mapsto [\lambda \mapsto (|n|, |\lambda|)(\lambda \otimes n)]$$

for $N \in {}_{gr-B}\mathcal{M}$, while the counit is given by the graded k-linear map (the evaluation map)

$$c_M : \Lambda \otimes_B {}^{\mathcal{C}}HOM(\Lambda, M) \to M; \lambda \otimes f \mapsto (|\lambda|, |f|)f(\lambda)$$

for $M \in {}^{gr-\mathcal{C}}\mathcal{M}$. The adjointness property means that we have

$$F'(c_M) \circ u_{F'(M)} = id_{F'(M)},\ c_{F(N)} \circ F(u_N) = id_{F(N)};\ M \in {}^{gr-\mathcal{C}}\mathcal{M},\ N \in {}_{gr-B}\mathcal{M}.\ (\star)$$

2. The main results

Let A be a colour algebra and $\mathcal{C}$ a graded A-coring. We keep the notations of the preceding sections.

Lemma 2.1. *Let Λ and N be graded left $\mathcal{C}$-comodules. Set $B = {}^{\mathcal{C}}END(\Lambda)$. For every $x \in G$, we have*

(1) ${}^{\mathcal{C}}HOM(\Lambda, N(x)) = {}^{\mathcal{C}}HOM(\Lambda, N)(x)$

(2) $\Lambda \otimes_B B(x) = \Lambda(x)$.

An object $\Lambda \in {}^{gr-\mathcal{C}}\mathcal{M}$ is called semi-quasiprojective if the functor ${}^{\mathcal{C}}HOM(\Lambda, -) : {}^{gr-\mathcal{C}}\mathcal{M} \to {}_{gr-k}\mathcal{M}$ sends an exact sequence of the form $\oplus_I \Lambda(x_i) \to \oplus_J \Lambda(x_j) \to N \to 0$ to an exact sequence (see [15]). A projective object in ${}^{gr-\mathcal{C}}\mathcal{M}$ is semi-quasiprojective in ${}^{gr-\mathcal{C}}\mathcal{M}$.

Lemma 2.2. *Assume that $\mathcal{C}$ is projective as a right A-module. Let Λ be a graded left $\mathcal{C}$-comodule and set $B = {}^{\mathcal{C}}END(\Lambda)$. Then the functor ${}^{\mathcal{C}}HOM(\Lambda, -)$ commutes with*

(1) *direct sums if Λ is finitely generated as a graded right $\mathcal{C}^*$-module,*

(2) *direct limits if Λ is finitely presented as a graded right $\mathcal{C}^*$-module.*

Proof. (2) We know that ${}^{gr-\mathcal{C}}\mathcal{M}$ is a Grothendieck category so the functor ${}^{\mathcal{C}}HOM(\Lambda, -)_0$ preserves direct limits. We also know from Lemma 1.4 that $HOM_{\mathcal{C}^*}(\Lambda, N)_x = HOM_{\mathcal{C}^*}(\Lambda, N(x))_0$ for every $x \in G$. Let $(N_i)_{i \in I}$ be a directed system of right graded $\mathcal{C}^*$-modules. It is easy to show that $(\varinjlim N_i)(x) = \varinjlim(N_i(x))$ for every $x \in G$. Now the result follows from the fact direct limit commutes with direct sum. □

Lemma 2.3. *Assume that $\mathcal{C}$ is projective as a right A-module. Let Λ be a graded left $\mathcal{C}$-comodule that is finitely generated as a graded right $\mathcal{C}^*$-module, and let $B = {}^{\mathcal{C}}END(\Lambda)$. For every index set I,*

(1) *$c_{\oplus_I \Lambda(x_i)}$ is an isomorphism for every $x_i \in G$;*

(2) *$u_{\oplus_I B(x_i)}$ is an isomorphism for every $x_i \in G$;*

(3) *if Λ is semi-quasiprojective in ${}^{gr-\mathcal{C}}\mathcal{M}$, then u is a natural isomorphism; in other words, the induction functor $F = \Lambda \otimes_B (-)$ is fully faithful.*

Proof. (1) By Lemma 2.1(1), ${}^{\mathcal{C}}HOM(\Lambda,\Lambda)(x_i) = {}^{\mathcal{C}}HOM(\Lambda,\Lambda(x_i))$ for every $i \in I$. This implies that $\oplus_I B(x_i) = \oplus_I {}^{\mathcal{C}}HOM(\Lambda,\Lambda(x_i))$. By Lemma 2.2(1), the natural map $\kappa : \oplus_I B(x_i) \to {}^{\mathcal{C}}HOM(\Lambda, \oplus_I \Lambda(x_i))$ is an isomorphism. Lemma 2.1(2) implies that $\Lambda \otimes_B (\oplus_I B(x_i)) \simeq \oplus_I \Lambda(x_i)$. It is easy to see that this isomorphism is just $c_{\oplus_I \Lambda(x_i)} \circ (id_\Lambda \otimes \kappa)$. So $c_{\oplus_I \Lambda(x_i)}$ is an isomorphism since κ is an isomorphism.

(2) Putting $M = \oplus_I \Lambda(x_i)$ in $(\star)$ and using (1), we find

$$ {}^{\mathcal{C}}HOM(\Lambda, c_{\oplus_I \Lambda(x_i)}) \circ uc_{HOM(\Lambda, \oplus_I \Lambda(x_i))} = idc_{HOM(\Lambda, \oplus_I \Lambda(x_i))}; i.e.,$$

$$ {}^{\mathcal{C}}HOM(\Lambda, c_{\oplus_I \Lambda(x_i)}) \circ u_{\oplus_I B(x_i)} = id_{\oplus_I B(x_i)}.$$

From (1), ${}^{\mathcal{C}}HOM(\Lambda, c_{\oplus_I \Lambda(x_i)})$ is an isomorphism, hence $u_{\oplus_I B(x_i)}$ is an isomorphism.

(3) Take a graded free resolution $\oplus_J B(x_j) \to \oplus_I B(x_i) \to N \to 0$ of a graded left B-module N. Since u is natural, we have a commutative diagram

$$\begin{array}{ccccccc}
\oplus_{\mathbf{J}} \mathbf{B}(\mathbf{x_j}) & \longrightarrow & \oplus_{\mathbf{I}} \mathbf{B}(\mathbf{x_i}) & \longrightarrow & \mathbf{N} & \longrightarrow & 0 \\
\Big\downarrow{\scriptstyle u_{\oplus_J B(x_j)}} & & \Big\downarrow{\scriptstyle u_{\oplus_I B(x_i)}} & & \Big\downarrow{\scriptstyle u_N} & & \\
\mathbf{F'F}(\oplus_{\mathbf{J}} \mathbf{B}(\mathbf{x_j})) & \longrightarrow & \mathbf{F'F}(\oplus_{\mathbf{I}} \mathbf{B}(\mathbf{x_i})) & \longrightarrow & \mathbf{F'F}(\mathbf{N}) & \longrightarrow & 0\,.
\end{array}$$

The top row is exact; the bottom row is exact, since

$$F'F(\oplus_I B(x_i)) = {}^{\mathcal{C}}HOM(\Lambda, \Lambda \otimes_B \oplus_I B(x_i)) = {}^{\mathcal{C}}HOM(\Lambda, \oplus_I \Lambda(x_i))$$

and Λ is semi-quasiprojective. By (2), $u_{\oplus_I B(x_i)}$ and $u_{\oplus_J B(x_j)}$ are isomorphisms; and it follows from the five lemma that u_N is an isomorphism. □

We can now give equivalent conditions for the projectivity and flatness of $P \in {}_{gr-B}\mathcal{M}$.

Theorem 2.4. *Assume that $\mathcal{C}$ is projective as a right A-module. Let Λ be a graded left $\mathcal{C}$-comodule that is finitely generated as a graded right $\mathcal{C}^*$-module, and let $B = {}^{\mathcal{C}}END(\Lambda)$. For $P \in {}_{gr-B}\mathcal{M}$, we consider the following statements.*

(1) *$\Lambda \otimes_B P$ is projective in ${}^{gr-\mathcal{C}}\mathcal{M}$ and u_P is injective;*

(2) *P is projective as a graded left B-module;*

(3) *$\Lambda \otimes_B P$ is a direct summand in ${}^{gr-\mathcal{C}}\mathcal{M}$ of some $\oplus_I \Lambda(x_i)$, and u_P is bijective;*

(4) *there exists $Q \in {}^{gr-\mathcal{C}}\mathcal{M}$ such that Q is a direct summand of some $\oplus_I \Lambda(x_i)$, and $P \cong {}^{\mathcal{C}}HOM(\Lambda, Q)$ in ${}_{gr-B}\mathcal{M}$;*

(5) *$\Lambda \otimes_B P$ is a direct summand in ${}^{gr-\mathcal{C}}\mathcal{M}$ of some $\oplus_I \Lambda(x_i)$.*

Then (1) $\Rightarrow$ (2) $\Leftrightarrow$ (3) $\Leftrightarrow$ (4) $\Rightarrow$ (5).

If Λ *is semi-quasiprojective in* ${}^{gr-\mathcal{C}}\mathcal{M}$, *then* (5) $\Rightarrow$ (3)*; if* Λ *is projective in* ${}^{gr-\mathcal{C}}\mathcal{M}$, *then* (3) $\Rightarrow$ (1).

Proof. (2) $\Rightarrow$ (3): If P is projective as a graded left B-module, then we can find an index set I and $P' \in {}_{gr-B}\mathcal{M}$ such that $\oplus_I B(x_i) \cong P \oplus P'$. Then obviously

$$\oplus_I \Lambda(x_i) \cong \Lambda \otimes_B (\oplus_I B(x_i)) \cong (\Lambda \otimes_B P) \oplus (\Lambda \otimes_B P').$$

Since u is a natural transformation, we have a commutative diagram:

$$\begin{array}{ccc}
\oplus_{\mathbf{I}}\mathbf{B}(\mathbf{x_i}) & \xrightarrow{\cong} & \mathbf{P} \oplus \mathbf{P'} \\
\Big\downarrow{\scriptstyle u_{\oplus_I B(x_i)}} & & \Big\downarrow{\scriptstyle u_P \oplus u_{P'}} \\
{}^{\mathcal{C}}\mathbf{HOM}(\mathbf{\Lambda}, \oplus_{\mathbf{I}}\mathbf{\Lambda}(\mathbf{x_i})) & \xrightarrow[\cong]{} & {}^{\mathcal{C}}\mathbf{HOM}(\mathbf{\Lambda}, \mathbf{\Lambda} \otimes_{\mathbf{B}} \mathbf{P}) \oplus {}^{\mathcal{C}}\mathbf{HOM}(\mathbf{\Lambda}, \mathbf{\Lambda} \otimes_{\mathbf{B}} \mathbf{P'})
\end{array}$$

From the fact that $u_{\oplus_I B(x_i)}$ is an isomorphism, it follows that u_P (and $u_{P'}$) are isomorphisms.

(3) $\Rightarrow$ (4): Take $Q = \Lambda \otimes_B P$.

(4) $\Rightarrow$ (2): Let $f : \oplus_I \Lambda(x_i) \to Q$ be a split epimorphism in ${}^{gr-\mathcal{C}}\mathcal{M}$. Then

$${}^{\mathcal{C}}HOM(\Lambda, f) : {}^{\mathcal{C}}HOM(\Lambda, \oplus_I \Lambda(x_i)) \cong \oplus_I B(x_i) \to {}^{\mathcal{C}}HOM(\Lambda, Q) \cong P$$

is also split surjective, hence P is projective as a graded left B-module.

(4) $\Rightarrow$ (5): If (4) is true, we know from the proof of (4) $\Rightarrow$ (2) that P is a direct summand of some $\oplus_I B(x_i)$ in ${}_{gr-B}\mathcal{M}$. So $\Lambda \otimes_B P$ is a direct summand of $\oplus_I \Lambda(x_i)$.

(1) $\Rightarrow$ (2): Take an epimorphism $f : \oplus_I B(x_i) \to P$ in ${}_{gr-B}\mathcal{M}$. Then

$$F(f) = id_\Lambda \otimes_B f : \Lambda \otimes_B (\oplus_I B(x_i)) \cong \oplus_I \Lambda(x_i) \to \Lambda \otimes_B P$$

is an epimorphism in ${}^{gr-\mathcal{C}}\mathcal{M}$, and it splits since $\Lambda \otimes_B P$ is projective in ${}^{gr-\mathcal{C}}\mathcal{M}$. Consider the commutative diagram:

$$\begin{array}{ccccc}
\oplus_{\mathbf{I}}\mathbf{B}(\mathbf{x_i}) & \xrightarrow{f} & \mathbf{P} & \longrightarrow & 0 \\
\Big\downarrow{\scriptstyle u_{\oplus_I B(x_i)}} & & \Big\downarrow{\scriptstyle u_P} & & \\
{}^{\mathcal{C}}\mathbf{HOM}(\mathbf{\Lambda}, \oplus_{\mathbf{I}}\mathbf{\Lambda}(\mathbf{x_i})) & \xrightarrow[F'F(f)]{} & {}^{\mathcal{C}}\mathbf{HOM}(\mathbf{\Lambda}, \mathbf{\Lambda} \otimes_{\mathbf{B}} \mathbf{P}) & \longrightarrow & 0\,.
\end{array}$$

The bottom row is split exact, since any functor, in particular, ${}^{\mathcal{C}}HOM(\Lambda,-)$ preserves split exact sequences. By Lemma 2.3(2), $u_{\oplus_I B(x_i)}$ is an isomorphism. A diagram chasing tells us that u_P is surjective. By assumption, u_P is injective, so u_P is bijective. We deduce that the top row is isomorphic to the bottom row, and therefore splits. Thus P is projective in $\in {}_{gr-B}\mathcal{M}$.

Under the assumption that Λ is semi-quasiprojective in ${}^{gr-\mathcal{C}}\mathcal{M}$, (5) $\Rightarrow$ (3) follows from Lemma 2.3(3).

(3) $\Rightarrow$ (1): By (3), $\Lambda\otimes_B P$ is a direct summand of some $\oplus_I\Lambda(x_i)$. If Λ is projective in ${}^{gr-\mathcal{C}}\mathcal{M}$, then $\oplus_I\Lambda(x_i)$ is projective in ${}^{gr-\mathcal{C}}\mathcal{M}$. So $\Lambda\otimes_B P$ being a direct summand of a projective object of ${}^{gr-\mathcal{C}}\mathcal{M}$ is projective in ${}^{gr-\mathcal{C}}\mathcal{M}$. □

Theorem 2.5. *Assume that $\mathcal{C}$ is projective as a right A-module. Let Λ be a graded left $\mathcal{C}$-comodule that is finitely presented as a graded right $\mathcal{C}^*$-module, and let $B = {}^{\mathcal{C}}END(\Lambda)$. For $P \in {}_{gr-B}\mathcal{M}$, the following assertions are equivalent.*

(1) *P is flat as a graded left B-module;*

(2) *$\Lambda \otimes_B P = \varinjlim Q_i$, where $Q_i \cong \oplus_{j\le n_i} B(x_{ij})$ in ${}^{gr-\mathcal{C}}\mathcal{M}$ for some positive integer n_i, and u_P is bijective;*

(3) *$\Lambda \otimes_B P = \varinjlim Q_i$, where $Q_i \in {}^{gr-\mathcal{C}}\mathcal{M}$ is a direct summand of some $\oplus_{j\in I_i}\Lambda(x_{ij})$ in ${}^{gr-\mathcal{C}}\mathcal{M}$, and u_P is bijective;*

(4) *there exists $Q = \varinjlim Q_i \in {}^{gr-\mathcal{C}}\mathcal{M}$, such that $Q_i \cong \oplus_{j\le n_i}\Lambda(x_{ij})$ for some positive integer n_i and ${}^{\mathcal{C}}HOM(\Lambda, Q) \cong P$ in ${}_{gr-B}\mathcal{M}$;*

(5) *there exists $Q = \varinjlim Q_i \in {}^{gr-\mathcal{C}}\mathcal{M}$, such that Q_i is a direct summand of some $\oplus_{j\in I_i}\Lambda(x_{ij})$ in ${}^{gr-\mathcal{C}}\mathcal{M}$, and ${}^{\mathcal{C}}HOM(\Lambda, Q) \cong P$ in ${}_{gr-B}\mathcal{M}$.*

If Λ is semi-quasiprojective in ${}^{gr-\mathcal{C}}\mathcal{M}$, these conditions are also equivalent to conditions (2) *and* (3)*, without the assumption that u_P is bijective.*

Proof. (1) $\Rightarrow$ (2): $P = \varinjlim N_i$, with $N_i = \oplus_{j\le n_i} B(x_{ij})$ for some positive integer n_i. Take $Q_i = \oplus_{j\le n_i}\Lambda(x_{ij})$, then

$$\varinjlim Q_i \cong \varinjlim(\Lambda \otimes_B N_i) \cong \Lambda \otimes_B \varinjlim N_i \cong \Lambda \otimes_B P.$$

Consider the following commutative diagram:

$$\begin{array}{ccc}
\mathbf{P} = \varinjlim \mathbf{N_i} & \xrightarrow{\lim(u_{N_i})} & \varinjlim {}^{\mathcal{C}}\mathbf{HOM}(\mathbf{\Lambda}, \mathbf{\Lambda} \otimes_{\mathbf{B}} \mathbf{N_i}) \\
\Big\downarrow u_P & & \Big\downarrow f \\
{}^{\mathcal{C}}\mathbf{HOM}(\mathbf{\Lambda}, \mathbf{\Lambda} \otimes_{\mathbf{B}} (\varinjlim \mathbf{N_i})) & \xrightarrow[\cong]{} & {}^{\mathcal{C}}\mathbf{HOM}(\mathbf{\Lambda}, \varinjlim(\mathbf{\Lambda} \otimes_{\mathbf{B}} \mathbf{N_i}))\,.
\end{array}$$

By Lemma 2.3(2), the u_{N_i} are isomorphisms; by Lemma 2.2, the natural homomorphism f is an isomorphism. Hence u_P is an isomorphism.

(2) $\Rightarrow$ (3) and (4) $\Rightarrow$ (5) are obvious.
(2) $\Rightarrow$ (4) and (3) $\Rightarrow$ (5): Put $Q = \Lambda \otimes_B P$. Then $u_P : P \to {}^{\mathcal{C}}HOM(\Lambda, \Lambda \otimes_B P)$ is the required isomorphism.
(5) $\Rightarrow$ (1): We have a split exact sequence $0 \to N_i \to P_i = \oplus_{j\in I_i}\Lambda(x_{ij}) \to Q_i \to 0$ in ${}^{gr-\mathcal{C}}\mathcal{M}$. Consider the following commutative diagram:

$$\begin{array}{ccccccccc}
\mathbf{0} & \longrightarrow & \mathbf{FF'(N_i)} & \longrightarrow & \mathbf{FF'(P_i)} & \longrightarrow & \mathbf{FF'(Q_i)} & \longrightarrow & \mathbf{0} \\
 & & \downarrow{\scriptstyle c_{N_i}} & & \downarrow{\scriptstyle c_{P_i}} & & \downarrow{\scriptstyle c_{Q_i}} & & \\
\mathbf{0} & \longrightarrow & \mathbf{N_i} & \longrightarrow & \mathbf{P_i} & \longrightarrow & \mathbf{Q_i} & \longrightarrow & \mathbf{0}\,.
\end{array}$$

We know from Lemma 2.3(1) that c_{P_i} is an isomorphism. Both rows in the diagram are split exact, so it follows that c_{N_i} and c_{Q_i} are also isomorphisms. Next consider the commutative diagram:

$$\begin{array}{ccc}
\mathbf{\Lambda} \otimes_{\mathbf{B}} \varinjlim {}^{\mathcal{C}}\mathbf{HOM}(\mathbf{\Lambda}, \mathbf{Q_i}) & \xrightarrow{id_\Lambda \otimes f} & \mathbf{\Lambda} \otimes_{\mathbf{B}} {}^{\mathcal{C}}\mathbf{HOM}(\mathbf{\Lambda}, \mathbf{Q}) \\
\uparrow{\scriptstyle h} & & \downarrow{\scriptstyle c_Q} \\
\varinjlim(\mathbf{\Lambda} \otimes_{\mathbf{B}} {}^{\mathcal{C}}\mathbf{HOM}(\mathbf{\Lambda}, \mathbf{Q_i})) & \xrightarrow[\lim c_{Q_i}]{} & \mathbf{Q}
\end{array}$$

where h and f are the natural homomorphisms. h is an isomorphism, because $\Lambda \otimes_B (-)$ preserves inductive limits; by Lemma 2.2, f is an isomorphism; and $limc_{Q_i}$ is an isomorphism because every c_{Q_i} is an isomorphism. It follows that c_Q is an isomorphism, hence ${}^{\mathcal{C}}HOM(\Lambda, c_Q)$ is an isomorphism. From $(\star)$, we get

$$ {}^{\mathcal{C}}HOM(\Lambda, c_Q) \circ uc_{HOM(\Lambda,Q)} = idc_{HOM(\Lambda,Q)}.$$

It follows that $uc_{HOM(\Lambda,Q)}$ is also an isomorphism. Since ${}^{\mathcal{C}}HOM(\Lambda, Q) \cong P$, u_P is an isomorphism. Consider the isomorphisms

$$P \cong {}^{\mathcal{C}}HOM(\Lambda, \Lambda \otimes_B P) \cong {}^{\mathcal{C}}HOM(\Lambda, \Lambda \otimes_B {}^{\mathcal{C}}HOM(\Lambda, Q)) \cong$$
$$ {}^{\mathcal{C}}HOM(\Lambda, Q) \cong \varinjlim {}^{\mathcal{C}}HOM(\Lambda, Q_i);$$

where the first isomorphism is u_P, the third is ${}^{\mathcal{C}}HOM(\Lambda, c_Q)$ and the last one is f. It follows from Lemmas 2.1(1)and 2.2 that ${}^{\mathcal{C}}HOM(\Lambda, P_i) \cong \oplus_{j\in I_i} B(x_{ij})$ is projective as a graded left B-module, hence ${}^{\mathcal{C}}HOM(\Lambda, Q_i)$ is also projective as a graded left B-module, and we conclude that P is flat in $\in {}_{gr-B}\mathcal{M}$. The final statement is an immediate consequence of Lemma 2.3(3). $\square$

3. Applications

3.1. $\mathcal{C}$ contains a grouplike element

A grouplike element of $\mathcal{C}$ is an element $X \in \mathcal{C}_0$ such that $\Delta_c(X) = X \otimes_A X$ and $\epsilon_{\mathcal{C}}(X) = 1_A$ (see [14]). If $\mathcal{C}$ contains a grouplike element X, then A is an object of ${}^{gr-\mathcal{C}}\mathcal{M}$: the $\mathcal{C}$-coaction is defined by $\rho_{A,\mathcal{C}}(a) = aX = aX \otimes_A 1_A; \forall a \in A$. Conversely, if A is an object of ${}^{gr-\mathcal{C}}\mathcal{M}$, then $\rho_{A,\mathcal{C}}(1_A) = X$ is a grouplike element of $\mathcal{C}$.

Assume that $\mathcal{C}$ contains a grouplike element X. Then A is an object of $\mathcal{M}_{gr-\mathcal{C}^*}$ and $a.\epsilon_{\mathcal{C}} = a$, that is, A is generated as a right $\mathcal{C}^*$-module by the homogeneous element $\epsilon_{\mathcal{C}}$ of degree 0. Lemma 1.2 implies that A is finitely generated in $\mathcal{M}_{gr-\mathcal{C}^*}$. For any graded left $\mathcal{C}$-comodule M, we call ${}^{co\mathcal{C},X}M = \{m \in M, \quad \rho_{M,\mathcal{C}}(m) = X \otimes_A m\}$ the vector space of $(\mathcal{C}, X)$-coinvariants of M. Clearly, ${}^{co\mathcal{C},X}A = \{a \in A, \quad Xa = aX\}$ is a colour subalgebra of A: the colour subalgebra of $(\mathcal{C}, X)$-coinvariants. For every $f \in {}^{\mathcal{C}}HOM(A, M)$, $f(1) \in {}^{co\mathcal{C},X}M$. The graded k-linear map $f \mapsto f(1)$ establishes an isomorphism ${}^{\mathcal{C}}HOM(A, M) \to {}^{co\mathcal{C},X}M$ with inverse the graded k-linear map ψ defined by $\psi(m)(a) = (|m|, |a|)am$. We have ${}^{\mathcal{C}}END(A) = {}^{co\mathcal{C},X}A$. Set $B = {}^{co\mathcal{C},X}A$. Then we get from Theorems 2.4 and 2.5 necessary and sufficient conditions for projectivity and flatness over the colour algebra $B = {}^{co\mathcal{C},X}A$.

3.2. A colour algebra as a trivial coring

A colour algebra A is a graded A-bimodule. Let us define $\Delta_A(a) = a \otimes_A 1_A$ and $\epsilon_A(a) = a$. Then A is an A-coring. A graded left A-comodule is just a graded left A-module. The product on A^* is defined by $f\#g(a) = \sum_{|a|}(|f|, |g|)g(f(a)1_A) = \sum_{|a|}(|f|, |g|)g(1_A)(f(a))$. It is easy to show that the algebra A^* is isomorphic to A^{op}, the opposite algebra of A: this isomorphism is defined by $f \mapsto f(1_A)$. For graded left A-modules M and N, we have ${}^{A}HOM(M, N) = {}_{A}HOM(M, N)$. Then Theorems 2.4 and 2.5 give necessary and sufficient conditions for projectivity and flatness over $B =_A END(\Lambda)$, where Λ is a finitely generated graded left A-module. When the gradation is trivial we recover [11]. In many examples, Λ will be a colour algebra and A will be a graded Λ-ring with a graded left grouplike character.

Definition 3.1. (see [6], Section 2) *Let A and Λ be two colour algebras and $i : \Lambda \to A$ a graded ring morphism. A graded k-linear map $\chi : A \to \Lambda$ is called a graded left grouplike character on A if χ is graded left Λ-linear and*

$$\chi(a\chi(a')) = \chi(aa') \quad \textit{and} \quad \chi(1_A) = 1_\Lambda \quad \forall \quad a, a' \in A.$$

We then say that A is a graded Λ-ring with a graded left grouplike character χ.

Let A be a graded Λ-ring with a graded left grouplike character χ. Then Λ is a graded left A-module: the action is given by $a \rightharpoonup \lambda = \chi(a\lambda)$. Furthermore, Λ is cyclic as a left A-module, since $\lambda = (\lambda 1_A) \rightharpoonup 1_\Lambda$. But 1_Λ is homogeneous of degree 0, so Λ is a finitely generated as a graded left A-module (Lemma 1.2). So

we get necessary and sufficient conditions for projectivity and flatness over the colour endomorphism ring ${}_AEND(\Lambda)$ of Λ.

Now we will give two examples of this situation. There are other examples in the literature.

• Let H be a colour algebra, the colour tensor product $H \otimes H$ is the G-graded vector space $H \otimes H = \oplus_{x\in G}(\oplus_{y+z=x} H_y \otimes H_z)$ with multiplication $(h \otimes l)(h' \otimes l') = (|l|/|h'|)hh' \otimes ll'$ for homogeneous elements h, h', l, $l' \in H$. By [9, Lemma 3.2], $H \otimes H$ is a colour algebra. A Hopf colour algebra is a colour algebra and a graded coalgebra such that Δ_H and ϵ_H are morphisms of colour algebras and there exists a graded k-linear map $S_H : H \to H$ (called antipode) such that $(S_H \otimes id_H) \circ \Delta_H = \epsilon_H = (id_H \otimes S_H) \circ \Delta_H$ or equivalently, $\sum_{|h|} \epsilon(h_1)h_2 = h = \sum_{|h|} h_1\epsilon(h_2)$ and $\sum_{|h|} S(h_1)h_2 = \epsilon(h) = \sum_{|h|} h_1 S(h_2)$.

Let H be a Hopf colour algebra over k with comultiplication Δ_H, counit ϵ_H and antipode S_H. A colour algebra Λ which is a graded left H-module such that $h.(\lambda\lambda') = \sum_{|h|}(|h_2|/|\lambda|)(h_1.\lambda)(h_2.\lambda')$ for all $h \in H$ and $\lambda, \lambda' \in \Lambda$ will be called a graded left H-module algebra. We denote by $A = \Lambda\#H$ the associated smash product; i.e., the colour algebra generated by Λ and H whose multiplication is defined by $(\lambda h)(\lambda' h') = \sum_{|h|}(|h_2|/|\lambda'|)\lambda(h_1.\lambda')(h_2 h')$ (see [12]). A graded vector space M is a graded left A-module if and only if it is a graded left Λ-module and a graded left H-module such that $h.(\lambda m) = \sum_{|h|}(|h_2|/|\lambda|)(h_1.\lambda)(h_2 m)$. Define a k-linear map $\chi : A \to \Lambda$ by $\chi(\lambda h) = \epsilon_H(h)\lambda$. Since $\epsilon_H(h) = 0$ for $|h| \neq 0$, χ is homogeneous of degree 0. Clearly, χ is left Λ-linear. It follows that $\Lambda\#H$ is a graded Λ-ring with a graded left grouplike character χ. Note that ${}_{\Lambda\#H}END(\Lambda)$ is exactly the colour subring of invariants of Λ; i.e., ${}_{\Lambda\#H}END(\Lambda) = \{\lambda \in \Lambda; h.\lambda = \epsilon_H(h)\lambda\}$.

• Assume that $\mathcal{C}$ contains a grouplike element X. The linear map $i : A \to \mathcal{C}^*$ defined by $i(a)(c) = a\epsilon_C(c)$ is a morphism of colour algebras. Define $\chi : \mathcal{C}^* \to A$ by $\chi(f) = f(X)$. Then χ is a graded left grouplike character on $\mathcal{C}^*$. So $\mathcal{C}^*$ is a graded A-ring.

3.3. $\mathcal{C}$ comes from a graded entwining structure

In this section, A is a colour algebra with multiplication μ and unit ι, and C is a graded coalgebra with comultiplication Δ_C and counit ϵ_C. We denote by $\tau : A\otimes C \to C\otimes A$ the twist map; that is $\tau(a\otimes c) = (|a|/|c|)c\otimes a$. If M is a left (resp. right C-comodule, we write $\rho_{M,C}(m) = m_{-1} \otimes m_0$ (resp. $\rho_{M,C}(m) = m_0 \otimes m_1$). We remind that C is a graded k-coring. Interesting examples of graded corings come from graded entwining structures. We will often refer to [5] for the ungraded case.

• Graded left-left entwined modules.

A graded left-left entwining structure over k is a triple (A, C, ψ) with a graded k-linear map $\psi : A\otimes C \to C\otimes A$; $a\otimes c \mapsto (|a_\alpha|/|c|)c^\alpha \otimes a_\alpha$ satisfying the following conditions [5, 32.1]:

$$\psi \circ (\mu \otimes id_C) = (id_C \otimes \mu) \circ (\psi \otimes id_A) \circ (id_A \otimes \psi)$$

$$(\Delta_C \otimes id_A) \circ \psi = (id_C \otimes \psi) \circ (\psi \otimes id_C) \circ (id_A \otimes \Delta_C)$$
$$(\epsilon_C \otimes id_A) \circ \psi = id_A \otimes \epsilon_C$$
$$\psi \circ (id_C \otimes \iota) = \iota \otimes id_C.$$

These relations are respectively equivalent to

$$(|(aa')_\alpha|/|c|)(c^\alpha \otimes (aa')_\alpha) = (|a'_\alpha|/|c|)(|a_\beta|/|c^\alpha|)(c^{\alpha\beta} \otimes a_\beta a'_\alpha)$$
$$(|a_\alpha|/|c|)(\Delta_C(c^\alpha) \otimes a_\alpha) = (|a_\alpha|/|c_1|)(|a_{\alpha\beta}|/|c_2|)({c_1}^\alpha \otimes {c_2}^\beta \otimes a_{\alpha\beta})$$
$$(|a_\alpha|/|c|)(\epsilon_C(c^\alpha)a_\alpha) = a\epsilon_C(c)$$
$$(|1_\alpha|/|c|)(c^\alpha \otimes 1_\alpha) = c \otimes 1.$$

The map ψ is called a graded entwining map, and A and C are said to be graded entwined by ψ. By [5, 32.1], $\mathcal{C} = C \otimes A$ is a graded A-coring with A-multiplications $a'(c \otimes a)a'' = \psi(a' \otimes c)aa''$, coproduct

$$\Delta_{\mathcal{C}} : C \otimes A \to C \otimes A \otimes_A C \otimes A \cong C \otimes C \otimes A; \quad c \otimes a \mapsto \Delta_C(c) \otimes a$$

and counit $\epsilon_{\mathcal{C}}(c \otimes a) = \epsilon_C(c)a$.

Let M be a graded left A-module. Then $C \otimes M$ becomes a graded left A-module if we set $a(c \otimes m) = (|a_\alpha|/|c|)|)c^\alpha \otimes (a_\alpha m)$. We say that a vector space M is a graded left-left (A, C, ψ)-entwined module if M is a graded left A-module and a graded left C-comodule such that $\rho_{M,C}$ is a graded left A-linear map; i.e.,

$$\rho_{M,C}(am) = (|a_\alpha|/|(m_{-1})|)(m_{-1})^\alpha \otimes (a_\alpha m_0).$$

We denote by ${}^{gr-C}_{gr-A}\mathcal{M}(\psi)$ the category of graded left-left (A, C, ψ)-entwined modules: its morphisms are the graded left A-linear maps and the graded left C-colinear maps. We can show that ${}^{gr-C}_{gr-A}\mathcal{M}(\psi)$ is isomorphic to ${}^{gr-\mathcal{C}}\mathcal{M}$.

• Graded right-right entwined modules.

A graded right-right entwining structure over k is a triple (A, C, ψ) with a graded k-linear map $C \otimes A \to A \otimes C$; $c \otimes a \mapsto (|c|/|a_\alpha|)a_\alpha \otimes c^\alpha$ satisfying the following conditions [5, 32.1]:

$$\psi \circ (id_C \otimes \mu) = (\mu \otimes id_C) \circ (id_A \otimes \psi) \circ (\psi \otimes id_A)$$
$$(id_A \otimes \Delta_C) \circ \psi = (\psi \otimes id_C) \circ (id_C \otimes \psi) \circ (\Delta_C \otimes id_A)$$
$$(id_A \otimes \epsilon_C) \circ \psi = \epsilon_C \otimes id_A$$
$$\psi \circ (id_C \otimes \iota) = \iota \otimes id_C.$$

These relations are respectively equivalent to

$$(|c|/|(aa')_\alpha|)((aa')_\alpha \otimes c^\alpha) = (|c|/|a_\alpha|)(|c^\alpha|/|a'_\beta|)(a_\alpha a'_\beta \otimes c^{\alpha\beta})$$
$$(|c|/|a_\alpha|)(a_\alpha \otimes \Delta_C(c^\alpha)) = (|(c_2)|/|a_\alpha|)(|(c_1)|/|a_{\alpha\beta}|)(a_{\alpha\beta} \otimes {c_1}^\beta \otimes {c_2}^\alpha)$$
$$(|c|/|a_\alpha|)(a_\alpha \epsilon_C(c^\alpha)) = \epsilon_C(c)a$$

$$(|c|/|1_\alpha|)(1_\alpha \otimes c^\alpha) = 1 \otimes c.$$

By [5, 32.1], $\mathcal{C} = A \otimes C$ is a graded A-coring with A-multiplications $a'(a \otimes c)a'' = a'a\psi(c \otimes a'')$, coproduct

$$\Delta_{\mathcal{C}} : A \otimes C \to A \otimes C \otimes_A A \otimes C \cong A \otimes C \otimes C; \quad a \otimes c \mapsto a \otimes \Delta_C(c)$$

and counit $\epsilon_{\mathcal{C}}(a \otimes c) = a\epsilon_C(c)$.

Let M be a graded right A-module. Then $M \otimes C$ becomes a graded right A-module if we set $(m \otimes c)a = (|c|/|a_\alpha|)(ma_\alpha) \otimes c^\alpha$. A vector space M is a graded right-right (A, C, ψ)-entwined module if M is a graded right A-module and a graded right C-comodule via such that $\rho_{M,C}$ is a graded right A-linear map; i.e.,

$$\rho_{M,C}(ma) = (|(m_1)|/|a_\alpha|)(m_0 a_\alpha) \otimes (m_1)^\alpha.$$

We denote by $\mathcal{M}(\psi)^{gr-C}_{gr-A}$ the category of graded right-right (A, C, ψ)-entwined modules: its morphisms are the graded right A-linear maps and the graded right C-colinear maps. We can show that this category is isomorphic to $\mathcal{M}^{gr-\mathcal{C}}$.

• Graded left-right entwined modules.

A graded left-right entwining structure over k is a triple (A, C, ψ) with a graded k-linear map $A \otimes C \to A \otimes C$; $a \otimes c \mapsto a_\alpha \otimes c^\alpha$ satisfying the following conditions [5, 32.1]:

$$\psi \circ (\mu \otimes id_C) = (\mu \otimes id_C) \circ (id_A \otimes \tau^{-1}) \circ \circ(\psi \otimes id_A) \circ (id_A \otimes \tau) \circ (id_A \otimes \psi)$$

$$(id_A \otimes \Delta_C) \circ \psi = (\tau^{-1} \otimes id_C) \circ (id_C \otimes \psi) \circ (\tau \otimes id_C) \circ (\psi \otimes id_C) \circ (id_A \otimes \Delta_C)$$

$$(id_A \otimes \epsilon_C) \circ \psi = id_A \otimes \epsilon_C$$

$$\psi \circ (\iota \otimes id_C) = \iota \otimes id_C.$$

These relations are respectively equivalent to

$$(aa')_\alpha \otimes c^\alpha = (|a'_\alpha|/|c^\alpha|)(|c^{\alpha\beta}|/|a'_\alpha|)(a_\beta a'_\alpha \otimes c^{\alpha\beta})$$

$$a_\alpha \otimes \Delta_C(c^\alpha) = (|a_\alpha|/|{c_1}^\alpha|)(|{c_1}^\alpha|/|a_{\alpha\beta}|)(a_{\alpha\beta} \otimes {c_1}^\alpha \otimes {c_2}^\beta)$$

$$a_\alpha \epsilon_C(c^\alpha) = a\epsilon_C(c)$$

$$1_\alpha \otimes c^\alpha = 1 \otimes c.$$

Let M be a graded left A-module. Then $M \otimes C$ becomes a graded left A-module if we set $a(m \otimes c) = (|m|/|c|)(|c^\alpha|/|m|)(a_\alpha m \otimes c^\alpha)$. A vector space M is a graded left-right (A, C, ψ)-entwined module if M is a graded left A-module and a graded right C-comodule such that $\rho_{M,C}$ is a graded left A-linear map; i.e.,

$$\rho_{M,C}(am) = (|m_0|/|m_1|)(|(m_1)^\alpha|/|m_0|)(a_\alpha m_0 \otimes (m_1)^\alpha.$$

We denote by ${}_{gr-A}\mathcal{M}(\psi)^{gr-C}$ the category of graded left-right (A, C, ψ)-entwined modules: its morphisms are the graded left A-linear maps and the graded right C-colinear maps.

• Graded right-left entwined modules.

A graded right-left entwining structure over k is a triple (A, C, ψ) with a graded k-linear map $C \otimes A \to C \otimes A$; $c \otimes a \mapsto c^{\alpha} \otimes a_{\alpha}$ satisfying the following conditions [5, 32.1]:

$$\psi \circ (id_C \otimes \mu) = (id_C \otimes \mu) \circ (\tau \otimes id_A) \circ (id_A \otimes \psi) \circ (\tau^{-1} \otimes id_A) \circ (\psi \otimes id_A)$$

$$(\Delta_C \otimes id_A) \circ \psi = (id_C \otimes \tau) \circ (\psi \otimes id_C) \circ (id_C \otimes \tau^{-1}) \circ (id_C \otimes \psi) \circ (\Delta_C \otimes id_A)$$

$$(\epsilon_C \otimes id_A) \circ \psi = \epsilon_C \otimes id_A$$

$$\psi \circ (id_C \otimes \iota) = id_C \otimes \iota$$

where $\tau : C \otimes A \to A \otimes C$; $c \otimes a \mapsto (|c|/|a|)a \otimes c$. These relations are respectively equivalent to

$$c^{\alpha} \otimes (aa')_{\alpha} = (|c^{\alpha}|/|a_{\alpha}|)(|a_{\alpha}|/|c^{\alpha\beta}|)(c^{\alpha\beta} \otimes a_{\alpha}a'_{\beta})$$

$$\Delta_C(c^{\alpha}) \otimes a_{\alpha} = (|{c_2}^{\alpha}|/|a_{\alpha}|)(|a_{\alpha\beta}|/|{c_2}^{\alpha}|)({c_1}^{\beta} \otimes {c_2}^{\alpha} \otimes a_{\alpha\beta})$$

$$\epsilon_C(c^{\alpha})a_{\alpha} = \epsilon_C(c)a$$

$$c_{\alpha} \otimes 1^{\alpha} = c \otimes 1.$$

Let M be a graded right A-module. Then $C \otimes M$ becomes a graded right A-module if we set $(c \otimes m)a = (|c|/|m|)(|m|/|c^{\alpha}|)(c^{\alpha} \otimes ma_{\alpha})$. A vector space M is a graded right-left (A, C, ψ)-entwined module if M is a graded right A-module and a graded left C-comodule such that $\rho_{M,C}$ is a graded right A-linear map; i.e.,

$$\rho_{M,C}(ma) = (|m_{-1}|/|m_0|)(|m_0|/|(m_{-1})^{\alpha}|)(m_{-1})^{\alpha} \otimes (m_0 a_{\alpha}).$$

We denote by ${}^{gr-C}\mathcal{M}(\psi)_{gr-A}$ the category of graded right-left (A, C, ψ)-entwined modules: its morphisms are the graded right A-linear maps and the graded left C-colinear maps.

3.3.1. Graded Doi-Hopf modules

In this section, H is a Hopf colour algebra with a bijective antipode S_H, A is a colour algebra and C is a graded coalgebra.

We say that A is a graded left H-comodule algebra if it is a graded left H-comodule via $\rho_{A,H}(a) = a_{[-1]} \otimes a_{[0]}$ such that $\rho_{A,H}(aa') = (|a_{[0]}|/|a'_{[-1]}|)(a_{[-1]}a'_{[-1]}) \otimes a_{[0]} \otimes a'_{[0]}$ and $\rho_{A,H}(1_A) = 1_H \otimes 1_A$. This is equivalent to say that the multiplication and the unit are graded left H-colinear, where the left H-coaction on $A \otimes A$ is defined by $(a \otimes a')_{[-1]} \otimes (a \otimes a')_{[0]} = (|a_{[0]}|/|a'_{[-1]}|)(a_{[-1]}a'_{[-1]}) \otimes a_{[0]} \otimes a'_{[0]}$.

We say that A is a graded right H-comodule algebra if it is a graded right H-comodule via $\rho_{A,H}(a) = a_{[0]} \otimes a_{[1]}$ such that $\rho_{A,H}(aa') = (|a_{[1]}|/|a'_{[0]}|)a_{[0]} \otimes a'_{[0]} \otimes (a_{[1]}a'_{[1]})$ and $\rho_{A,H}(1_A) = 1_A \otimes 1_H$. This is equivalent to say that the multiplication and the unit are graded right H-colinear, where the right H-coaction on $A \otimes A$ is defined by $(a \otimes a')_{[0]} \otimes (a \otimes a')_{[1]} = (|a_{[1]}|/|a'_{[0]}|)a_{[0]} \otimes a'_{[0]} \otimes (a_{[1]}a'_{[1]})$.

We say that C is a graded left H-module coalgebra if C is a graded left H-module such that $\Delta_C(h \rightharpoonup c) = (|h_2|/|c_1|)(h_1 \rightharpoonup c_1) \otimes (h_2 \rightharpoonup c_2)$ and $\epsilon_C(h \rightharpoonup c) = \epsilon_H(h)\epsilon_C(c)$. This is equivalent to say that Δ_C and ϵ_C are graded left H-linear, where the left H-action on $C \otimes C$ is defined by

$$h \rightharpoonup (c \otimes c') = (|h_2|/|c_1|)(h_1 \rightharpoonup c_1) \otimes (h_2 \rightharpoonup c_2).$$

We say that C is a graded right H-module coalgebra if C is a graded right H-module such that $\Delta_C(c \leftharpoonup h) = (|c_2|/|h_1|)(c_1 \leftharpoonup h_1) \otimes (c_2 \leftharpoonup h_2)$ and $\epsilon_C(c \leftharpoonup h) = \epsilon_H(h)\epsilon_C(c)$. This is equivalent to say that Δ_C and ϵ_C are graded right H-linear, where the right H-action on $C \otimes C$ is defined by

$$(c \otimes c') \leftharpoonup h = (|c_2|/|h_1|)(c_1 \leftharpoonup h_1) \otimes (c_2 \leftharpoonup h_2).$$

• Graded left-left Doi-Hopf modules.

Let A be a graded left H-comodule algebra and C a graded left H-module coalgebra. According to [5], we call the triple (H, A, C) a graded left-left Doi-Hopf datum.

The category ${}^{gr-C}_{gr-A}\mathcal{M}(H)$ of graded left-left Doi-Hopf modules is the category whose objects are the graded left A-modules and the graded left C-comodules M such that $\rho_{M,C}(am) = (|a_{[0]}|/|m_{-1}|)(a_{[-1]} \rightharpoonup m_{-1}) \otimes (a_{[0]}m_0)$. The morphisms of this category are the graded left A-linear maps and the graded left C-colinear maps. Any graded left-left Doi-Hopf datum (H, A, C) gives rise to a graded left-left entwining structure (A, C, ψ): the map ψ is defined by $\psi(a \otimes c) = (|a_{[0]}|/|c|)(a_{[-1]} \rightharpoonup c) \otimes a_{[0]}$. The corresponding category of graded left-left entwined modules coincides with the category ${}^{gr-C}_{gr-A}\mathcal{M}(H)$.

• Graded right-right Doi-Hopf modules.

Let A be a graded right H-comodule algebra and C a graded right H-module coalgebra. According to [5], we call the triple (H, A, C) a graded right-right Doi-Hopf datum.

The category $\mathcal{M}(H)^{gr-C}_{gr-A}$ of graded right-right Doi-Hopf modules is the category whose objects are the graded right A-modules and the graded right C-comodules M such that $\rho_{M,C}(ma) = (|m_1|/|a_{[0]}|)(m_0a_{[0]}) \otimes (m_1 \leftharpoonup a_{[1]})$. The morphisms of this category are the graded right A-linear maps and the graded right C-colinear maps. Any graded right-right Doi-Hopf datum (H, A, C) gives rise to a graded right-right entwining structure (A, C, ψ): the map ψ is defined by $\psi(c \otimes a) = (|c|/|a_{[0]}|)a_{[0]} \otimes (c \leftharpoonup a_{[1]})$. The corresponding category of graded right-right entwined modules coincides with $\mathcal{M}(H)^{gr-C}_{gr-A}$.

• Graded left-right Doi-Hopf modules.

Let A be a graded right H-comodule algebra and C a graded left H-module coalgebra. According to [5], we call the triple (H, A, C) a graded left-right Doi-Hopf datum.

The category ${}_{gr-A}\mathcal{M}(H)^{gr-C}$ of graded left-right Doi-Hopf modules is the category whose objects are the graded left A-modules and the graded right C-comodules M such that $\rho_{M,C}(am) = (|a_{[1]}|/|m_0|)(a_{[0]}m_0) \otimes (a_{[1]} \rightharpoonup m_1)$. The

morphisms of this category are the graded left A-linear maps and the graded right C-colinear maps. Any graded left-right Doi-Hopf datum (H, A, C) gives rise to a graded left-right entwining structure (A, C, ψ): the map ψ is defined by $\psi(a \otimes c) = a_{[0]} \otimes (a_{[1]} \rightharpoonup c)$. The corresponding category of graded left-right entwined modules coincides with ${}_{gr-A}\mathcal{M}(H)^{gr-C}$.

• Graded right-left Doi-Hopf modules.

Let A be a graded left H-comodule algebra and C a graded right H-module coalgebra. According to [5], we call the triple (H, A, C) a graded right-left Doi-Hopf datum.

The category ${}^{gr-C}\mathcal{M}(H)_{gr-A}$ of graded right-left Doi-Hopf modules is the category whose objects are the graded right A-modules and the graded left C-comodules M such that $\rho_{M,C}(ma) = (|m_0|/|a_{[-1]}|)(m_{-1} \leftharpoonup a_{[-1]}) \otimes (m_0 a_{[0]})$. The morphisms of this category are the graded right A-linear maps and the graded left C-colinear maps. Any graded right-left Doi-Hopf datum (H, A, C) gives rise to a graded right-left entwining structure (A, C, ψ): the map ψ is defined by $\psi(c \otimes a) = (c \leftharpoonup a_{[-1]}) \otimes a_{[0]}$. The corresponding category of graded right-left entwined modules coincides with ${}^{gr-C}\mathcal{M}(H)_{gr-A}$.

References

[1] Brzeziński, T.: *Galois comodules*. J. Algebra **290** (2005), 505–535. Zbl 1078.16039

[2] Brzeziński, T.: *The structure of corings. Induction functors, Maschke-type theorem, and Frobenius and Galois properties*. Algebr. Represent. Theory **5** (2002), 389–410. Zbl 1025.16017

[3] Brzeziński, T.: *The structure of corings with a grouplike element.* In: P. M. Hajac (ed.) et al., Noncommutative Geometry and Quantum Groups, Banach Cent. Publ. **61**, Polish Academy of Sciences, Warsaw (2003), 21–35. Zbl 1066.16034

[4] Brzeziński, T.; Wisbauer, R.: *Corings and Comodules*. London Mathematical Society Lecture Note Series **309**, Cambridge University Press, Cambridge 2003. Zbl 1035.16030

[5] Caenepeel, S.; Militaru, G.; Zhu, S.: *Frobenius and Separable Functors for Generalized Module Categories and Nonlinear Equations*. Lecture Notes in Mathematics **1787**, Springer Verlag, Berlin 2002. Zbl 1008.16036

[6] Caenepeel, S.; Vercruysse, J.; Shuanhong Wang: *Morita theory for corings and cleft entwining structures*. J. Algebra **276** (2004), 210–235. Zbl 1064.16037

[7] Caenepeel, S.; Guédénon, T.: *Projectivity and flatness over the endomorphism ring of a finitely generated module*. Int. J. Math. Math. Sci. **29–32** (2004), 1581–1588. Zbl 1081.16003

[8] Caenepeel, S.; Janssen, K.; Wang, S. H.: *Group corings*. Appl. Categ. Struct. **16** (2008), 65–96. Zbl 1145.16017

[9] Feldvoss, J.: *Representations of Lie colour algebras*. Adv. Math. **157** (2001), 95–137. Zbl 0998.17032

[10] García, J. J.; Del Río, A.: *On flatness and projectivity of a ring as a module over a fixed subring*. Math. Scand. **76** (1995), 179–192. Zbl 0842.16023

[11] Guédénon, T.: *Projectivity and flatness over the endomorphism ring of a finitely generated comodule*. Beitr. Algebra Geom. **49**(2) (2008), 399–408. Zbl pre05345496

[12] Guédénon, T.: *Semisimplicity of locally finite R#H-modules*. Algebra Colloq. **16**(1) (2009), 109–122. Zbl 1170.16029

[13] Năstăsescu, C.; Van Oystaeyen, F.: *Graded ring theory*. North-Holland, Amsterdam-New York-Oxford 1982. Zbl 0494.16001

[14] Palmieri, J. H.: *Stable Homotopy over the Steenrod Algebra*. Mem. Am. Math. Soc. **716** (2001). Zbl 0966.55013

[15] Sato, M.: *Fuller's theorem on equivalences*. J. Algebra **52** (1978), 274–284. Zbl 0374.16024

[16] Scheunert, M.; Zhang, R. B.: *The general linear supergroup and its Hopf superalgebra of regular functions*. J. Algebra **254** (2002), 44–83. Zbl 1047.17016

[17] Scheunert, M.: *Introduction to the cohomology of Lie superalgebras and some applications*. In: I. Bajo (ed.) et al., Recent advances in Lie theory, Res. Expo. Math. **25**, Heldermann, Lemgo 2002, 77–107. Zbl 1022.17013

Received January 15, 2009