General Mathematics Vol. 9, No. 1–2 (2001), 49–52

On the exponential diophantine equations of the form $a^x - b^y \cdot c^z = \pm 1$

Ana-Maria Acu, Mugur Acu

Abstract

In this short paper we obtain by elementary methods some general results for some exponential diophantine equations of the form $a^x - b^y \cdot c^z = \pm 1$

2000 Mathematical Subject Classification: 11D72

1 Introduction

This type of exponential diophantine equations was studied by D.Acu, L.J.Alex, L.S.Chen, J.H.Teng, Y.B.Wang.

Theorem 1 *[3] If p_1, p_2 are distinct prime numbers and not equal with 3, then the equation $7^x - p_1^y \cdot p_2^z = 1$ has no nonnegative integer solutions.*

Theorem 2 *[7] The equation $a^x - p_1^{y_1} p_2^{y_2} \dots p_k^{y_k} = \pm 1$ where a is a positive integer with $a > 1$ and $p_1, p_2, ..., p_k$ are distinct primes with $gcd(a, p_1 p_2 ... p_k) = 1$, has only finitely many positive integer solutions.*

Theorem 3 *[1] If a, b, c are distinct prime numbers and not equal with 2, then the equation $a^x - b^y \cdot c^z = \pm 1$ has no nonnegative integer solutions.*

2 New results

Theorem 4 *If a is a natural numbers, $a \equiv 1$ (mod p), and b, c are natural numbers with $bc \not\equiv 0$ (mod p), where p is a natural number with $p \geq 2$, then the equations $a^x - b^y \cdot c^z = 1$ have no nonnegative integer solutions.*

Proof. The equation $a^x - b^y \cdot c^z = 1$ is equivalent with

$$a^x - 1 = b^y \cdot c^z \tag{1}$$

From $bc \not\equiv 0$ (mod p) we obtain

$$b^y \cdot c^z \not\equiv 0 \ (mod\ p) \tag{2}$$

Because $a \equiv 1$ (mod p) we obtain $a^x \equiv 1$ (mod p), and thus we conclude

$$a^x - 1 \equiv 0 \ (mod\ p) \tag{3}$$

Using (1), (2), (3) we find a contradiction. We conclude that the equations $a^x - b^y \cdot c^z = 1$ have no nonnegative integer solutions.

Remark 1 *From theorem 4 with $a = 7, p = 3, b = p_1, c = p_2$, where p_1, p_2 are distinct prime numbers and not equal with 3, we obtain the result from theorem 1.*

Remark 2 *From theorem 4 with a, b, c distinct prime numbers and not equal with 2, and $p = 2$, we obtain a half of results from theorem 3.*

Theorem 5 *If a is a natural number, $a \equiv -1$ (mod p), and b, c are natural numbers with $bc \equiv 0$ (mod p), where p is a natural number with $p \geq 2$, then the equations $a^x - b^y \cdot c^z = -1$ have no nonnegative integer solutions with the form $(2k+1, y, z)$.*

Proof. By using congruences with p modulus, in a similar way with the proof of the theorem 4, we obtain the assertion.

Theorem 6 *If a is a natural number, $a \equiv 1$ (mod p), and $p_1, p_2, ..., p_k$ are natural numbers with $\prod_{i=1}^{k} p_i \not\equiv 0$ (mod p), where p is a natural number with $p \geq 2$, then the equations $a^x - \prod_{i=1}^{k} p_i^{y_i} = 1$ have no nonnegative integer solutions*

Proof. The equation $a^x - \prod_{i=1}^{k} p_i^{y_i} = 1$ is equivalent with $a^x - 1 = \prod_{i=1}^{k} p_i^{y_i}$. Using congruences with p modulus we show that $a^k - 1 \equiv 0 \pmod{p}$ and $\prod_{i=1}^{k} p_i^{y_i} \not\equiv 0 \pmod{p}$. Thus we find a contradiction and we obtain the assertion.

Remark 3 *It is easy to see that the Theorem 6 is complementary to the Theorem 2.*

Theorem 7 *The only solutions of the equation $3^x - 5^y \cdot 2^z = -1$ nonnegative integers are (0, 0, 1), (1, 0, 2) and (2, 1, 1).*

Proof. The equation $3^x - 5^y \cdot 2^z = -1$ is equivalent with

$$3^x + 1 = 5^y \cdot 2^z \tag{4}$$

If $z \geq 3$ then $2^z \equiv 0 \pmod{8}$. Using (4) we obtain $3^x + 1 \equiv 0 \pmod{8}$.

By using congruences with 8 modulus for $x = 0$, $x = 2p$ and respectively $x = 2p + 1$, we conclude that $3^x + 1 \not\equiv 0 \pmod{8}$.

If $y \geq 2$ then $5^y \equiv 0 \pmod{25}$. Using (4) we obtain $3^x + 1 \equiv 0 \pmod{25}$. By using congruences with 25 modulus $x = 4q$, $x = 4q+1$, $x = 4q+2$, and respectively $x = 4q+3$, we conclude that $3^x + 1 \not\equiv 0 \pmod{25}$.

Now we conclude that

$$y \in \{0, 1\} \text{ and } z \in \{0, 1, 2\} \tag{5}$$

By a simple calculation we obtain from (4) and (5) the only solutions (0, 0, 1), (1, 0, 2) and (2, 1, 1).

References

[1] A.M.Acu, M.Acu, *On the exponential diophantine equations of the forms $a^x - b^y \cdot c^z = \pm 1$ with a, b, c prime numbers*, General Mathematics, vol.8/2000, no.3-4.

[2] D.Acu, *On some exponential diophantine equations of the form $a^x - b^y \cdot c^z = \pm 1$*, Anal. Univ. de Vest(Timişoara), Mat.-Inf., Vol.XXXIV, Fasc.2 (1996), 167-172.

[3] D.Acu, *On some exponential diophantine equations of the form $a^x - b^y \cdot c^z = \pm 1$*, submitted.

[4] L.J. Alex, *On simple groups of order* $2^a3^b7^cp$, Journal of Algebra, 25(1)(1973), pag.113-124.

[5] L.J. Alex, *On simple groups of order* $2^a3^b7^cp$, Trans. Amar. Math. Soc., 173(1972), pag.389-399

[6] L.S.Chen, Y.B.Wang *On the exponential diophantine equation* $a^x - b^y \cdot c^z = \underline{+}1$, Heilongjiang Dexue Ziran Kexue Xuebao, 11 (2) (1994), pag.18-20 (MR 96B:11039).

[7] Y.H.Teng *On the diophantine equation* $a^x - p_1^{y_1}p_2^{y_2}...p_k^{y(k)} = \underline{+}1$ *(Chinese)*, Heilongjiang Dexue Ziran Kexue Xuebao, 8 (4) (1991), 57-59 (MR93C:11019).

Department of Mathematics
Faculty of Sciences
"Lucian Blaga" University of Sibiu
2400 Sibiu, Romania
E-mail address: *acu_mugur@yahoo.com*