INTEGERS: ELECTRONIC JOURNAL OF COMBINATORIAL NUMBER THEORY 7 (2007), #A46

# RECURRENCE FORMULAE FOR MULTI-POLY-BERNOULLI NUMBERS

**Y. Hamahata**
*Department of Mathematics, Tokyo University of Science, Noda, Chiba 278-8510, Japan*
hamahata@ma.noda.tus.ac.jp

**H. Masubuchi**
*Department of Mathematics, Tokyo University of Science, Noda, Chiba 278-8510, Japan*
yoshinori@ma.noda.tus.ac.jp

*Received: 4/3/07, Revised: 10/10/07, Accepted: 10/15/07, Published: 11/3/07*

**Abstract**

In this paper we establish recurrence formulae for multi-poly-Bernoulli numbers.

*—Dedicated to Professor Ryuichi Tanaka on the occasion of his sixtieth birthday*

## 1. Introduction and Background

Let us briefly recall poly-Bernoulli numbers.

For an integer $k \in \mathbb{Z}$, put

$$Li_k(z) = \sum_{n=1}^{\infty} \frac{z^n}{n^k}.$$

The formal power series $Li_k(z)$ is the $k$-th polylogarithm if $k \geq 1$, and a rational function if $k \leq 0$. When $k = 1$, $Li_1(z) = -\log(1-z)$. The formal power series $Li_k(z)$ can be used to introduce poly-Bernoulli numbers. The rational numbers $B_n^{(k)}$ $(n = 0, 1, 2, \ldots)$ are said to be *poly-Bernoulli numbers* if they satisfy

$$\frac{Li_k(1-e^{-x})}{1-e^{-x}} = \sum_{n=0}^{\infty} B_n^{(k)} \frac{x^n}{n!}.$$

In addition, for any $n \geq 0$, $B_n^{(1)}$ is the classical Bernoulli number, $B_n$. It was shown in [1] that special values of certain zeta functions at non-positive integers can be described in terms of poly-Bernoulli numbers. Furthermore, Kaneko [8] presented the following recurrences for poly-Bernoulli numbers.

**Theorem 1 (Kaneko)** (1) *For any $k \in \mathbb{Z}$ and $n \geq 0$,*

$$B_n^{(k)} = \frac{1}{n+1}\left\{ B_n^{(k-1)} - \sum_{m=1}^{n-1} \left( \begin{array}{c} n \\ m-1 \end{array} \right) B_m^{(k)} \right\}.$$

(2) *For any $k \geq 1$ and $n \geq 0$,*

$$B_n^{(k)} = \sum_{m=0}^{n} (-1)^m \left( \begin{array}{c} n \\ m \end{array} \right) B_{n-m}^{(k-1)} \left\{ \sum_{l=0}^{m} \frac{(-1)^l}{n-l+1} \left( \begin{array}{c} m \\ l \end{array} \right) B_l^{(1)} \right\}.$$

In this paper we consider generalized poly-Bernoulli numbers, which we refer to as *multi-poly-Bernoulli numbers*. Kim-Kim [9] introduced these numbers and proved that special values of certain zeta functions at non-positive integers can be described in terms of these numbers. In [5], we established a closed formula, and a duality property for special multi-poly-Bernoulli numbers. Bernoulli numbers satisfy certain recurrence relationships, which are used in many computations involving Bernoulli numbers. Obtaining a recurrence formula for multi-poly-Bernoulli numbers therefore seems to be a natural and important problem. The objective of this paper is thus to establish some recurrence formulae for multi-poly-Bernoulli numbers. It will be apparent that these recurrence formulae (Theorems 6, 7) are similar to those in the theorem above.

## 2. Multi-poly-Bernoulli Numbers

We first define a generalization of $Li_k(z)$. Let $r$ be an integer with a value greater than one.

**Definition 2** Let $k_1, k_2, \ldots, k_r$ be integers. Define

$$Li_{k_1,k_2,\ldots,k_r}(z) = \sum_{\substack{m_1,m_2,\ldots,m_r \in \mathbb{Z} \\ 0<m_1<m_2<\cdots<m_r}} \frac{z^{m_r}}{m_1^{k_1} \cdots m_r^{k_r}}.$$

Next let us establish the following fundamental result.

**Lemma 3**

$$\frac{d}{dz} Li_{k_1,k_2,\ldots,k_r}(z) = \left\{ \begin{array}{ll} \frac{1}{z} Li_{k_1,k_2,\ldots,k_{r-1},k_r-1}(z) & (k_r \neq 1) \\ \frac{1}{1-z} Li_{k_1,k_2,\ldots,k_{r-1}}(z) & (k_r = 1) \end{array} \right. . \tag{1}$$

The former case may be proven by means of a simple calculation. The latter case follows from

$$\begin{aligned} \sum_{0<m_1<\cdots<m_{r-1}<m_r} \frac{z^{m_r-1}}{m_1^{k_1} \cdots m_{r-1}^{k_{r-1}}} &= \sum_{0<m_1<\cdots<m_{r-1}} \sum_{m_r=m_{r-1}+1}^{\infty} \frac{z^{m_r-1}}{m_1^{k_1} \cdots m_{r-1}^{k_{r-1}}} \\ &= \sum_{0<m_1<\cdots<m_{r-1}} \frac{1}{m_1^{k_1} \cdots m_{r-1}^{k_{r-1}}} \cdot \frac{z^{m_{r-1}}}{1-z}. \end{aligned}$$

This lemma will be required later.

Let us now introduce a generalization of poly-Bernoulli numbers, making use of $Li_{k_1,k_2,\ldots,k_r}(z)$.

**Definition 4** Multi-poly-Bernoulli numbers $B_n^{(k_1,k_2,\ldots,k_r)}$ $(n = 0, 1, 2, \ldots)$ are defined for each integer $k_1, k_2, \ldots, k_r$ by the generating series

$$\frac{Li_{k_1,k_2,\ldots,k_r}(1-e^{-t})}{(1-e^{-t})^r} = \sum_{n=0}^{\infty} B_n^{(k_1,k_2,\ldots,k_r)} \frac{t^n}{n!}. \tag{2}$$

By definition, the left-hand side of (2) is

$$\frac{1}{1^{k_1}2^{k_2}\cdots r^{k_r}} + \sum_{\substack{0<m_1<\cdots<m_r \\ m_r\neq r}} \frac{(1-e^{-t})^{m_r-r}}{m_1^{k_1}\cdots m_r^{k_r}}.$$

Hence we have,

**Proposition 5**

$$B_0^{(k_1,k_2,\ldots,k_r)} = \frac{1}{1^{k_1}2^{k_2}\cdots r^{k_r}}.$$

The following tables show the values of $B_n^{(k_1,k_2)}$ and $B_n^{(k_1,k_2,k_3)}$ for small $n, k_i$.

| $\backslash n$ | 0 | 1 | 2 | 3 | 4 | 5 | 6 | 7 |
|---|---|---|---|---|---|---|---|---|
| $B_n^{(1,1)}$ | $\frac{1}{2}$ | $\frac{1}{2}$ | $\frac{5}{12}$ | $\frac{1}{4}$ | $\frac{1}{20}$ | $-\frac{1}{12}$ | $\frac{5}{84}$ | $\frac{1}{12}$ |
| $B_n^{(1,0)}$ | 1 | $\frac{3}{2}$ | $\frac{13}{6}$ | 3 | $\frac{119}{30}$ | 5 | $\frac{253}{42}$ | 7 |
| $B_n^{(0,1)}$ | $\frac{1}{2}$ | $\frac{2}{3}$ | $\frac{5}{6}$ | $\frac{29}{30}$ | $\frac{31}{30}$ | $\frac{43}{42}$ | $\frac{41}{42}$ | $\frac{29}{30}$ |
| $B_n^{(0,0)}$ | 1 | 2 | 4 | 8 | 16 | 32 | 64 | 128 |
| $B_n^{(0,-1)}$ | 2 | 6 | 18 | 54 | 162 | 486 | 1458 | 2574 |
| $B_n^{(-1,0)}$ | 1 | 3 | 9 | 27 | 81 | 243 | 729 | 1287 |
| $B_n^{(-1,-1)}$ | 2 | 9 | 39 | 165 | 687 | 2829 | 11505 | 46965 |

| $\backslash n$ | 0 | 1 | 2 | 3 | 4 | 5 | 6 | 7 |
|---|---|---|---|---|---|---|---|---|
| $B_n^{(1,1,1)}$ | $\frac{1}{6}$ | $\frac{1}{4}$ | $\frac{1}{3}$ | $\frac{3}{8}$ | $\frac{19}{60}$ | $\frac{1}{8}$ | $\frac{8}{63}$ | $\frac{5}{24}$ |
| $B_n^{(1,1,0)}$ | $\frac{1}{2}$ | 1 | $\frac{23}{12}$ | $\frac{7}{2}$ | $\frac{121}{20}$ | $\frac{59}{6}$ | $\frac{1255}{84}$ | $\frac{127}{6}$ |
| $B_n^{(1,0,1)}$ | $\frac{1}{3}$ | $\frac{5}{8}$ | $\frac{133}{120}$ | $\frac{221}{120}$ | $\frac{2383}{840}$ | $\frac{673}{168}$ | $\frac{4321}{840}$ | $\frac{145}{24}$ |
| $B_n^{(0,1,1)}$ | $\frac{1}{6}$ | $\frac{7}{24}$ | $\frac{19}{40}$ | $\frac{17}{24}$ | $\frac{53}{56}$ | $\frac{185}{168}$ | $\frac{303}{280}$ | $\frac{109}{120}$ |
| $B_n^{(1,0,0)}$ | 1 | $\frac{5}{2}$ | $\frac{37}{6}$ | 15 | $\frac{1079}{30}$ | 85 | $\frac{8317}{42}$ | 455 |
| $B_n^{(0,1,0)}$ | $\frac{1}{2}$ | $\frac{7}{6}$ | $\frac{8}{3}$ | $\frac{179}{30}$ | $\frac{147}{10}$ | $\frac{1177}{42}$ | $\frac{1238}{21}$ | $\frac{3659}{30}$ |
| $B_n^{(0,0,1)}$ | $\frac{1}{3}$ | $\frac{3}{4}$ | $\frac{33}{20}$ | $\frac{71}{20}$ | $\frac{1047}{140}$ | $\frac{433}{28}$ | $\frac{4411}{140}$ | $\frac{1271}{20}$ |
| $B_n^{(0,0,0)}$ | 1 | 3 | 9 | 27 | 81 | 243 | 729 | 2187 |
| $B_n^{(0,0,-1)}$ | 3 | 12 | 48 | 192 | 768 | 3072 | 12288 | 49152 |
| $B_n^{(0,-1,0)}$ | 2 | 8 | 32 | 128 | 512 | 2048 | 8192 | 32768 |
| $B_n^{(-1,0,0)}$ | 1 | 4 | 16 | 64 | 256 | 1024 | 4096 | 16384 |
| $B_n^{(0,-1,-1)}$ | 6 | 32 | 168 | 872 | 4488 | 22952 | 116808 | 592232 |
| $B_n^{(-1,0,-1)}$ | 3 | 16 | 84 | 436 | 2244 | 11476 | 58404 | 296116 |
| $B_n^{(-1,-1,0)}$ | 2 | 11 | 59 | 311 | 1619 | 8351 | 42779 | 217991 |
| $B_n^{(-1,-1,-1)}$ | 6 | 44 | 306 | 2054 | 13446 | 86414 | 547686 | 3434174 |

The remainder of this paper deals with recurrences for multi-poly-Bernoulli numbers.

## 3. Recurrence Formulae

We present two kinds of recurrence formulae for multi-poly-Bernoulli numbers. The first formula is as follows.

**Theorem 6**(1) *If $k_r \neq 1$ and $n \geq 1$, then*

$$B_n^{(k_1,k_2,\ldots,k_r)} = \frac{1}{n+r}\left\{B_n^{(k_1,\ldots,k_{r-1},k_r-1)} - \sum_{m=1}^{n-1}\binom{n}{m-1} B_m^{(k_1,k_2,\ldots,k_r)}\right\}.$$

(2) *If $k_r = 1$ and $n \geq 1$, then*

$$B_n^{(k_1,\ldots,k_{r-1},1)} = \frac{1}{n+r}\left[B_n^{(k_1,\ldots,k_{r-1})} - \sum_{m=0}^{n-1}(-1)^{n-m}\left\{r\binom{n}{m} + \binom{n}{m-1}\right\} B_m^{(k_1,\ldots,k_{r-1},1)}\right].$$

*Proof.* (1) Differentiate both sides of

$$Li_{k_1,k_2,\ldots,k_r}(1-e^{-t}) = (1-e^{-t})^r \sum_{n=0}^{\infty} B_n^{(k_1,k_2,\ldots,k_r)}\frac{t^n}{n!}.$$

By (1), we have

$$\text{L.H.S.} = \frac{e^{-t}}{1-e^{-t}} Li_{k_1,\ldots,k_{r-1},k_r-1}(1-e^{-t}).$$

On the other hand,

$$\begin{aligned}\text{R.H.S.} &= r(1-e^{-t})^{r-1}e^{-t}\sum_{n=0}^{\infty} B_n^{(k_1,k_2,\ldots,k_r)}\frac{t^n}{n!} \\ &\quad +(1-e^{-t})^r \sum_{n=1}^{\infty} B_n^{(k_1,k_2,\ldots,k_r)}\frac{t^{n-1}}{(n-1)!}.\end{aligned}$$

Hence,

$$\begin{aligned}&\frac{Li_{k_1,k_2,\ldots,k_r-1}(1-e^{-t})}{(1-e^{-t})^r} \\ &= r\sum_{n=0}^{\infty} B_n^{(k_1,k_2,\ldots,k_r)}\frac{t^n}{n!} + (e^t-1)\sum_{n=1}^{\infty} B_n^{(k_1,k_2,\ldots,k_r)}\frac{t^{n-1}}{(n-1)!} \\ &= r\sum_{n=0}^{\infty} B_n^{(k_1,k_2,\ldots,k_r)}\frac{t^n}{n!} + \sum_{n=1}^{\infty}\frac{t^n}{n!}\sum_{m=1}^{\infty} B_m^{(k_1,k_2,\ldots,k_r)}\frac{t^{m-1}}{(m-1)!} \\ &= r\sum_{n=0}^{\infty} B_n^{(k_1,k_2,\ldots,k_r)}\frac{t^n}{n!} + \sum_{m=1}^{\infty}\sum_{n=1}^{\infty} B_m^{(k_1,k_2,\ldots,k_r)}\frac{t^{n+m-1}}{n!(m-1)!}.\end{aligned}$$

Here we put $l = n+m-1$. The right-hand side of the last equation is then equal to

$$\begin{aligned}&r\sum_{n=0}^{\infty} B_n^{(k_1,k_2,\ldots,k_r)}\frac{t^n}{n!} + \sum_{m=1}^{\infty}\sum_{l=m}^{\infty} B_m^{(k_1,k_2,\ldots,k_r)}\frac{l!}{(l-(m-1))!(m-1)!}\cdot\frac{t^l}{l!} \\ &= r\sum_{n=0}^{\infty} B_n^{(k_1,k_2,\ldots,k_r)}\frac{t^n}{n!} + \sum_{l=1}^{\infty}\sum_{m=1}^{l} B_m^{(k_1,k_2,\ldots,k_r)}\left(\begin{array}{c} l \\ m-1 \end{array}\right)\frac{t^l}{l!} \\ &= r\sum_{n=0}^{\infty} B_n^{(k_1,k_2,\ldots,k_r)}\frac{t^n}{n!} + \sum_{n=1}^{\infty}\left(\sum_{m=1}^{n}\left(\begin{array}{c} n \\ m-1 \end{array}\right) B_m^{(k_1,k_2,\ldots,k_r)}\right)\frac{t^n}{n!}.\end{aligned}$$

Comparing the coefficients of both sides, for each $n \geq 1$,

$$\begin{aligned}B_n^{(k_1,\ldots,k_{r-1},k_r-1)} &= rB_n^{(k_1,k_2,\ldots,k_r)} + \sum_{m=1}^{n}\left(\begin{array}{c} n \\ m-1 \end{array}\right) B_m^{(k_1,k_2,\ldots,k_r)} \\ &= (n+r)B_n^{(k_1,k_2,\ldots,k_r)} + \sum_{m=1}^{n-1}\left(\begin{array}{c} n \\ m-1 \end{array}\right) B_m^{(k_1,k_2,\ldots,k_r)}.\end{aligned}$$

This implies the result.

(2) Differentiate both sides of

$$Li_{k_1,\ldots,k_{r-1},1}(1-e^{-t}) = (1-e^{-t})^r\sum_{n=0}^{\infty} B_n^{(k_1,\ldots,k_{r-1},1)}\frac{t^n}{n!}.$$

$$\begin{aligned}
\text{L.H.S.} &= \frac{e^{-t}}{1-(1-e^{-t})} Li_{k_1,\ldots,k_{r-1}}(1-e^{-t}) \\
&= Li_{k_1,\ldots,k_{r-1}}(1-e^{-t}), \\
\text{R.H.S.} &= r(1-e^{-t})^{r-1}e^{-t}\sum_{n=0}^{\infty} B_n^{(k_1,\ldots,k_{r-1},1)}\frac{t^n}{n!} + (1-e^{-t})^r \sum_{n=1}^{\infty} B_n^{(k_1,\ldots,k_{r-1},1)}\frac{t^{n-1}}{(n-1)!}.
\end{aligned}$$

Hence,

$$\begin{aligned}
&\sum_{n=0}^{\infty} B_n^{(k_1,\ldots,k_{r-1})}\frac{t^n}{n!} \\
&\quad = re^{-t}\sum_{n=0}^{\infty} B_n^{(k_1,\ldots,k_{r-1},1)}\frac{t^n}{n!} + (1-e^{-t})\sum_{n=1}^{\infty} B_n^{(k_1,\ldots,k_{r-1},1)}\frac{t^{n-1}}{(n-1)!} \\
&\quad = r\sum_{n=0}^{\infty}\frac{(-t)^n}{n!}\sum_{m=0}^{\infty} B_m^{(k_1,\ldots,k_{r-1},1)}\frac{t^m}{m!} - \sum_{n=1}^{\infty}\frac{(-t)^n}{n!}\sum_{m=0}^{\infty} B_{m+1}^{(k_1,\cdots,k_{r-1},1)}\frac{t^m}{m!} \\
&\quad = r\sum_{m=0}^{\infty}\sum_{n=0}^{\infty}(-1)^n B_m^{(k_1,\ldots,k_{r-1},1)}\frac{t^{n+m}}{n!m!} - \sum_{m=0}^{\infty}\sum_{n=1}^{\infty}(-1)^n B_{m+1}^{(k_1,\ldots,k_{r-1},1)}\frac{t^{n+m}}{n!m!}.
\end{aligned}$$

Here we put $l = n+m$. The right-hand side of the last equation then becomes

$$\begin{aligned}
&r\sum_{m=0}^{\infty}\sum_{l=m}^{\infty}(-1)^{l-m}B_m^{(k_1,\ldots,k_{r-1},1)}\frac{l!}{(l-m)!m!}\cdot\frac{t^l}{l!} \\
&\quad -\sum_{m=0}^{\infty}\sum_{l=m+1}^{\infty}(-1)^{l-m}B_{m+1}^{(k_1,\ldots,k_{r-1},1)}\frac{l!}{(l-m)!m!}\cdot\frac{t^l}{l!} \\
&= r\sum_{l=0}^{\infty}\sum_{m=0}^{l}(-1)^{l-m}B_m^{(k_1,\ldots,k_{r-1},1)}\binom{l}{m}\frac{t^l}{l!} \\
&\quad -\sum_{l=1}^{\infty}\sum_{m=0}^{l-1}(-1)^{l-m}B_{m+1}^{(k_1,\ldots,k_{r-1},1)}\binom{l}{m}\frac{t^l}{l!} \\
&= rB_0^{(k_1,\ldots,k_{r-1},1)} \\
&\quad +\sum_{n=1}^{\infty}\left\{\sum_{m=0}^{n}(-1)^{n-m}rB_m^{(k_1,\ldots,k_{r-1},1)}\binom{n}{m} - \sum_{m=0}^{n-1}(-1)^{n-m}B_{m+1}^{(k_1,\ldots,k_{r-1},1)}\binom{n}{m}\right\}\frac{t^n}{n!}.
\end{aligned}$$

Comparing both sides, for each $n \geq 1$,

$$\begin{aligned}
&B_n^{(k_1,\ldots,k_{r-1})} \\
&= \sum_{m=0}^{n}(-1)^{n-m}r\binom{n}{m}B_m^{(k_1,\ldots,k_{r-1},1)} + \sum_{m=1}^{n}(-1)^{n-m}\binom{n}{m-1}B_m^{(k_1,\ldots,k_{r-1},1)} \\
&= (n+r)B_n^{(k_1,\ldots,k_{r-1},1)} + \sum_{m=0}^{n-1}(-1)^{n-m}\left\{r\binom{n}{m}+\binom{n}{m-1}\right\}B_m^{(k_1,\ldots,k_{r-1},1)}.
\end{aligned}$$

These computations imply the result.

We obtain the second formula using the integral representation of $Li_{k_1,k_2,\ldots,k_r}(1-e^{-t})$.

**Theorem 7** (1) *If $k_r \neq 1$, then for any $n \geq 0$,*

$$B_n^{(k_1,k_2,\ldots,k_r)}$$
$$= (-1)^{r-1}\frac{n!}{(n+r-1)!}$$
$$\times\sum_{m=0}^{n}\left\{\sum_{p=0}^{n-m}\sum_{\substack{j_1+\cdots+j_{r-1}\\=n-m+r-1-p}}\frac{(-1)^{n-m+r-1-p}(n-m+r-1-p)!}{j_1!\cdots j_{r-1}!}\binom{n-m+r-1}{p}B_p^{(k_1,\ldots,k_{r-1},k_r-1)}\right\}$$
$$\times(-1)^m\binom{n+r-1}{m}\sum_{l=0}^{m}\frac{(-1)^l}{n-l+r}\left(\begin{array}{c} m \\ l \end{array}\right)\left\{\sum_{\substack{i_1+\cdots+i_r\\=l}}B_{i_1}^{(1)}\cdots B_{i_r}^{(1)}\frac{l!}{i_1!\cdots i_r!}\right\}.$$

(2) *If $k_r = 1$, then for any $n \geq 0$,*

$$B_n^{(k_1,k_2,\ldots,k_r)}$$
$$= (-1)^{r-1}\frac{n!}{(n+r-1)!}$$
$$\times\sum_{m=0}^{n}\left\{\sum_{p=0}^{n-m}\sum_{\substack{j_1+\cdots+j_{r-1}\\=n-m+r-1-p}}\frac{(-1)^{n-m+r-1-p}(n-m+r-1-p)!}{j_1!\cdots j_{r-1}!}\binom{n-m+r-1}{p}B_p^{(k_1,\ldots,k_{r-1})}\right\}$$
$$\times\binom{n+r-1}{m}\frac{1}{n-m+r}\left\{\sum_{\substack{i_1+\cdots+i_r\\=m}}B_{i_1}^{(1)}\cdots B_{i_r}^{(1)}\frac{m!}{i_1!\cdots i_r!}\right\}.$$

*Proof.* (1) Since

$$\frac{d}{dt}Li_{k_1,k_2,\ldots,k_r}(1-e^{-t}) = \frac{e^{-t}}{1-e^{-t}}Li_{k_1,\ldots,k_{r-1},k_r-1}(1-e^{-t}),$$

we have

$$Li_{k_1,k_2,\ldots,k_r}(1-e^{-t}) = \int_0^t \frac{e^{-s}}{1-e^{-s}}Li_{k_1,\ldots,k_{r-1},k_r-1}(1-e^{-s})ds.$$

By this equation,

$$\sum_{n=0}^{\infty}B_n^{(k_1,k_2,\ldots,k_r)}\frac{t^n}{n!}$$
$$= \frac{1}{(1-e^{-t})^r}\int_0^t\frac{e^{-s}}{1-e^{-s}}Li_{k_1,\ldots,k_{r-1},k_r-1}(1-e^{-s})ds$$
$$= \left(\frac{e^t}{e^t-1}\right)^r\int_0^t e^{-s}(1-e^{-s})^{r-1}\cdot\frac{Li_{k_1,\ldots,k_{r-1},k_r-1}(1-e^{-s})}{(1-e^{-s})^r}ds$$
$$= \left(\sum_{n=0}^{\infty}B_n^{(1)}\frac{t^{n-1}}{n!}\right)^r\int_0^t\sum_{n=0}^{\infty}\frac{(-s)^n}{n!}\left(-\sum_{n=1}^{\infty}\frac{(-s)^n}{n!}\right)^{r-1}\sum_{n=0}^{\infty}B_n^{(k_1,\ldots,k_{r-1},k_r-1)}\frac{s^n}{n!}ds$$

$$= (-1)^{r-1}\left(\sum_{n=0}^{\infty} B_n^{(1)}\frac{t^{n-1}}{n!}\right)^r \times \int_0^t \sum_{n=0}^{\infty}\frac{(-s)^n}{n!}\sum_{n=r-1}^{\infty}\sum_{\substack{j_1+\cdots+j_{r-1}\\=n}}\frac{(-1)^n s^n}{j_1!\cdots j_{r-1}!}\sum_{n=0}^{\infty} B_n^{(k_1,\ldots,k_{r-1},k_r-1)}\frac{s^n}{n!}ds.$$

We write $I_1$ for the integral part of the last equation.

$$I_1 = \int_0^t \sum_{n=0}^{\infty}\frac{(-s)^n}{n!}\sum_{m=0}^{\infty}\sum_{n=r-1}^{\infty}\sum_{\substack{j_1+\cdots+j_{r-1}\\=n}}\frac{(-1)^n n!}{j_1!\cdots j_{r-1}!}B_m^{(k_1,\ldots,k_{r-1},k_r-1)}\frac{s^{n+m}}{n!m!}ds.$$

Putting $l = n+m$,

$$I_1 = \int_0^t \sum_{n=0}^{\infty}\frac{(-s)^n}{n!}\sum_{m=0}^{\infty}\sum_{l=m+r-1}^{\infty}\sum_{\substack{j_1+\cdots+j_{r-1}\\=l-m}}\frac{(-1)^{l-m}(l-m)!}{j_1!\cdots j_{r-1}!}B_m^{(k_1,\ldots,k_{r-1},k_r-1)}\frac{l!}{(l-m)!m!}\cdot\frac{s^l}{l!}ds$$

$$= \int_0^t \sum_{n=0}^{\infty}\frac{(-s)^n}{n!}\sum_{l=r-1}^{\infty}\sum_{m=0}^{l-r+1}\sum_{\substack{j_1+\cdots+j_{r-1}\\=l-m}}\frac{(-1)^{l-m}(l-m)!}{j_1!\cdots j_{r-1}!}B_m^{(k_1,\ldots,k_{r-1},k_r-1)}\binom{l}{m}\frac{s^l}{l!}ds$$

$$= \int_0^t \sum_{l=r-1}^{\infty}\sum_{n=0}^{\infty}\sum_{m=0}^{l-r+1}(-1)^n\sum_{\substack{j_1+\cdots+j_{r-1}\\=l-m}}\frac{(-1)^{l-m}(l-m)!}{j_1!\cdots j_{r-1}!}B_m^{(k_1,\ldots,k_{r-1},k_r-1)}\binom{l}{m}\frac{s^{n+l}}{n!l!}ds.$$

We put $a = l+n$. Then

$$I_1 = \int_0^t \sum_{l=r-1}^{\infty}\sum_{a=l}^{\infty}\sum_{m=0}^{l-r+1}(-1)^{a-l}\sum_{\substack{j_1+\cdots+j_{r-1}\\=l-m}}\frac{(-1)^{l-m}(l-m)!}{j_1!\cdots j_{r-1}!}B_m^{(k_1,\ldots,k_{r-1},k_r-1)}\binom{l}{m}\frac{a!}{(a-l)!l!}\cdot\frac{s^a}{a!}ds$$

$$= \int_0^t \sum_{a=r-1}^{\infty}\sum_{l=r-1}^{a}\sum_{m=0}^{l-r+1}(-1)^{a-l}\sum_{\substack{j_1+\cdots+j_{r-1}\\=l-m}}\frac{(-1)^{l-m}(l-m)!}{j_1!\cdots j_{r-1}!}B_m^{(k_1,\ldots,k_{r-1},k_r-1)}\binom{l}{m}\binom{a}{l}\frac{s^a}{a!}ds$$

$$= \sum_{a=r-1}^{\infty}\sum_{l=r-1}^{a}\sum_{m=0}^{l-r+1}(-1)^{a-l}\sum_{\substack{j_1+\cdots+j_{r-1}\\=l-m}}\frac{(-1)^{l-m}(l-m)!}{j_1!\cdots j_{r-1}!}B_m^{(k_1,\ldots,k_{r-1},k_r-1)}\binom{l}{m}\binom{a}{l}\frac{t^{a+1}}{(a+1)!}.$$

Hence,

$$(-1)^{r-1}\left(\sum_{n=0}^{\infty} B_n^{(1)}\frac{t^{n-1}}{n!}\right)^r I_1$$
$$= (-1)^{r-1}\sum_{n=0}^{\infty}\sum_{\substack{i_1+\cdots+i_r\\=n}} B_{i_1}^{(1)}\cdots B_{i_r}^{(1)}\frac{t^{n-r}}{i_1!\cdots i_r!}$$
$$\times\sum_{a=r-1}^{\infty}\sum_{l=r-1}^{a}\sum_{m=0}^{l-r+1}(-1)^{a-l}\sum_{\substack{j_1+\cdots+j_{r-1}\\=l-m}}\frac{(-1)^{l-m}(l-m)!}{j_1!\cdots j_{r-1}!}B_m^{(k_1,\ldots,k_{r-1},k_r-1)}\binom{l}{m}\binom{a}{l}\frac{t^{a+1}}{(a+1)!}$$

$$= (-1)^{r-1} \sum_{a=r-1}^{\infty} \sum_{n=0}^{\infty} \sum_{l=r-1}^{a} \sum_{m=0}^{l-r+1} (-1)^{a-l} \sum_{\substack{i_1+\cdots+i_r \\ =n}} B_{i_1}^{(1)} \cdots B_{i_r}^{(1)} \frac{n!}{i_1! \cdots i_r!}$$
$$\times \sum_{\substack{j_1+\cdots+j_{r-1} \\ =l-m}} \frac{(-1)^{l-m}(l-m)!}{j_1! \cdots j_{r-1}!} B_m^{(k_1,\ldots,k_{r-1},k_r-1)} \binom{l}{m} \binom{a}{l} \frac{t^{n+a+1}}{n!(a+1)!} \cdot t^{-r}$$

$$= (-1)^{r-1} \sum_{a=r-1}^{\infty} \sum_{b=a+1}^{\infty} \sum_{l=r-1}^{a} \sum_{m=0}^{l-r+1} (-1)^{a-l} \sum_{\substack{i_1+\cdots+i_r \\ =b-a-1}} B_{i_1}^{(1)} \cdots B_{i_r}^{(1)} \frac{(b-a-1)!}{i_1! \cdots i_r!}$$
$$\times \sum_{\substack{j_1+\cdots+j_{r-1} \\ =l-m}} \frac{(-1)^{l-m}(l-m)!}{j_1! \cdots j_{r-1}!} B_m^{(k_1,\ldots,k_{r-1},k_r-1)} \binom{l}{m} \binom{a}{l} \frac{b!}{(b-(a+1))!(a+1)!} \cdot \frac{t^b}{b!} \cdot t^{-r}$$
$$(\text{put } b = n+a+1)$$

$$= (-1)^{r-1} \sum_{b=r}^{\infty} \sum_{a=r-1}^{b-1} \sum_{l=r-1}^{a} \sum_{m=0}^{l-r+1} (-1)^{a-l} \sum_{\substack{i_1+\cdots+i_r \\ =b-a-1}} B_{i_1}^{(1)} \cdots B_{i_r}^{(1)} \frac{(b-a-1)!}{i_1! \cdots i_r!}$$
$$\times \sum_{\substack{j_1+\cdots+j_{r-1} \\ =l-m}} \frac{(-1)^{l-m}(l-m)!}{j_1! \cdots j_{r-1}!} B_m^{(k_1,\ldots,k_{r-1},k_r-1)} \binom{l}{m} \binom{a}{l} \binom{b}{a+1} \frac{t^{b-r}}{b!}$$

$$= (-1)^{r-1} \sum_{n=0}^{\infty} \sum_{a=r-1}^{n+r-1} \sum_{l=r-1}^{a} \sum_{m=0}^{l-r+1} (-1)^{a-l} \sum_{\substack{i_1+\cdots+i_r \\ =n-a+r-1}} B_{i_1}^{(1)} \cdots B_{i_r}^{(1)} \frac{(n-a+r-1)!}{i_1! \cdots i_r!}$$
$$\times \sum_{\substack{j_1+\cdots+j_{r-1} \\ =l-m}} \frac{(-1)^{l-m}(l-m)!}{j_1! \cdots j_{r-1}!} B_m^{(k_1,\ldots,k_{r-1},k_r-1)} \binom{l}{m} \binom{a}{l} \binom{n+r}{a+1} \frac{t^n}{(n+r)!}$$
$$(\text{put } n = b-r)$$

$$= \sum_{n=0}^{\infty} \left\{ (-1)^{r-1} \frac{n!}{(n+r)!} \sum_{a=r-1}^{n+r-1} \left( \sum_{\substack{i_1+\cdots+i_r \\ =n-a+r-1}} B_{i_1}^{(1)} \cdots B_{i_r}^{(1)} \frac{(n-a+r-1)!}{i_1! \cdots i_r!} \right) \right.$$
$$\left. \times \sum_{l=r-1}^{a} (-1)^{a-l} \binom{n+r}{a+1} \binom{a}{l} \sum_{m=0}^{l-r+1} \sum_{\substack{j_1+\cdots+j_{r-1} \\ =l-m}} \frac{(-1)^{l-m}(l-m)!}{j_1! \cdots j_{r-1}!} \binom{l}{m} B_m^{(k_1,\ldots,k_{r-1},k_r-1)} \right\} \frac{t^n}{n!}.$$

If we compare the coefficients of both sides,

$$
\begin{aligned}
&B_n^{(k_1,k_2,\ldots,k_r)} \\
&\quad = (-1)^{r-1}\frac{n!}{(n+r)!}\sum_{a=r-1}^{n+r-1}\left(\sum_{\substack{i_1+\cdots+i_r\\=n-a+r-1}} B_{i_1}^{(1)}\cdots B_{i_r}^{(1)}\frac{(n-a+r-1)!}{i_1!\cdots i_r!}\right) \\
&\quad\times\sum_{l=r-1}^{a}(-1)^{a-l}\left(\begin{array}{c} n+r \\ a+1 \end{array}\right)\left(\begin{array}{c} a \\ l \end{array}\right)\sum_{m=0}^{l-r+1}\sum_{\substack{j_1+\cdots+j_{r-1}\\=l-m}}\frac{(-1)^{l-m}(l-m)!}{j_1!\cdots j_{r-1}!}\left(\begin{array}{c} l \\ m \end{array}\right)B_m^{(k_1,\ldots,k_{r-1},k_r-1)} \\
&\quad = (-1)^{r-1}\frac{n!}{(n+r-1)!}\sum_{l=r-1}^{n+r-1}\sum_{m=0}^{l-r+1}\sum_{\substack{j_1+\cdots+j_{r-1}\\=l-m}}\frac{(-1)^{l-m}(l-m)!}{j_1!\cdots j_{r-1}!}\left(\begin{array}{c} l \\ m \end{array}\right)B_m^{(k_1,\ldots,k_{r-1},k_r-1)} \\
&\quad\times\sum_{a=l}^{n+r-1}\frac{(-1)^{a-l}}{a+1}\binom{n+r-1}{a}\left(\begin{array}{c} a \\ l \end{array}\right)\left(\sum_{\substack{i_1+\cdots+i_r\\=n-a+r-1}} B_{i_1}^{(1)}\cdots B_{i_r}^{(1)}\frac{(n-a+r-1)!}{i_1!\cdots i_r!}\right) \\
&\qquad\qquad\left(\text{by } \tbinom{n+r}{a+1} = \frac{n+r}{a+1}\tbinom{n+r-1}{a} \text{ and } \sum_{a=r-1}^{n+r-1}\sum_{l=r-1}^{a} = \sum_{l=r-1}^{n+r-1}\sum_{a=l}^{n+r-1}\right) \\
&\quad = (-1)^{r-1}\frac{n!}{(n+r-1)!} \\
&\quad\times\sum_{l=r-1}^{n+r-1}(-1)^{l}\binom{n+r-1}{l}\left\{\sum_{m=0}^{l-r+1}\sum_{\substack{j_1+\cdots+j_{r-1}\\=l-m}}\frac{(-1)^{l-m}(l-m)!}{j_1!\cdots j_{r-1}!}\left(\begin{array}{c} l \\ m \end{array}\right)B_m^{(k_1,\ldots,k_{r-1},k_r-1)}\right\} \\
&\quad\times\sum_{a=l}^{n+r-1}\frac{(-1)^{a}}{a+1}\left(\begin{array}{c} n+r-1-l \\ n+r-1-a \end{array}\right)\left\{\sum_{\substack{i_1+\cdots+i_r\\=n-a+r-1}} B_{i_1}^{(1)}\cdots B_{i_r}^{(1)}\frac{(n-a+r-1)!}{i_1!\cdots i_r!}\right\} \\
&\qquad\qquad\left(\text{by } \tbinom{n+r-1}{a}\tbinom{a}{l} = \tbinom{n+r-1}{l}\left(\begin{array}{c} n+r-1-l \\ n+r-1-a \end{array}\right)\right) \\
&\quad = (-1)^{r-1}\frac{n!}{(n+r-1)!}\sum_{l'=0}^{n}(-1)^{l'+r-1}\binom{n+r-1}{l'+r-1} \\
&\quad\times\left\{\sum_{m=0}^{l'}\sum_{\substack{j_1+\cdots+j_{r-1}\\=l'+r-1-m}}\frac{(-1)^{l'+r-1-m}(l'+r-1-m)!}{j_1!\cdots j_{r-1}!}\left(\begin{array}{c} l'+r-1 \\ m \end{array}\right)B_m^{(k_1,\ldots,k_{r-1},k_r-1)}\right\} \\
&\quad\times\sum_{a=l'+r-1}^{n+r-1}\frac{(-1)^{a}}{a+1}\left(\begin{array}{c} n-l' \\ n+r-1-a \end{array}\right)\left\{\sum_{\substack{i_1+\cdots+i_r\\=n-a+r-1}} B_{i_1}^{(1)}\cdots B_{i_r}^{(1)}\frac{(n-a+r-1)!}{i_1!\cdots i_r!}\right\} \\
&\qquad\qquad(\text{put } l' = l-r+1)
\end{aligned}
$$

$$= \frac{n!}{(n+r-1)!}\sum_{l'=0}^{n}(-1)^{l'}\binom{n+r-1}{l'+r-1}$$
$$\times\left\{\sum_{m=0}^{l'}\sum_{\substack{j_1+\cdots+j_{r-1}\\=l'+r-1-m}}\frac{(-1)^{l'+r-1-m}(l'+r-1-m)!}{j_1!\cdots j_{r-1}!}\left(\begin{array}{c} l'+r-1 \\ m \end{array}\right)B_m^{(k_1,\ldots,k_{r-1},k_r-1)}\right\}$$
$$\times\sum_{a'=l'}^{n}\frac{(-1)^{a'+r-1}}{a'+r}\left(\begin{array}{c} n-l' \\ n-a' \end{array}\right)\left\{\sum_{\substack{i_1+\cdots+i_r\\=n-a'}}B_{i_1}^{(1)}\cdots B_{i_r}^{(1)}\frac{(n-a')!}{i_1!\cdots i_r!}\right\}$$
(put $a' = a-r+1$)

$$= \frac{n!}{(n+r-1)!}\sum_{l''=0}^{n}(-1)^{n-l''}\binom{n+r-1}{n+r-1-l''}$$
$$\times\left\{\sum_{m=0}^{n-l''}\sum_{\substack{j_1+\cdots+j_{r-1}\\=n-l''+r-1-m}}\frac{(-1)^{n-l''+r-1-m}(n-l''+r-1-m)!}{j_1!\cdots j_{r-1}!}\binom{n-l''+r-1}{m}B_m^{(k_1,\ldots,k_{r-1},k_r-1)}\right\}$$
$$\times\sum_{a'=n-l''}^{n}\frac{(-1)^{a'+r-1}}{a'+r}\left(\begin{array}{c} l'' \\ n-a' \end{array}\right)\left\{\sum_{\substack{i_1+\cdots+i_r\\=n-a'}}B_{i_1}^{(1)}\cdots B_{i_r}^{(1)}\frac{(n-a')!}{i_1!\cdots i_r!}\right\}$$
(put $l'' = n-l'$)

$$= \frac{n!}{(n+r-1)!}\sum_{l''=0}^{n}(-1)^{n-l''}\binom{n+r-1}{l''}$$
$$\times\left\{\sum_{m=0}^{n-l''}\sum_{\substack{j_1+\cdots+j_{r-1}\\=n-l''+r-1-m}}\frac{(-1)^{n-l''+r-1-m}(n-l''+r-1-m)!}{j_1!\cdots j_{r-1}!}\binom{n-l''+r-1}{m}B_m^{(k_1,\ldots,k_{r-1},k_r-1)}\right\}$$
$$\times\sum_{a''=0}^{l''}\frac{(-1)^{n+r-1-a''}}{n-a''+r}\left(\begin{array}{c} l'' \\ a'' \end{array}\right)\left\{\sum_{\substack{i_1+\cdots+i_r\\=a''}}B_{i_1}^{(1)}\cdots B_{i_r}^{(1)}\frac{a''!}{i_1!\cdots i_r!}\right\}$$
(put $a'' = n-a'$)

$$= (-1)^{r-1}\frac{n!}{(n+r-1)!}\sum_{l''=0}^{n}(-1)^{l''}\binom{n+r-1}{l''}$$

$$\times\left\{\sum_{m=0}^{n-l''}\sum_{\substack{j_1+\cdots+j_{r-1}\\=n-l''+r-1-m}}\frac{(-1)^{n-l''+r-1-m}(n-l''+r-1-m)!}{j_1!\cdots j_{r-1}!}\binom{n-l''+r-1}{m}B_m^{(k_1,\ldots,k_{r-1},k_r-1)}\right\}$$

$$\times\sum_{a''=0}^{l''}\frac{(-1)^{a''}}{n-a''+r}\left(\begin{array}{c} l''\\ a''\end{array}\right)\left\{\sum_{\substack{i_1+\cdots+i_r\\=a''}}B_{i_1}^{(1)}\cdots B_{i_r}^{(1)}\frac{a''!}{i_1!\cdots i_r!}\right\}.$$

(put $a''=n-a'$)

This shows the claim.

(2) Since

$$\frac{d}{dt}Li_{k_1,k_2,\ldots,k_{r-1},1}(1-e^{-t}) = Li_{k_1,\ldots,k_{r-1}}(1-e^{-t}),$$

we have

$$Li_{k_1,\ldots,k_{r-1},1}(1-e^{-t}) = \int_0^t Li_{k_1,\ldots,k_{r-1}}(1-e^{-s})ds.$$

Hence

$$\sum_{n=0}^{\infty}B_n^{(k_1,\ldots,k_{r-1},1)}\frac{t^n}{n!}$$

$$= \left(\frac{e^t}{e^t-1}\right)^r\int_0^t(1-e^{-s})^{r-1}\cdot\frac{Li_{k_1,\ldots,k_{r-1}}(1-e^{-s})}{(1-e^{-s})^{r-1}}ds$$

$$= \left(\sum_{n=0}^{\infty}B_n^{(1)}\frac{t^{n-1}}{n!}\right)^r\int_0^t\left(-\sum_{n=1}^{\infty}\frac{(-s)^n}{n!}\right)^{r-1}\sum_{n=0}^{\infty}B_n^{(k_1,\ldots,k_{r-1})}\frac{s^n}{n!}ds$$

$$= (-1)^{r-1}\left(\sum_{n=0}^{\infty}B_n^{(1)}\frac{t^{n-1}}{n!}\right)^r\int_0^t\sum_{n=r-1}^{\infty}\sum_{\substack{j_1+\cdots+j_{r-1}\\=n}}\frac{(-1)^n s^n}{j_1!\cdots j_{r-1}!}\sum_{m=0}^{\infty}B_m^{(k_1,\ldots,k_{r-1})}\frac{s^m}{m!}ds.$$

Denote by $I_2$ the integral part of the last equation.

$$I_2 = \int_0^t\sum_{m=0}^{\infty}\sum_{n=r-1}^{\infty}\sum_{\substack{j_1+\cdots+j_{r-1}\\=n}}\frac{(-1)^n n!}{j_1!\cdots j_{r-1}!}B_m^{(k_1,\ldots,k_{r-1})}\frac{s^{n+m}}{n!m!}ds$$

$$= \int_0^t\sum_{m=0}^{\infty}\sum_{l=m+r-1}^{\infty}\sum_{\substack{j_1+\cdots+j_{r-1}\\=l-m}}\frac{(-1)^{l-m}(l-m)!}{j_1!\cdots j_{r-1}!}B_n^{(k_1,\ldots,k_{r-1})}\frac{l!}{(l-m)!m!}\cdot\frac{s^l}{l!}ds$$

(put $l=n+m$)

$$= \int_0^t\sum_{l=r-1}^{\infty}\sum_{m=0}^{l-r+1}\sum_{\substack{j_1+\cdots+j_{r-1}\\=l-m}}\frac{(-1)^{l-m}(l-m)!}{j_1!\cdots j_{r-1}!}B_m^{(k_1,\ldots,k_{r-1})}\left(\begin{array}{c} l\\ m\end{array}\right)\frac{s^l}{l!}ds$$

$$= \sum_{l=r-1}^{\infty} \sum_{m=0}^{l-r+1} \sum_{\substack{j_1+\cdots+j_{r-1} \\ =l-m}} \frac{(-1)^{l-m}(l-m)!}{j_1!\cdots j_{r-1}!} B_m^{(k_1,\ldots,k_{r-1})} \left( \begin{array}{c} l \\ m \end{array} \right) \frac{t^{l+1}}{(l+1)!}.$$

Thus,

$$(-1)^{r-1} \left( \sum_{n=0}^{\infty} B_n^{(1)} \frac{t^{n-1}}{n!} \right)^r I_2$$
$$= (-1)^{r-1} \sum_{n=0}^{\infty} \sum_{\substack{i_1+\cdots+i_r \\ =n}} B_{i_1}^{(1)} \cdots B_{i_r}^{(1)} \frac{t^{n-r}}{i_1!\cdots i_r!}$$
$$\times \sum_{l=r-1}^{\infty} \sum_{m=0}^{l-r+1} \sum_{\substack{j_1+\cdots+j_{r-1} \\ =l-m}} \frac{(-1)^{l-m}(l-m)!}{j_1!\cdots j_{r-1}!} B_m^{(k_1,\ldots,k_{r-1})} \left( \begin{array}{c} l \\ m \end{array} \right) \frac{t^{l+1}}{(l+1)!}$$

$$= (-1)^{r-1} \sum_{l=r-1}^{\infty} \sum_{n=0}^{\infty} \sum_{m=0}^{l-r+1} \sum_{\substack{i_1+\cdots+i_r \\ =n}} B_{i_1}^{(1)} \cdots B_{i_r}^{(1)} \frac{n!}{i_1!\cdots i_r!} \sum_{\substack{j_1+\cdots+j_{r-1} \\ =l-m}} \frac{(-1)^{l-m}(l-m)!}{j_1!\cdots j_{r-1}!}$$
$$\times B_m^{(k_1,\ldots,k_{r-1})} \left( \begin{array}{c} l \\ m \end{array} \right) \frac{t^{n+l+1}}{n!(l+1)!} \cdot t^{-r}$$

$$= (-1)^{r-1} \sum_{l=r-1}^{\infty} \sum_{a=l+1}^{\infty} \sum_{m=0}^{l-r+1} \sum_{\substack{i_1+\cdots+i_r \\ =a-l-1}} B_{i_1}^{(1)} \cdots B_{i_r}^{(1)} \frac{(a-l-1)!}{i_1!\cdots i_r!} \sum_{\substack{j_1+\cdots+j_{r-1} \\ =l-m}} \frac{(-1)^{l-m}(l-m)!}{j_1!\cdots j_{r-1}!}$$
$$\times B_m^{(k_1,\ldots,k_{r-1})} \left( \begin{array}{c} l \\ m \end{array} \right) \frac{a!}{(a-(l+1))!(l+1)!} \cdot \frac{t^a}{a!} \cdot t^{-r}$$
(put $a = n+l+1$)

$$= (-1)^{r-1} \sum_{a=r}^{\infty} \sum_{l=r-1}^{a-1} \sum_{m=0}^{l-r+1} \sum_{\substack{i_1+\cdots+i_r \\ =a-l-1}} B_{i_1}^{(1)} \cdots B_{i_r}^{(1)} \frac{(a-l-1)!}{i_1!\cdots i_r!} \sum_{\substack{j_1+\cdots+j_{r-1} \\ =l-m}} \frac{(-1)^{l-m}(l-m)!}{j_1!\cdots j_{r-1}!}$$
$$\times B_m^{(k_1,\ldots,k_{r-1})} \left( \begin{array}{c} l \\ m \end{array} \right) \left( \begin{array}{c} a \\ l+1 \end{array} \right) \frac{t^{a-r}}{a!}$$

$$= (-1)^{r-1} \sum_{n=0}^{\infty} \sum_{l=r-1}^{n+r-1} \sum_{m=0}^{l-r+1} \sum_{\substack{i_1+\cdots+i_r \\ =n-l+r-1}} B_{i_1}^{(1)} \cdots B_{i_r}^{(1)} \frac{n!}{i_1!\cdots i_r!} \sum_{\substack{j_1+\cdots+j_{r-1} \\ =l-m}} \frac{(-1)^{l-m}(l-m)!}{j_1!\cdots j_{r-1}!}$$
$$\times B_m^{(k_1,\ldots,k_{r-1})} \left( \begin{array}{c} l \\ m \end{array} \right) \left( \begin{array}{c} n+r \\ l+1 \end{array} \right) \frac{t^n}{(n+r)!}$$
(put $n = a-r$)

$$= \sum_{n=0}^{\infty} \left\{ \frac{(-1)^{r-1} n!}{(n+r)!} \sum_{l=r-1}^{n+r-1} \binom{n+r}{l+1} \sum_{\substack{i_1+\cdots+i_r \\ =n-l+r-1}} B_{i_1}^{(1)} \cdots B_{i_r}^{(1)} \frac{(n-l+r-1)!}{i_1! \cdots i_r!} \right.$$

$$\left. \times \sum_{m=0}^{l-r+1} \left( \begin{array}{c} l \\ m \end{array} \right) \sum_{\substack{j_1+\cdots+j_{r-1} \\ =l-m}} \frac{(-1)^{l-m}(l-m)!}{j_1! \cdots j_{r-1}!} B_m^{(k_1,\ldots,k_{r-1})} \right\} \frac{t^n}{n!}.$$

If we compare the coefficients of both sides of the equation,

$$B_n^{(k_1,k_2,\ldots,k_r)} = \frac{(-1)^{r-1} n!}{(n+r)!} \sum_{l=r-1}^{n+r-1} \binom{n+r}{l+1} \left( \sum_{\substack{i_1+\cdots+i_r \\ =n-l+r-1}} B_{i_1}^{(1)} \cdots B_{i_r}^{(1)} \frac{(n-l+r-1)!}{i_1! \cdots i_r!} \right)$$

$$\times \sum_{m=0}^{l-r+1} \left( \sum_{\substack{j_1+\cdots+j_{r-1} \\ =l-m}} \frac{(-1)^{l-m}(l-m)!}{j_1! \cdots j_{r-1}!} \right) \left( \begin{array}{c} l \\ m \end{array} \right) B_m^{(k_1,\ldots,k_{r-1})}$$

$$= \frac{(-1)^{r-1} n!}{(n+r-1)!} \sum_{l=r-1}^{n+r-1} \frac{1}{l+1} \binom{n+r-1}{l} \left( \sum_{\substack{i_1+\cdots+i_r \\ =n+r-1-l}} B_{i_1}^{(1)} \cdots B_{i_r}^{(1)} \frac{(n-l+r-1)!}{i_1! \cdots i_r!} \right)$$

$$\times \sum_{m=0}^{l-r+1} \left( \sum_{\substack{j_1+\cdots+j_{r-1} \\ =l-m}} \frac{(-1)^{l-m}(l-m)!}{j_1! \cdots j_{r-1}!} \right) \left( \begin{array}{c} l \\ m \end{array} \right) B_m^{(k_1,\ldots,k_{r-1})}$$

$$\left( \text{by } \tbinom{n+r}{l+1} = \frac{n+r}{l+1} \tbinom{n+r-1}{l} \right)$$

$$= \frac{(-1)^{r-1} n!}{(n+r-1)!} \sum_{a=0}^{n} \frac{1}{a+r} \binom{n+r-1}{a+r-1} \left( \sum_{\substack{i_1+\cdots+i_r \\ =n-a}} B_{i_1}^{(1)} \cdots B_{i_r}^{(1)} \frac{(n-a)!}{i_1! \cdots i_r!} \right)$$

$$\times \sum_{m=0}^{a} \left( \sum_{\substack{j_1+\cdots+j_{r-1} \\ =a+r-1-m}} \frac{(-1)^{a+r-1-m}(a+r-1-m)!}{j_1! \cdots j_{r-1}!} \right) \left( \begin{array}{c} a+r-1 \\ m \end{array} \right) B_m^{(k_1,\ldots,k_{r-1})}$$

$$(\text{put } a = l-r+1)$$

$$= \frac{(-1)^{r-1} n!}{(n+r-1)!} \sum_{l'=0}^{n} \frac{1}{n-l'+r} \binom{n+r-1}{n+r-1-l'} \left( \sum_{\substack{i_1+\cdots+i_r \\ =l'}} B_{i_1}^{(1)} \cdots B_{i_r}^{(1)} \frac{l'!}{i_1! \cdots i_r!} \right)$$

$$\times \sum_{m=0}^{n-l'} \left( \sum_{\substack{j_1+\cdots+j_{r-1} \\ =n-l'+r-1-m}} \frac{(-1)^{n-l'+r-1-m}(n-l'+r-1-m)!}{j_1! \cdots j_{r-1}!} \right) \binom{n-l'+r-1}{m} B_m^{(k_1,\ldots,k_{r-1})}.$$

$$(\text{put } l' = n-a)$$

This shows the claim.

**Remark 8** Some recurrences exist for the original Bernoulli numbers $B_n^{(1)} = B_n$ $(n \geq 0)$, the case in which the upper index is fixed with the value (1). The problem of finding a recurrence among $B_n^{(k_1,k_2,\ldots,k_r)}$ $(n \geq 0)$ for fixed $(k_1, k_2, \ldots, k_r)$, is therefore intriguing.

## References

[1] T. Arakawa, M. Kaneko, Multiple zeta values, poly-Bernoulli numbers, and related zeta functions. Nagoya J. Math. **153** (1999), 189-209.

[2] T. Arakawa, M. Kaneko, On poly-Bernoulli numbers. Comment. Math. Univ. St. Pauli **48** (1999), 159-167.

[3] L. Carlitz, Some theorems on Bernoulli numbers of higher order. Pacific J. Math. **2** (1952), 127-139.

[4] R. Graham, D. Knuth, O. Patashnik, Concrete Mathematics. Addison-Wesley, 1989.

[5] Y. Hamahata, H. Masubuchi, Special multi-poly-Bernoulli numbers. Journal of Integer Sequences **10** (2007), Article 07.4.1. 6pp.

[6] K. Ireland, M. Rosen, A Classical Introduction to Modern Number Theory. Second Edition, Graduate Texts in Mathematics **84** Springer-Verlag, 1990.

[7] M. Kaneko, Poly-Bernoulli numbers. J. Théorie de Nombres **9** (1997), 221-228.

[8] M. Kaneko, Multiple Zeta Values and Poly-Bernoulli Numbers. Tokyo Metropolitan University Seminar Report, 1997.

[9] M.-S. Kim, T. Kim, An explicit formula on the generalized Bernoulli number with order $n$. Indian J. Appl. Math. **31** (2000), 1455-1461.