INTEGERS: ELECTRONIC JOURNAL OF COMBINATORIAL NUMBER THEORY 7 (2007), #A52

TWO GENERALIZED CONSTANTS RELATED TO ZERO-SUM PROBLEMS FOR TWO SPECIAL SETS

Xingwu Xia
School of Mathematics and Computational Science, Sun Yat-Sen University, Guangzhou 510275, P.R. China
xxwsjtu@yahoo.com.cn

Received: 3/24/07, Revised: 11/8/07, Accepted: 11/13/07, Published: 11/28/07

Abstract

Let $n \in \mathbb{N}$ and $A \subseteq \mathbb{Z}_n$ be such that A is non–empty and does not contain 0. Adhikari et al proposed two generalized constants related to the zero-sum problem. One is $D_A(n)$, which denotes the least natural number k such that for any sequence $(x_1, \cdots, x_k) \in \mathbb{Z}^k$, there exists a non-empty subsequence $(x_{j_1}, \cdots, x_{j_l})$ and $(a_1, \cdots, a_l) \in A^l$ such that $\sum_{i=1}^{l} a_i x_{j_i} \equiv 0 \pmod{n}$. The other is $E_A(n)$, defined as the smallest $t \in \mathbb{N}$ such that for all sequences $(x_1, \ldots, x_t) \in \mathbb{Z}^t$, there exist indices $j_1, \ldots, j_n \in \mathbb{N}, 1 \leq j_1 < \cdots < j_n \leq t$ and $(\vartheta_1, \cdots, \vartheta_n) \in A^n$ with $\sum_{i=1}^{n} \vartheta_i x_{j_i} \equiv 0 \pmod{n}$. S. D. Adhikari et al proposed characterizing any other sets for which $E_A(n) = n + 1$ or even those for which $E_A(n) = n + j$ for specific small values of j. In this paper we give two kinds of sets, calculate $D_A(n)$ and $E_A(n)$ for these sets, and partially solve Adhikari's problem.

1. Introduction

Let G be an additive finite abelian group. A finite sequence $S = (g_1, g_2, \cdots, g_l) = g_1 g_2 \cdots g_l$ of elements of G, where repetition of elements is allowed and their order is disregarded, is called a zero-sum sequence if $g_1 + g_2 + \cdots + g_l = 0$.

For a finite abelian group G of cardinality n, the Davenport constant $D(G)$ is the smallest natural number t such that any sequence of t elements in G has a non-empty zero-sum subsequence. Another interesting constant, $E(G)$, is the smallest natural number k such that any sequence of k elements in G has a zero-sum subsequence of length n.

For the particular group $\mathbb{Z}_n$, the following generalization of $E(G)$ has been considered in [1] and [2] recently. Let $n \in \mathbb{N}$ and assume $A \subseteq \mathbb{Z}_n$. Then $E_A(n)$ is the least $t \in \mathbb{N}$ such that for all sequences $(x_1, \ldots, x_t) \in \mathbb{Z}^t$, there exist indices $j_1, \ldots, j_n \in \mathbb{N}, 1 \leq j_1 <$

$\cdots < j_n \leq t$ and $(\vartheta_1, \cdots, \vartheta_n) \in A^n$ with

$$\sum_{i=1}^{n} \vartheta_i x_{j_i} \equiv 0 \pmod{n}.$$

To avoid trivial cases, one assumes that the weight set A does not contain 0 and it is non–empty.

Similarly, for any such set $A \subseteq \mathbb{Z}_n \setminus \{0\}$ of weights, we define the Davenport constant of $\mathbb{Z}_n$ with weight A, denoted by $D_A(n)$, as the least natural number k such that for any sequence $(x_1, \cdots, x_k) \in \mathbb{Z}^k$, there exists a non-empty subsequence $(x_{j_1}, \cdots, x_{j_l})$ and $(a_1, \cdots, a_l) \in A^l$ such that

$$\sum_{i=1}^{l} a_i x_{j_i} \equiv 0 \pmod{n}.$$

Thus, for the group $G = \mathbb{Z}_n$, if we take $A = \{1\}$, then $E_A(n)$ and $D_A(n)$ are, respectively, $E(G)$ and $D(G)$ as defined earlier.

$E_A(n)$ and $D_A(n)$ were studied in [1], [2] and [3].

It is not difficult to observe the following result.

Lemma 1 *$D_A(n) + n - 1 \leq E_A(n) \leq 2n - 1$ for any $A \subseteq \mathbb{Z}_n \setminus \{0\}$.*

Lemma 2 ([2]) *Let $A = \mathbb{Z}_n \setminus \{0\}$. Then $E_A(n) = n + 1$.*

In [2] Adhikari et al proposed characterizing any other sets for which $E_A(n) = n + 1$ or even those for which $E_A(n) = n + j$ for specific small values of j. It is easy to see that if $A \subseteq B$, then $D_A(n) \geq D_B(n)$.

In this paper we prove the following results:

Theorem 1 *Let n be a positive integer and p be a prime satisfying $p^k \| n$. If $A = \{a \mid a \not\equiv 0 \pmod{p}\}$, then $D_A(n) = k + 1$ and $E_A(n) = n + k$.*

Theorem 2 *If A is an arithmetic progression with length $l = \lceil \frac{n}{2} \rceil$, where for any real number x, $\lceil x \rceil$ denotes the smallest integer $\geq x$, and common difference 1, that is, A is the set of the form $\{a + i | i = 1, 2, \cdots, l\}$ where $1 \leq a < a + l \leq n - 1$, then $D_A(n) = 2$, $E_A(n) = n + 1$.*

2. Proofs of Theorem 1 and Theorem 2

In order to prove the theorems, we need the following result.

Lemma 3 ([4]) *Let A, B be subsets of a finite group G such that $|A| + |B| \geq |G| + 1$. Then $A + B = G$.*

Proof of Theorem 1. (1) We will prove that $D_A(n) = k + 1$.

First, we prove that $D_A(n) > k$. We assert that $0 \notin \sum_{i\in I} \frac{n}{p^i}A$ with $I \subseteq \{1, 2, \cdots, k\}$. We proceed by induction on the cardinality of I. Note that for $|I| = 1$, the result follows trivially. Inductively, assume the result holds true for $1 \leq |I| < k$. Now consider $|I| = k$. If $0 \in \sum_{i=1}^{k} \frac{n}{p^i}A$, then there must exist $a_i \in A$ for $i = 1, 2, \cdots, k$ such that

$$\frac{n}{p}a_1 + \frac{n}{p^2}a_2 + \cdots + \frac{n}{p^k}a_k \equiv 0 \pmod{n}.$$

Multiplying the above equation by p, we get

$$\frac{n}{p}a_2 + \frac{n}{p^2}a_3 + \cdots + \frac{n}{p^{k-1}}a_k \equiv 0 \pmod{n}.$$

Hence $0 \in \frac{n}{p}A + \frac{n}{p^2}A + \cdots + \frac{n}{p^{k-1}}A$, which contradicts to the inductive hypothesis.

Next we prove that $D_A(n) \leq k + 1$. Let $S = (s_1, \cdots, s_N)$ be a sequence of elements in $\mathbb{Z}_n$ of length $N = k+1$. We will prove that S has a zero-sum subsequence with weight A. We distinguish two cases:

Case 1. If there exist two elements s_1 and s_2 such that $p^i \parallel s_1$, $p^i \parallel s_2$ for some $i = 0, 1, \cdots, k-1$, then $\frac{s_2}{p^i}$, $n - \frac{s_1}{p^i} \in A$. Hence,

$$s_1\frac{s_2}{p^i} + s_2(n - \frac{s_1}{p^i}) \equiv 0 \pmod{n}.$$

Case 2. If Case 1 does not hold, then there must exist one element, say s_{i_0}, satisfying $p^k \| s_{i_0}$. Since $\frac{n}{p^k} \in A$, we have

$$s_{i_0}\frac{n}{p^k} \equiv 0 \pmod{n}.$$

Thus, we have proved that $D_A(n) = k + 1$.

(2) We will prove that $E_A(n) = n + k$. Assume that $S = (s_1, \cdots, s_{N'})$ is a sequence of elements in $\mathbb{Z}_n$ of length $N' = n + k$. To prove $E_A(n) = n + k$, because of Lemma 1 it suffices to prove that S has a zero-sum subsequence of length n with weight A. We partition S into the following multisets (sets with repetitions allowed).

$$M_i = \big\{s_j |\ p^i \parallel s_j,\ s_j \in S\big\},\ \text{for } i = 0, 1, 2, \cdots, k.$$

Note that every pair of elements $s_i^{(1)}, s_i^{(2)}$ in M_i constitutes a zero-sum subsequence of S with weight A since

$$s_i^{(1)} \cdot \frac{s_i^{(2)}}{p^i} + s_i^{(2)}\big(n - \frac{s_i^{(1)}}{p^i}\big) \equiv 0 \pmod{n},$$

where $\frac{s_i^{(2)}}{p^i},\ n - \frac{s_i^{(1)}}{p^i} \in A,$ for $i = 0, 1, \cdots, k-1$.

Since every element $s_k^{'}$ in M_k produces a zero-sum subsequence of S of length 1 with weight A since $s_k^{'}\frac{n}{p^k} \equiv 0 \pmod{n},\ \frac{n}{p^k} \in A$. We consider two cases:

Case 1. n is even. We can choose m $(0 \le m \le k)$ integers $l_1, l_2, \cdots, l_m$ satisfying $l_1 + l_2 + \cdots + l_m = \frac{n-t}{2}$, where $t = |M_k|$, $l_1 = [\frac{|M_{i_1}|}{2}], l_2 = [\frac{|M_{i_2}|}{2}], \cdots, l_m = [\frac{|M_{i_m}|}{2}]$, $0 \le i_1 \le i_2 \le i_m \le k-1$. Hence, we can obtain $\frac{n-t}{2}$ pairs of disjoint zero-sum subsequences of S with weight A and t disjoint zero-sum subsequences of S of length 1 with weight A, and it follows that we can get a zero-sum subsequence of S of length n with weight A.

Case 2. n is odd. If n is a prime, the result follows because of Lemma 2. For $n > 3$ and composite, since $2(k+1) < n$ we know that there must exist some M_i $(0 \le i \le k)$ satisfying $|M_i| \ge 3$. Let $s_i^{(1)}, s_i^{(2)}, s_i^{(3)} \in M_i$. We conclude that there must exist $x, y, z \in A$ satisfying $xs_i^{(1)} + ys_i^{(2)} + zs_i^{(3)} \equiv 0 \pmod{n}$.

Indeed, choose $x = 1,\ y = 1$ if $\frac{s_i^{(1)}}{p^i} + \frac{s_i^{(2)}}{p^i} \not\equiv 0 \pmod{p}$, and $x = 1,\ y = n-1$ if $\frac{s_i^{(1)}}{p^i} + \frac{s_i^{(2)}}{p^i} \equiv 0 \pmod{p}$. Then the equation $\frac{s_i^{(3)}}{p_i}z = -(x\frac{s_i^{(1)}}{p_i} + y\frac{s_i^{(2)}}{p_i}) \pmod{\frac{n}{p^i}}$ has a solution in A, and the result follows as before. □

Corollary 1 *Let n be a positive integer and p be a prime satisfying $p\|n$. If $A = \{a \mid a \not\equiv 0 \pmod{p}\}$, then $D_A(n) = 2$ and $E_A(n) = n+1$.*

Proof of Theorem 2. (1) We prove that $D_A(n) = 2$. Let $S = (s_1, s_2)$ be a sequence of elements in $\mathbb{Z}_n$ of length 2. It suffices to show that S has a zero-sum subsequence with weight A. We distinguish two cases:

Case 1. n is even. We see that $\frac{n}{2} \in A$ since $|A| = \lceil \frac{n}{2} \rceil$. So if $2|s_1$ or $2|s_2$, then $s_1\frac{n}{2} \equiv 0 \pmod{n}$ or $s_2\frac{n}{2} \equiv 0 \pmod{n}$. If s_1 and s_2 are both odd, then $\frac{n}{2} \in s_1A, s_2A$. Therefore, $0 \in s_1A + s_2A$.

Case 2. n is odd. If $\gcd(s_1, n) = \gcd(s_2, n) = 1$, then $|s_1A| = |s_2A| = \frac{n+1}{2}$. Hence $|s_1A| + |s_2A| = 2\frac{n+1}{2} = n+1 > n$. From Lemma 3 it follows that $s_1A + s_2A = \mathbb{Z}_n$. Therefore, $0 \in s_1A + s_2A$. If $\gcd(s_1, n) = d \ge 1$, that is $3 \le d \le \frac{n}{3}$, then there must exist $i (1 \le i \le d-1)$ such that $\frac{in}{d} \in A$ since $|A| = \frac{n+1}{2}$. It follows that $s_1\frac{in}{d} \equiv 0 \pmod{n}$.

(2). We now prove that $E_A(n) = n+1$. Assume that $S = (s_1, \cdots, s_N)$ is a sequence of elements in $\mathbb{Z}_n$ of length $N = n+1$. To prove $E_A(n) = n+1$, because of Lemma 1 it suffices to prove that S has a zero-sum subsequence of length n with weight A.

Partition S into the two multi-sets M_1, M_2 where $M_1 = \{s_i|\ \gcd(s_i, n) = 1, s_i \in S\}$ and $M_2 = \{s_i|\ \gcd(\mathrm{s_i}, \mathrm{n}) \neq 1, \mathrm{s_i} \in \mathrm{S}\}$. We note the two following facts.

Fact 1. If $s_1, s_2 \in M_1$, then $0 \notin s_1A, s_2A$. Thus, from $D_A(n) = 2$, we conclude that there exist two elements $a_1, a_2 \in A$ such that $a_1s_1 + a_2s_2 \equiv 0 \pmod{n}$.

Fact 2. If $s_{i_0} \in M_2$, then there must exist one element $a_0 \in A$ such that $a_0s_i \equiv 0 \pmod{n}$. We distinguish two cases:

Case 1. n is even.

(1) $|M_1| \geq n$. Using Fact 1, it is easy to see that we can get a zero-sum subsequence of length n with weight A.

(2) $|M_2| < n$. Using Fact 1 and Fact 2, it is easy to see that we can get a zero-sum subsequence of length n with weight A.

Case 2. n is odd.

(1) $|M_2| \geq 1$. Using Fact 1 and Fact 2, it is easy to see that we can get a zero-sum subsequence of length n with weight A.

(2) $M_2 = \emptyset$. Then $|M_1| = n + 1$ and $\gcd(s_i, n) = 1$ for $i = 1, 2, \cdots, n + 1$. Set $A_i = s_iA$. Therefore, $|A_i| = \lceil \frac{n}{2} \rceil = \frac{n+1}{2}$. Since $|A_1| + |A_2| > n$, the result follows that $\sum_{i=1}^{n} A_i \supseteq A_1 + A_2 = \mathbb{Z}_n$. Therefore, $0 \in \sum_{i=1}^{n} A_i = \mathbb{Z}_n$. □

Acknowledgement. The author would like to thank the anonymous referee for his or her helpful comments which have greatly improved the presentation of the paper. He is very appreciative and sincerely thanks the referee.

References

[1] Sukumar Das Adhikari and Purusottam Rath, *Davenport constant with weights and some related questions,* Integers **6** (2006), #A30.

[2] S. D. Adhikari, Y. G. Chen, J. B. Friedlander, S. V. Konyagin and F. Pappalardi, *Contributions to zero–sum problems*, Discrete Math. **306**, 1–10 (2006).

[3] Florian Luca, *A generalization of a classical zero-sum problem*, Discrete Math. **307**, 1672–1678 (2007).

[4] H. B. Mann, *Addition Theorems,* Wiley, New York (1965).