INTEGERS: ELECTRONIC JOURNAL OF COMBINATORIAL NUMBER THEORY 8 (2008), #A26

A MULTIPLICITY PROBLEM RELATED TO SCHUR NUMBERS

Daniel Schaal
Department of Mathematics and Statistics, South Dakota State University, Brookings, SD 57007, USA
daniel.schaal@sdstate.edu

Hunter Snevily
Department of Mathematics, University of Idaho, Moscow, ID 43844, USA
snevily@uidaho.edu

Received: 9/18/06, Revised: 6/5/08, Accepted: 6/6/08, Published: 6/30/08

Abstract

For each natural number n, let C_n represent the set of all 2-colorings of the set $\{1, 2, \ldots, n\}$. Given a natural number n and a coloring $\Delta \in C_n$, let $S(\Delta)$ represent the set

$$S(\Delta) = \{x_3 \mid \exists\ x_1, x_2\ s.t.\ x_1 + x_2 = x_3\ \ and\ \Delta(x_1) = \Delta(x_2) = \Delta(x_3)\}.$$

Given a natural number n, let

$$f(n) = \min_{\Delta \in C_n} S(\Delta).$$

For all natural numbers n and r where $\frac{n}{2} \leq r \leq n$, let $C_{n,r}$ represent the set of all 2-colorings of the set $\{1, 2, \ldots, n\}$ where $\max\{|\Delta^{-1}(0)|, |\Delta^{-1}(1)|\} = r$. Given natural numbers n and r where $\frac{n}{2} \leq r \leq n$, let

$$f(n, r) = \min_{\Delta \in C_{n,r}} S(\Delta).$$

In this paper it is determined that for all natural numbers n,

$$f(n) = \begin{cases} 0 & 1 \leq n \leq 4 \\ \left\lfloor \frac{n-3}{2} \right\rfloor & n \geq 5 \end{cases}$$

and for all natural numbers n and r where $n \geq 5$ and $\frac{n}{2} \leq r \leq n$,

$$f(n, r) = \begin{cases} r - 2 & r < n \\ r - 1 & r = n. \end{cases}$$

1. Introduction

Let $\mathbb{N}$ represent the set of natural numbers and let $[a, b]$ denote the set $\{n \in \mathbb{N} | \, a \leq n \leq b\}$. A function $\Delta : [1, n] \to [0, t-1]$ is referred to as a *t-coloring of the set* $[1, n]$ or as a *t-coloring*

of length n. For every natural number n, let C_n represent the set of all 2-colorings of length n. For a given natural number n' where $n' < n$, a coloring Δ restricted to the set $[1, n']$ will be denoted by $\Delta|_{n'}$. Given a t-coloring Δ and a linear equation L in m variables, a solution $(x_1, x_2, \ldots, x_m)$ to L is *monochromatic* if and only if $\Delta(x_1) = \Delta(x_2) = \cdots = \Delta(x_m)$.

In 1916, I. Schur [21] proved that for every $t \geq 2$, there exists a least integer $n = Schur(t)$ such that for every t-coloring of length n, there exists a monochromatic solution to

$$x_1 + x_2 = x_3. \tag{1}$$

Note that the integers x_1 and x_2 need not be distinct. The integers $Schur(t)$ are called *Schur numbers*. It is known that $Schur(2) = 5$, $Schur(3) = 14$ and $Schur(4) = 45$, but no other Schur numbers are known [22]. A monochromatic solution to equation (1) is called a *monochromatic Schur triple*. In 1933, R. Rado found necessary and sufficient conditions to determine if an arbitrary system of linear equations admits a monochromatic solution for every coloring of the natural numbers with a finite number of colors [5, 15, 16, 17].

Recently, several other problems related to Schur numbers have been considered [1, 2, 3, 4, 7, 8, 9, 10, 11, 12, 13, 14]. In 1996, R. Graham, V. Rödl and A. Rucinski proposed the following problem [6]. Find (asymptotically) the least number of monochromatic Schur triples that must occur in an arbitrary 2-coloring of length n. A problem of this nature where the number of monochromatic solutions is to be determined is referred to as a *multiplicity* problem. This problem was solved independently by A. Robertson and D. Zeilberger [19] and by T. Schoen [20] and was found to be $\frac{1}{22}n^2 + O(n)$.

In this paper we modify the problem of Graham, Rödl and Rucinski by asking the following question. In an arbitrary 2-coloring of length n, how many integers must there be that are the third integer (i.e. x_3) in at least one monochromatic Schur triple? For convenience in this paper, we shall refer to the third integer in any monochromatic Schur triple as *special*, formally defined below.

Definition 1. *For every 2-coloring $\Delta : [1, n] \to [0, 1]$, an integer $s \in [1, n]$ is* special *if and only if there exist integers $x_1, x_2 \in [1, n]$, such that $x_1 + x_2 = s$ and $\Delta(x_1) = \Delta(x_2) = \Delta(s)$. The set of all special integers for a 2-coloring Δ is denoted by $S(\Delta)$.*

Definition 2. *For every natural number n, let*

$$f(n) = \min_{\Delta \in C_n} S(\Delta).$$

In this paper it is determined that

$$f(n) = \begin{cases} 0 & 1 \leq n \leq 4 \\ \left\lfloor \frac{n-3}{2} \right\rfloor & n \geq 5. \end{cases}$$

More importantly, and perhaps more surprisingly, it is found that the number of special integers in a 2-coloring Δ is more directly related to the number of monochromatic integers

in Δ, rather than the length of Δ. For instance, the number of special integers that must occur in an arbitrary 2-coloring of length seventy with fourty integers colored 0 is the same as the number that must occur in an arbitrary 2-coloring of length sixty with fourty integers colored 0 or an arbitrary 2-coloring of length fifty with fourty integers colored 0. We say that a coloring Δ has r monochromatic integers if $\max\{|\Delta^{-1}(0)|, |\Delta^{-1}(1)|\} = r$. For all natural numbers n and r where $\frac{n}{2} \leq r \leq n$, let $C_{n,r}$ represent the set of all 2-colorings of length n with r monochromatic integers.

Definition 3. *For all integers n and r where $n \geq 5$ and $\frac{n}{2} \leq r \leq n$, let*

$$f(n,r) = \min_{\Delta \in C_{n,r}} S(\Delta).$$

In this paper it is determined that

$$f(n,r) = \begin{cases} r-2 & r < n \\ r-1 & r = n. \end{cases}$$

The above formula for $f(n)$ follows immediately from this result.

2. Main Results

Theorem 1. *If n and r are natural numbers such that $n \geq 5$ and $\frac{n}{2} \leq r \leq n$, then*

$$f(n,r) = \begin{cases} r-2 & \text{if } r < n \\ r-1 & \text{if } r = n. \end{cases}$$

Proof. First we shall prove the case where $r = n$. Let a natural number $n \geq 5$ be given, let $r = n$ and let $\Delta : [1,n] \to [0,1]$ be a coloring with r monochromatic integers. Clearly $S(\Delta) = [2,n]$, so $f(n,r) = r-1$.

We shall now consider the case where $r < n$. First we shall show that $f(n,r) \leq r-2$. Let integers n and r be given such that $n \geq 5$ and $\frac{n}{2} \leq r < n$. We shall exhibit a coloring $\Delta : [1,n] \to [0,1]$ where $|\Delta^{-1}(0)| = r$ and $|S(\Delta)| = r-2$. If $r = n-1$ let Δ be defined by

$$\Delta(x) = \begin{cases} 1 & x = 1 \\ 0 & 2 \leq x \leq n. \end{cases}$$

It is clear that $S(\Delta) = [4,n]$ and that $|S(\Delta)| = n-3 = r-2$. If $r \leq n-2$, let Δ be defined by

$$\Delta(x) = \begin{cases} 1 & 1 \leq x \leq n-r-1 \\ 0 & n-r \leq x \leq n-1 \\ 1 & x = n. \end{cases}$$

If $r = n-2$, then $\Delta(S) = [4, n-1]$; if $r = \frac{n}{2}$, then $\Delta(S) = [2, n-r-1]$; and if $\frac{n}{2} < r < n-2$, then $\Delta(S) = [2, n-r-1] \cup [2n-2r, n-1]$. Since in each case $|S(\Delta)| = r-2$, it follows that $f(n,r) \leq r-2$.

Next we shall show that $f(n,r) \geq r-2$ for all integers n and r where $n \geq 5$ and $\frac{n}{2} \leq r < n$. We will use induction on the integer n. When $n = 5$ we must consider the two cases of $r = 3$ and $r = 4$. Since the 2-color Schur number is 5, every 2-coloring of the set $[1,5]$ with 3 monochromatic integers has at least 1 special integer, so $f(5,3) \geq 1$. It is easy to check that every 2-coloring of the set $[1,5]$ with 4 monochromatic integers has as least 2 special integers, so $f(5,4) \geq 2$. In both cases $f(5,r) \geq r-2$, so the basis step is complete. Now, let an integer $n_0 \geq 6$ be given. We may assume that

$$f(n,r) \geq r-2 \text{ for every } n \in [5, n_0-1] \text{ and for every } r \text{ such that } \frac{n}{2} \leq r < n. \qquad (2)$$

We must show that $f(n_0,r) \geq r-2$ for all integers r such that $\frac{n_0}{2} \leq r < n_0$. Let an integer r such that $\frac{n_0}{2} \leq r < n_0$ be given and let a coloring $\Delta : [1,n_0] \to [0,1]$ with r monochromatic integers be given. We must show that $|S(\Delta)| \geq r-2$.

Without loss of generality, we may assume that Δ has r integers colored 0. If $\Delta(n_0) = 1$, then the coloring $\Delta|_{n_0-1}$ has r integers colored 0. If $r \leq n_0 - 2$, then by (2) it follows that $\Delta|_{n_0-1}$ has at least $r-2$ special integers, so Δ does as well. If $r = n_0 - 1$ it has been previously shown that $\Delta|_{n_0-1}$ has $r-1$ special integers. In each case $|S(\Delta)| \geq r-2$, so we may assume that $\Delta(n_0) = 0$.

We shall consider three cases.

Case 1: Assume that $r > \frac{n_0+1}{2}$.

Since Δ has r integers colored 0 and $\Delta(n_0) = 0$, it follows that $\Delta|_{n_0-1}$ has $r-1$ integers colored 0. Hence, from (2) we have that

$$|S(\Delta|_{n_0-1})| \geq r-3,$$

so it suffices to show that n_0 is special. We shall consider the two subcases where n_0 is even and where n_0 is odd.

(i) Assume that n_0 is even. It follows that $r > \frac{n_0}{2}$.

If $\Delta\left(\frac{n_0}{2}\right) = 0$, then since $\Delta(n_0) = 0$ and $\frac{n_0}{2} + \frac{n_0}{2} = n_0$, it follows that n_0 is special. In this case we are done, so we may assume that $\Delta\left(\frac{n_0}{2}\right) = 1$.

Let $A = \left[1, \frac{n_0}{2} - 1\right] \cup \left[\frac{n_0}{2} + 1, n_0 - 1\right]$. Since there are $r-1$ integers colored 0 in the set A and $r > \frac{n_0}{2}$, there are at least $\frac{n_0}{2}$ integers in the set A colored 0. For every $i \in \left[1, \frac{n_0}{2} - 1\right]$ let $A_i = \{i, n_0 - i\}$. Hence the set $\left\{A_1, A_2, \ldots, A_{\frac{n_0}{2}-1}\right\}$ is a partition of the set A into $\frac{n_0}{2} - 1$ subsets and there exists an $i \in \left[1, \frac{n_0}{2} - 1\right]$ such that $\Delta(i) = \Delta(n_0 - i) = 0$. Then $(i, n_0 - 1, n_0)$ is a monochromatic Schur triple, so n_0 is special.

(ii) Assume that n_0 is odd.

This subcase is very similar to the n_0 even subcase. Let $A = [1, n_0 - 1]$ and for every $i \in [1, \frac{n_0-1}{2}]$, let $A_i = \{i, n_0 - i\}$. Since the set A has $r-1$ integers colored 0 and $r - 1 \geq \frac{n_0+1}{2}$,

there exists an $i \in [1, \frac{n_0-1}{2}]$ such that $(i, n_0 - i, n_0)$ is a monochromatic Schur triple, so n_0 is special.

Since in both subcases n_0 is special and it was previously shown that $|S(\Delta|_{n_0-1})| \geq r-3$, we have that $|S(\Delta)| \geq r-2$.

Case 2: Assume that n_0 is even and $r = \frac{n_0}{2}$.

It follows that $\Delta|_{n_0-1}$ has $r-1$ integers colored 0 and r integers colored 1. From (2) it follows that $\Delta|_{n_0-1}$ has at least $r-2$ special integers, so Δ does as well and $|S(\Delta)| \geq r-2$.

Case 3: Assume that n_0 is odd and $r = \frac{n_0+1}{2}$.

Since $n_0 - 2$ is odd, it follows that $\Delta|_{n_0-2}$ has $\frac{n_0-1}{2} = r-1$ monochromatic integers. From (2) it follows that $\Delta|_{n_0-2}$ has at least $r-3$ special integers, so it will be sufficient to show that $S(\Delta) \cap \{n_0 - 1, n_0\} \neq \emptyset$.

Recall that Δ has $r = \frac{n_0+1}{2}$ integers colored 0 and $\Delta(n_0) = 0$. Hence, exactly $\frac{n_0-1}{2}$ integers in the set $[1, n_0 - 1]$ are colored 0. Let $A = [1, n_0 - 1]$ and for every $i \in \left[1, \frac{n_0-1}{2}\right]$ let $A_i = \{i, n_0 - i\}$. Hence the set $\left\{A_1, A_2, \ldots, A_{\frac{n_0-1}{2}}\right\}$ is a partition of the set A into $\frac{n_0-1}{2}$ subsets. If, for any $i \in \left[1, \frac{n_0-1}{2}\right]$, the set A_i is monochromatic in 0, then $(i, n_0 - i, n_0)$ is a monochromatic Schur triple and n_0 is special. In this case we are done, so we may assume that, for every $i \in \left[1, \frac{n_0-1}{2}\right]$, A_i contains one integer colored 0 and one integer colored 1.

Now we will prove the following claim.

Claim. *Let $\Delta(n_0 - 1) = a$ and let $b = 1 - a$. If $S(\Delta) \cap \{n_0 - 1, n_0\} = \emptyset$, then $\Delta(x) = b$ for every $x \in \left[1, \frac{n_0-1}{2}\right]$.*

Proof. Assume that the hypothesis of the claim is satisfied and note that $\{a, b\} = \{0, 1\}$. We shall show that $\Delta\left(\frac{n_0+1}{2} - i\right) = b$ for every $i \in \left[1, \frac{n_0-1}{2}\right]$ by using induction on i. First we shall show that

$$\Delta\left(\frac{n_0+1}{2} - 1\right) = b.$$

If $\Delta\left(\frac{n_0+1}{2} - 1\right) = a$, then $\left(\frac{n_0+1}{2} - 1, \frac{n_0+1}{2} - 1, n_0 - 1\right)$ is a monochromatic Schur triple and $n_0 - 1 \in S(\Delta)$, which is a contradiction. Hence, we may assume that $\Delta\left(\frac{n_0+1}{2} - 1\right) = b$.

Now, let $i_0 \in \left[2, \frac{n_0-1}{2}\right]$ be given and assume that $\Delta\left(\frac{n_0+1}{2} - (i_0 - 1)\right) = b$. We will show that

$$\Delta\left(\frac{n_0+1}{2} - i_0\right) = b.$$

Now, $A_{\frac{n_0+3}{2}-i_0} = \left\{\frac{n_0+1}{2} - (i_0 - 1), \frac{n_0-3}{2} + i_0\right\}$. Since $A_{\frac{n_0+3}{2}-i_0}$ contains an integer colored a, it follows that

$$\Delta\left(\frac{n_0-3}{2} + i_0\right) = a.$$

If $\Delta\left(\frac{n_0+1}{2} - i_0\right) = a$, then $\left(\frac{n_0-3}{2} + i_0, \frac{n_0+1}{2} - i_0, n_0 - 1\right)$ is a monochromatic Schur triple and $n_0 - 1 \in S_3(\Delta)$, which is a contradiction. Hence, we may assume that $\Delta\left(\frac{n_0+1}{2} - i_0\right) = b$ and the proof of the claim is complete. □

Now, as noted above, if $S(\Delta) \cap \{n_0 - 1, n_0\} \neq \emptyset$, then $|S(\Delta)| \geq r - 2$. If $S(\Delta) \cap \{n_0 - 1, n_0\} = \emptyset$, then from the claim we have that $\Delta(x) = b$ for every $x \in \left[1, \frac{n_0-1}{2}\right]$. This implies that $\left[2, \frac{n_0-1}{2}\right] \subseteq S(\Delta)$ and that $|S(\Delta)| \geq \frac{n_0-3}{2} = r - 2$. In either case we have that $|S(\Delta)| \geq r - 2$, so we are done with Case 3.

Since in all three cases we showed that $|S(\Delta)| \geq r - 2$, the proof of Theorem 1 is complete. □

We are now ready to state and prove Theorem 2. The theorem follows directly from previous results.

Theorem 2. *For every natural number n,*

$$f(n) = \begin{cases} 0 & 1 \leq n \leq 4 \\ \left\lfloor \frac{n-3}{2} \right\rfloor & n \geq 5. \end{cases}$$

Proof. The case where $1 \leq n \leq 4$ follows directly from the fact that the 2-color Schur number is 5. If $n \geq 5$, then every coloring of the set $[1, n]$ contains at least $\left\lfloor \frac{n+1}{2} \right\rfloor$ monochromatic integers. From Theorem 1 it follows that every coloring of the set $[1, n]$ contains at least $\left\lfloor \frac{n+1}{2} \right\rfloor - 2 = \left\lfloor \frac{n-3}{2} \right\rfloor$ special integers. □

Acknowledgments

This material is based upon work supported by a South Dakota Governor's 2010 Individual Research Seed Grant.

References

[1] A. Beutelspacher, W. Brestovansky, Generalized Schur numbers, Lecture Notes in Mathematics, Springer-Verlag, Berlin, 969 (1982), 30-38.

[2] A. Bialostocki, G. Bialostocki, D. Schaal, A zero-sum theorem, Journal of Combinatorial Theory Series A, 101 (2003), no. 1, 147-152.

[3] A. Bialostocki, H. Lefmann, T. Meerdink, On the degree of regularity of some equations. Selected papers in honour of Paul Erdös on the occasion of his 80th birthday (Keszthely, 1993), Discrete Mathematics, 150 (1996), no. 1-3, 49-60.

[4] A. Bialostocki, D. Schaal, On a Variation of Schur Numbers, Graphs and Combinatorics, vol. 16 (2000), 139-147.

[5] W. Deuber, Developments Based on Rado's Dissertation "Studien zur Kombinatorik," Survey in Combinatorics (1989), 52-74, Cambridge University Press.

[6] R. Graham, V. Rödl, A. Rucinski, On Schur properties of random subsets of integers. Journal of Number Theory, 61 (1996), no. 2, 388-408.

[7] H. Harborth, S. Maasberg, Rado numbers for Fibonacci sequences and a problem of S. Rabinowitz, in: G. E. Bergum at al., eds., Applications of Fibonacci Numbers, Vol. 6 (Cluwer Acad. Publ.) 143-153.

[8] H. Harborth, S. Maasberg, Rado numbers for homogeneous second order linear recurrences - degree of partition regularity, Congressus Numerantium, Vol. 108 (1995) 109-118.

[9] H. Harborth, S. Maasberg, Rado numbers for $a(x+y)=bz$, Journal of Combinatorial Theory Series A, Vol. 80, num. 2 (1997) 356-363.

[10] H. Harborth, S. Maasberg, All two-color Rado numbers for $a(x+y)=bz$, Discrete Mathematics, 197/198 (1999), 397-407.

[11] B. Hopkins, D. Schaal, On Rado numbers for $\sum_{i=1}^{m-1} a_i x_i = x_m$, Advances in Applied Mathematics, 35 (2005), no. 4, 433-441.

[12] B. Johnson, D. Schaal, Disjunctive Rado numbers, Journal of Combinatorial Theory Series A, 112 (2005), no. 2, 263-276.

[13] S. Jones, D. Schaal, A class of two-color Rado numbers, Discrete Mathematics, 289 (2004) no. 1-3, 63-69.

[14] W. Kosek, D. Schaal, Rado Numbers for the equation $\sum_{i=1}^{m-1} x_i + c = x_m$ for negative values of c, Advances in Applied Mathematics, vol. 27 (2001) 805-815.

[15] R. Rado, Verallgemeinerung eines Satzes von van der Waerden mit Anwendungen auf ein Problem der Zahlentheorie, Sonderausg. Sitzungsber. Preuss. Akad. Wiss. Phys.- Math. Klasse, 17 (1933) 1-10.

[16] R. Rado, Studien zur Kombinatorik, Math. Z. 36 (1933), 242-280.

[17] R. Rado, Note on Combinatorial Analysis, Proc. London Math. Soc. 48 (1936) 122-160.

[18] A. Robertson, D. Schaal, Off-Diagonal Generalized Schur Numbers, Advances in Applied Mathematics, vol. 26 (2001) 252-257.

[19] A. Robertson, D. Zeilberger, A 2-coloring of $[1,N]$ can have $(1/22)N^2+O(N)$ monochromatic Schur triples, but not less, Electronic Journal of Combinatorics, 5 (1998), R19, 4 pp.

[20] T. Schoen, The number of monochromatic Schur triples. European Journal of Combinatorics, 20 (1999), no. 8, 855-866.

[21] I. Schur, Über die Kongruenz $x^m + y^m \equiv z^m (\mathrm{mod}\ p)$. Jahresber. Deutsch. Math. Verein. 25 (1916), 114-117.

[22] W. Wallis, A. Street, J. Wallis, Combinatorics: Room Squares, Sum-free Sets, Hadamard Matrices, Lecture Notes in Math., vol. 292, Springer-Verlag, Berlin and New York, 1972.