Journal of Integer Sequences, Vol. 19 (2016),
Article 16.5.3

# A Proof of Dixon's Identity

Jovan Mikić
J.U. SŠC "Jovan Cvijić"
74480 Modriča
Bosnia and Herzegovina
jnmikic@gmail.com

**Abstract**

We give a proof of Dixon's binomial coefficient identity using recurrence equations and induction.

## 1 Introduction

In 1912, Dixon [1] established the following famous identity

$$\sum_{k=-a}^{a} (-1)^k \binom{a+b}{a+k} \binom{b+c}{b+k} \binom{c+a}{c+k} = \frac{(a+b+c)!}{a! \cdot b! \cdot c!}, \tag{1}$$

where $a$, $b$, $c$ are nonnegative integers.

There are many proofs of Eq. (1). In 1916, MacMahon proved his master theorem. In 1962, Good [4] found a short proof of Eq. (1) using MacMahon's master theorem. Gessel and Stanton [3] gave a short proof using Laurent series in 1985. In 1990, Ekhad [2] gave a very short proof using induction. In 2003, Guo [5] gave a short proof using polynomials.

We give an elementary proof of Eq. (1) using recurrence equations and induction. In order to obtain recurrence equations, we will use some so-called auxiliary sums.

## 2 Proof of Eq. (1)

*Proof.* Let $\mathbb{N}$ denote the set of positive integers, and let $\mathbb{N}_0$ denote the set of nonnegative integers. We let $S(a, b, c)$ denote the left side of Eq. (1), where $a, b, c \in \mathbb{N}_0$. We introduce the auxiliary sums $P(a,b,c)$, $Q(a,b,c)$ and $R(a,b,c)$ as follows

$$P(a,b,c) = \sum_{k=-a}^{a} (-1)^k (a^2-k^2) \binom{a+b}{a+k}\binom{b+c}{b+k}\binom{c+a}{c+k}, \tag{2}$$

$$Q(a,b,c) = \sum_{k=-a}^{a} (-1)^k (b^2-k^2) \binom{a+b}{a+k}\binom{b+c}{b+k}\binom{c+a}{c+k}, \tag{3}$$

$$R(a,b,c) = \sum_{k=-a}^{a} (-1)^k (c^2-k^2) \binom{a+b}{a+k}\binom{b+c}{b+k}\binom{c+a}{c+k}. \tag{4}$$

Let $a,b,c \in \mathbb{N}$. We will use the well-known binomial identity $k\binom{n}{k} = n\binom{n-1}{k-1}$. Then

$$\begin{aligned} P(a,b,c) &= \sum_{k=-a+1}^{a-1} (-1)^k (a+k)\binom{a+b}{a+k}\binom{b+c}{b+k}(a-k)\binom{c+a}{a-k} \\ &= \sum_{k=-a+1}^{a-1} (-1)^k (a+b)\binom{a+b-1}{a+k-1}\binom{b+c}{b+k}(c+a)\binom{c+a-1}{a-k-1} \\ &= (a+b)(a+c)\sum_{k=-a+1}^{a-1} (-1)^k \binom{a-1+b}{a-1+k}\binom{b+c}{b+k}\binom{c+a-1}{c+k}. \end{aligned}$$

From the last equation above, it follows that

$$P(a,b,c) = (a+b)(a+c)S(a-1,b,c). \tag{5}$$

Similarly, we obtain that

$$Q(a,b,c) = (a+b)(b+c)S(a,b-1,c), \tag{6}$$

and

$$R(a,b,c) = (a+c)(b+c)S(a,b,c-1). \tag{7}$$

From Eqns. (2) and (3), we have that

$$P(a,b,c) - Q(a,b,c) = (a^2-b^2)S(a,b,c). \tag{8}$$

Let $a \neq b$. From Eqns. (5), (6) and (8), it follows that

$$S(a,b,c) = \frac{1}{a-b}\big((a+c)S(a-1,b,c) - (b+c)S(a,b-1,c)\big). \tag{9}$$

Similarly, if $a \neq c$, then it follows that

$$S(a,b,c) = \frac{1}{a-c}\big((a+b)S(a-1,b,c) - (b+c)S(a,b,c-1)\big). \tag{10}$$

We treat the case when $a = b = c$ separately.

$$\begin{aligned}
S(a,a,a) &= \sum_{k=-a}^{a} (-1)^k \binom{2a}{a+k}^3 \\
&= \sum_{k=-a}^{a} (-1)^k \left( \binom{2a-1}{a+k} + \binom{2a-1}{a-1+k} \right)^3 \\
&= \sum_{k=-a}^{a} (-1)^k \left( \binom{2a-1}{a+k}^3 + 3\binom{2a}{a+k}\binom{2a-1}{a+k}\binom{2a-1}{a-1+k} + \binom{2a-1}{a-1+k}^3 \right) \\
&= \sum_{k=-a}^{a} (-1)^k \binom{2a-1}{a+k}^3 + \sum_{k=-a}^{a} (-1)^k \binom{2a-1}{a-1+k}^3 + 3S(a,a,a-1) \\
&= \sum_{k=-a}^{a-1} (-1)^k \binom{2a-1}{a+k}^3 + \sum_{k=-a+1}^{a} (-1)^k \binom{2a-1}{a-1+k}^3 + 3S(a,a,a-1) \\
&= \sum_{k=-a}^{a-1} (-1)^k \binom{2a-1}{a+k}^3 + \sum_{t=-a}^{a-1} (-1)^{t+1} \binom{2a-1}{a+t}^3 + 3S(a,a,a-1) \\
&= 3S(a,a,a-1).
\end{aligned}$$

Therefore,

$$S(a,a,a) = 3S(a,a,a-1). \tag{11}$$

Now we give a proof of Eq. (1) using induction:

Eq. (1) is true if $a = b = c = 0$. Let $n \in \mathbb{N}_0$ be fixed. Assume that Eq. (1) holds for all nonnegative integers $a$, $b$ and $c$, such that $a + b + c = n$. Let $a$, $b$ and $c$ be nonnegative integers such that $a + b + c = n + 1$. Then we have three cases:

Case 1: If, at least, one of numbers $a$, $b$ or $c$ is equal to zero, then Eq. (1) obviously holds. Therefore, we may assume that $a, b, c \in \mathbb{N}$.

Case 2: If $a = b = c$, then we use Eq. (11). From the induction hypothesis, we have that

$$S(a,a,a-1) = \frac{(3a-1)!}{a! \cdot a! \cdot (a-1)!} .$$

$$\begin{aligned}
\text{Then } \; S(a,a,a) &= 3 \cdot \frac{(3a-1)!}{a! \cdot a! \cdot (a-1)!} \\
&= 3a \cdot \frac{(3a-1)!}{a! \cdot a! \cdot (a-1)! \cdot a} \\
&= \frac{(3a)!}{a! \cdot a! \cdot a!}, \text{ as desired.}
\end{aligned}$$

Case 3: If $a \neq b$, then we use Eq. (9). From the induction hypothesis, we have that

$$S(a-1,b,c) = \frac{(a+b+c-1)!}{(a-1)! \cdot b! \cdot c!}, \text{ and } S(a,b-1,c) = \frac{(a+b+c-1)!}{a! \cdot (b-1)! \cdot c!}.$$

$$\begin{aligned}
S(a,b,c) &= \frac{1}{a-b}\big((a+c)S(a-1,b,c) - (b+c)S(a,b-1,c)\big) \\
&= \frac{1}{a-b}\big((a+c)\frac{(a+b+c-1)!}{(a-1)! \cdot b! \cdot c!} - (b+c)\frac{(a+b+c-1)!}{a! \cdot (b-1)! \cdot c!}\big) \\
&= \frac{1}{a-b} \cdot \frac{(a+b+c-1)!}{(a-1)! \cdot (b-1)! \cdot c!}\big(\frac{a+c}{b} - \frac{b+c}{a}\big) \\
&= \frac{(a+b+c-1)! \cdot (a^2 - b^2 + c(a-b))}{(a-b) \cdot a! \cdot b! \cdot c!} \\
&= \frac{(a+b+c-1)! \cdot (a-b)(a+b+c)}{(a-b) \cdot a! \cdot b! \cdot c!} \\
&= \frac{(a+b+c)!}{a! \cdot b! \cdot c!}, \text{ as desired.}
\end{aligned}$$

If $a = b$ then $a \neq c$. We do similarly as before using Eq. (10). This proves Case 3. By induction, Eq. (1) follows. □

# 3 Acknowledgments

I want to thank to my professor Duško Jojić.

# References

[1] A. C. Dixon, Summation of a certain series, *Proc. London Math. Soc.* **35** (1) (1903) 285–289.

[2] S. B. Ekhad, A very short proof of Dixon's theorem, *J. Combin. Theory*, Ser. A **54** (1990) 141–142.

[3] I. Gessel and D. Stanton, Short proofs of Saalschütz's and Dixon's theorems, *J. Combin. Theory*, Ser. A **38** (1985) 87–90.

[4] I. J. Good, Proofs of some 'binomial' identities by means of MacMahon's 'Master Theorem' , *Proc. Cambridge Philos. Soc.* **58** (1962) 161–162.

[5] Victor J. W. Guo, A simple proof of Dixon's identity, *Discrete Math.* **268** (2003) 309–310.

---

2010 *Mathematics Subject Classification*: Primary 11B65; Secondary 05A10.
*Keywords:* recurrence equation, binomial coefficient, Dixon's identity, auxiliary sum.

---

Received March 18 2016; revised version received May 10 2016. Published in *Journal of Integer Sequences*, June 2 2016.

---

Return to Journal of Integer Sequences home page.