PUBLICATIONS DE L'INSTITUT MATHÉMATIQUE
Nouvelle série, tome 57 (71), 1995, 81–90
Đuro Kurepa memorial volume

EXPANSIONS OF THE KUREPA FUNCTION

G.V. Milovanović

Dedicated to the memory of Professor Đ. Kurepa

Abstract. The Taylor series expansions of the Kurepa function $K(a+z)$, $a \geq 0$, and numerical determination of their coefficients $b_\nu(a)$ for $a=0$ and $a=1$ are given. An asymptotic behaviour of $b_\nu(a)$ as well as that $|b_\nu(a)/b_{\nu+1}(a)| \sim a+1$, when $\nu \to \infty$, are shown. Using this fact, a transformation of series with much faster convergence is done. Numerical values of coefficients in such a transformed series for $a=0$ and $a=1$ are given with 30 decimal digits. Also, the Chebyshev expansions of $K(1+z)$ and $1/K(1+z)$ are obtained.

1. Introduction

In 1971 Professor Đ. Kurepa (see [**8**–**9**]) defined so-called left factorial as

$$!n = 0! + 1! + \cdots + (n-1)!$$

and extended it to the complex plane

$$K(z) = \int_0^\infty \frac{t^z - 1}{t-1}\, e^{-t}\, dt \qquad (\operatorname{Re} z > 0). \tag{1.1}$$

This function can be extended analytically to the hole complex plane by

$$K(z) = K(z+1) - \Gamma(z+1), \tag{1.2}$$

where $\Gamma(z)$ is the gamma function defined by

$$\Gamma(z) = \int_0^\infty e^{-t} t^{z-1}\, dt \quad (\operatorname{Re} z > 0) \quad \text{and} \quad z\Gamma(z) = \Gamma(z+1).$$

AMS Subject Classification (1991): Primary 33B15, 65D20
Research partly supported by Science Fund of Serbia under grant 0401F.

Kurepa [**9**] proved that $K(z)$ is a meromorphic function with simple poles at the points $z_k = -k$ ($k \in \mathbb{N} \setminus \{2\}$). Slavić [**17**] found the representation

$$K(z) = -\frac{\pi}{e}\cot \pi z + \frac{1}{e}\left(\sum_{n=1}^{\infty}\frac{1}{n!n} + \gamma\right) + \sum_{n=0}^{\infty}\Gamma(z-n),$$

where γ is Euler's constant. These formulas were mentioned also in the book [**11**]. A number of problems and hypotheses, especially in number theory, were posed by Kurepa and then considered by several mathematicians. For details and a complete list of references see a recent survey written by Ivić and Mijajlović [**6**].

Evaluation of $K(z)$ for some specific z in $(0,1)$, using quadrature formulas with relatively small accuracy, was done by Slavić and the author of this paper (see [**9**]).

In this paper we give power series expansions of the Kurepa function $K(a+z)$, $a \geq 0$, and determine numerical values of their coefficients $b_\nu(a)$ for $a = 0$ and $a = 1$. Using an asymptotic behaviour of $b_\nu(a)$, when $\nu \to \infty$, we give a transformation of series with much faster convergence. Also, we obtain the Chebyshev expansions for $K(1+z)$ and $1/K(1+z)$. For similar expansions of the gamma function see e.g. Davis [**3**], Luke [**10**], Fransén and Wrigge [**4**], and Bohman and Fröberg [**1**].

2. Power series expansions

There are many investigations about the gamma function $\Gamma(z)$, including numerical calculations. Some old references can be found in Luke [**10**]. In 1980 Fransén and Wrigge [**4**] determined numerical values of the coefficients in the Taylor series expansion

$$\Gamma(a+x)^m = \sum_{k=0}^{\infty} g_k(m,a)x^k$$

for certain values of m and a and used these values to calculate $\Gamma(p/q)$ ($p, q = 1(1)10$; $p < q$) with a high precision. Recently, Bohman and Fröberg [**1**] have given power series expansions of the form

$$\Gamma(n+1+z) = n!(1 + d_1 z + d_2 z^2 + \cdots) \qquad (n = 2, 3, 4, 10)$$

and

$$(-1)^n n!\Gamma(-n+z) = \frac{n}{1-z} - \frac{1}{(n+1)(1+z)} + \frac{1}{z}\left(1 + f_1 z + f_2 z^2 + \cdots\right)$$

for $n = 0$ 1, 2, 10.

In this section we investigate the Kurepa function in a similar way.

Let $a \geq 0$. Differentiating (1.1) we find

$$K^{(\nu)}(a) = \int_0^{\infty} \frac{t^a \log^\nu t}{t-1}\, e^{-t}\, dt \qquad (\nu \geq 1), \tag{2.1}$$

so that the Taylor series expansion is given by

$$K(a+z) = b_0(a) + b_1(a)z + b_2(a)z^2 + b_3(a)z^3 + \cdots, \tag{2.2}$$

where

$$b_0(a) = K(a), \qquad b_\nu(a) = \frac{1}{\nu!}\,K^{(\nu)}(a) \qquad (\nu \geq 1). \tag{2.3}$$

Taking a to be an integer, we have that $b_0(a) =!a$. For example, $b_0(0) = 0$, $b_0(1) = 1$, $b_0(2) = 2$, $b_0(3) = 4$, etc.

The following theorem gives an asymptotic behaviour of $b_\nu(a)$.

THEOREM 2.1. *For every $a > 0$, $b_\nu(a) \to 0$, when $\nu \to \infty$. If $a = 0$ we have*

$$\lim_{\nu\to\infty} b_{2\nu}(0) = -1, \qquad \lim_{\nu\to\infty} b_{2\nu+1}(0) = 1.$$

Precisely,

$$b_\nu(a) \sim \frac{(-1)^{\nu+1}}{(a+1)^{\nu+1}}, \qquad \textit{when} \quad \nu \to \infty.$$

Proof. We start with the identity

$$K(a) = R_1(a) + R_2(a), \tag{2.4}$$

where

$$R_1(a) = \int_0^1 \frac{t^a - 1}{t-1}\,e^{-t}\,dt \quad \text{and} \quad \int_1^\infty \frac{t^a-1}{t-1}\,e^{-t}\,dt.$$

Differentiating (2.4) ν times with respect to a, we obtain

$$b_\nu(a) = \frac{K^{(\nu)}(a)}{\nu!} = -\frac{1}{\nu!}\int_0^1 \frac{t^a \log^\nu t}{1-t}\sum_{k=0}^{\infty}\frac{(-1)^k t^k}{k!}\,dt + \frac{1}{\nu!}\,R_2^{(\nu)}(a), \tag{2.5}$$

i.e.,

$$b_\nu(a) = -\frac{1}{\nu!}\sum_{k=0}^{\infty}\frac{(-1)^k}{k!}\int_0^1 \frac{t^{k+a}\log^\nu t}{1-t}\,dt + \frac{1}{\nu!}\,R_2^{(\nu)}(a). \tag{2.6}$$

Using the Karamata inequality [**7**] (cf. [**14**, p. 272]),

$$\frac{\log t}{t-1} \leq \frac{1}{\sqrt{t}} \qquad (t \geq 1),$$

for the last term on the right in (2.6), we have

$$\begin{aligned} 0 \leq \frac{1}{\nu!}\,R_2^{(\nu)}(a) &= \frac{1}{\nu!}\int_1^\infty \frac{t^a \log^\nu t}{t-1}\,e^{-t}\,dt \\ &\leq \frac{1}{\nu!}\int_1^\infty \frac{(t-1)^{\nu-1}}{t^{\nu/2-a}}\,e^{-t}\,dt \\ &\leq \frac{1}{\nu! e}\int_0^\infty \frac{z^{\nu-1}}{(z+1)^{\nu/2-a}}\,e^{-z}\,dz, \end{aligned}$$

i.e.,

$$\frac{1}{\nu!}\,R_2^{(\nu)}(a) < \frac{1}{\nu! e}\int_0^\infty z^{\nu/2+a-1}e^{-z}\,dz = \frac{1}{\nu! e}\,\Gamma\Big(\frac{\nu}{2}+a\Big) \to 0,$$

when $\nu \to \infty$.

On the other side, we have that (cf. [**16**, p. 491])

$$\int_0^1 \frac{t^{k+a}\log^\nu t}{1-t}\,dt = (-1)^\nu \nu!\zeta(\nu+1, k+a+1),$$

where $\zeta(z,\alpha) = \sum\limits_{m=0}^{\infty}(\alpha+m)^{-z}$ is the generalized Riemman function (the Hurwitz Zeta function).

Therefore,

$$b_r(a) = (-1)^{\nu+1}\sum_{k=0}^{\infty}\frac{(-1)^k}{k!}\,\zeta(\nu+1,k+a+1) + \frac{1}{\nu!}\,R_2^{(\nu)}(a). \tag{2.7}$$

When $\nu \to \infty$, we see that

$$\zeta(\nu+1,q+1) = \frac{1}{(q+1)^{\nu+1}} + \sum_{m=1}^{\infty}\frac{1}{(q+1+m)^{\nu+1}} \to \begin{cases} 0, & \text{if } q>0,\\ 1, & \text{if } q=0. \end{cases}$$

Thus, when $\nu \to \infty$ all terms in the right-hand side of (2.7) approaches zero except the term $(-1)^{\nu+1}$ in the case when $a=0$ ($k=0,\ q=k+a=0$). The last statement of the theorem follows immediately from (2.7). □

From this theorem we obtain:

COROLLARY 2.2. *For the coefficients $b_\nu(0)$ in the series* (2.2) *we have that $b_\nu(0) = (-1)^{\nu+1}(1+\varepsilon_\nu)$, where $\lim\limits_{\nu\to\infty}\varepsilon_\nu = 0$.*

Remark 2.1. We note that the radius of convergence of the series (2.2) increases as a increases, because of $|b_\nu(a)/b_{\nu+1}(a)| \sim a+1$, when $\nu \to \infty$.

The following theorem gives an expansion with a faster convergence.

THEOREM 2.3. *For $|z|<1$ we have*

$$K(z) = \frac{1}{1+z}\sum_{\nu=1}^{\infty}\alpha_\nu z^\nu, \tag{2.8}$$

where $\alpha_\nu = b_\nu(0) + b_{\nu-1}(0) = (-1)^{\nu-1}\Delta\varepsilon_{\nu-1}$, $\nu \ge 1$. The coefficients $b_\nu(0)$ and ε_ν are given in Corollary 2.2.

Proof. Applying the Euler-Abel transformation (cf. Milovanović [**12**, pp. 49–50]) to (2.2), with $a=0$, we obtain (2.8). □

Remark 2.2. The series (2.8) converges also for $z=1$. Notice that $\sum\limits_{\nu=1}^{\infty}\alpha_\nu = 2$.

Similar to Theorem 2.3 we can state and prove the following more general result:

THEOREM 2.4. *For $|z| < a+1$ we have*

$$K(a+z) = \frac{1}{a+1+z} \sum_{\nu=0}^{\infty} \beta_\nu(a) z^\nu, \tag{2.9}$$

where

$$\beta_0(a) = (a+1)b_0(a), \quad \beta_\nu(a) = (a+1)b_{\nu+1}(a) + b_\nu(a), \quad \nu \geq 1. \tag{2.10}$$

Proof. Put $K(a+z) = b_0(a) + \big(z/(a+1)\big)g(z)$, where

$$g(z) = \sum_{\nu=1}^{\infty} (a+1)^\nu b_\nu(a) \Big(\frac{z}{a+1}\Big)^{\nu-1}.$$

Since

$$\Big(1 + \frac{z}{a+1}\Big) g(z) = \sum_{\nu=0}^{\infty} \big\{(a+1)b_{\nu+1}(a) + b_\nu(a)\big\} z^\nu - b_0(a),$$

we obtain (2.9), where the coefficients $\beta_\nu(a)$ are given by (2.10). This is, in fact, the Euler-Abel transformation of the series (2.2). □

Remark 2.3. Suppose that $a \geq 1$. For $z = 1$ and $z = -1$, (2.9) gives

$$\sum_{\nu=0}^{\infty} \beta_\nu(a) = (a+2)K(a+1) \quad \text{and} \quad \sum_{\nu=0}^{\infty} (-1)^\nu \beta_\nu(a) = aK(a-1),$$

respectively. If $a = 1$, we have that

$$\sum_{\nu=0}^{\infty} \beta_\nu(1) = 6 \quad \text{and} \quad \sum_{\nu=0}^{\infty} (-1)^\nu \beta_\nu(1) = 0.$$

Numerical calculation of $b_\nu(a)$ for $\nu \geq 1$ is not so easy, because of singular integrand in (2.1). Some Gaussian type of quadrature formulas, e.g. Gauss-Laguerre formulas or, after changing variable, Gauss-Einstein formulas on $(0, \infty)$ (cf. Gautschi and Milovanović [**5**]) show a slowly convergence. Using the fact that for an integral of an analytic function over $(-\infty, \infty)$ the trapezoidal rule with an equal mesh size is asymptotically optimal among formulas with the same density of sampling points (see Takahasi and Mori [**18**]), we will transform our integrals to ones over $(-\infty, \infty)$, taking $t = \exp(\pi \sinh x)$, and then apply the trapezoidal quadrature formula

$$I = \int_{-\infty}^{\infty} g(x)\, dx \approx I_h = h \sum_{n=-\infty}^{\infty} g(nh) \tag{2.11}$$

For a such kind of quadratures obtained by variable transformation, especially for the DE-rule, see a survey written by Mori [**15**].

TABLE 2.1: The coefficients in the series (2.8) and (2.9) when $a = 1$

ν	$\beta_\nu(0)$	$\beta_\nu(1)$
0	0.00000 00000 00000 00000 00000 00000	2.00000 00000 00000 00000 00000 00000
1	1.43220 57346 53224 41481 10310 06215	2.70998 01395 03383 10840 90378 32265
2	0.46888 45450 62632 23498 89857 40662	0.90645 96812 26452 30535 12196 88028
3	0.06386 42753 51364 60958 84436 58189	0.26514 75834 59521 37636 12078 98719
4	0.02916 62163 10236 51587 58769 27935	0.08712 40652 66430 32799 74653 52816
5	0.00384 29068 21880 78393 78013 20684	0.02344 30113 03591 62453 71921 03613
6	0.00183 20459 86444 14916 92261 41329	0.00610 50968 58694 42506 90720 59033
7	0.00009 73017 96964 77704 35972 03248	0.00137 03067 97499 58760 40919 95658
8	0.00010 88168 64034 75461 36273 53524	0.00029 98495 87774 13386 75427 15149
9	-0.00000 83124 80989 14474 80756 83316	0.00005 71279 77327 82762 62249 42045
10	0.00000 76093 36747 52906 90149 05470	0.00001 09812 77997 28649 23721 64909
11	-0.00000 16730 50119 56080 52965 01901	0.00000 17875 37686 00232 08394 11611
12	0.00000 06860 95605 33952 08525 32143	0.00000 03177 80645 24735 74359 66125
13	-0.00000 02068 17935 17460 02890 90873	0.00000 00427 02999 64944 38064 00163
14	0.00000 00715 53267 36586 97321 77786	0.00000 00078 08364 44191 12066 23204
15	-0.00000 00233 56503 79198 66091 37138	0.00000 00007 17143 23915 78700 00764
16	0.00000 00078 18902 04675 37681 86066	0.00000 00001 87654 45234 13299 33159
17	-0.00000 00025 94278 60137 72193 76381	0.00000 00000 01586 36642 25452 36344
18	0.00000 00008 64332 38788 78858 80196	0.00000 00000 05714 87184 82397 51814
19	-0.00000 00002 87697 66083 26943 17995	-0.00000 00000 00680 75693 40826 49802
20	0.00000 00000 95832 46989 44643 89904	0.00000 00000 00256 93394 02012 50700
21	-0.00000 00000 31923 54355 14763 58582	-0.00000 00000 00054 25128 23079 80583
22	0.00000 00000 10636 48853 89104 25473	0.00000 00000 00014 61950 44184 99559
23	-0.00000 00000 03544 28133 05353 85378	-0.00000 00000 00003 53643 92780 66646
24	0.00000 00000 01181 12896 81039 46306	0.00000 00000 00000 89493 51613 20739
25	-0.00000 00000 00393 63482 92231 22875	-0.00000 00000 00000 22239 88173 56536
26	0.00000 00000 00131 19298 21915 64240	0.00000 00000 00000 05567 76999 51411
27	-0.00000 00000 00043 72633 98654 72079	-0.00000 00000 00000 01390 20407 68723
28	0.00000 00000 00014 57428 48201 29110	0.00000 00000 00000 00347 51114 79725
29	-0.00000 00000 00004 85780 47285 87821	-0.00000 00000 00000 00086 84201 06978
30	0.00000 00000 00001 61919 57473 07582	0.00000 00000 00000 00021 70582 21355
31	-0.00000 00000 00000 53971 38024 88922	-0.00000 00000 00000 00005 42530 90710
32	0.00000 00000 00000 17990 00746 96841	0.00000 00000 00000 00001 35611 61567
33	-0.00000 00000 00000 05996 55604 63516	-0.00000 00000 00000 00000 33898 54027
34	0.00000 00000 00000 01998 82374 65548	0.00000 00000 00000 00000 08473 77501
35	-0.00000 00000 00000 00666 26751 66870	-0.00000 00000 00000 00000 02118 27096
36	0.00000 00000 00000 00222 08740 61997	0.00000 00000 00000 00000 00529 53328
37	-0.00000 00000 00000 00074 02869 39615	-0.00000 00000 00000 00000 00132 37643
38	0.00000 00000 00000 00024 67612 09747	0.00000 00000 00000 00000 00033 09273
39	-0.00000 00000 00000 00008 22534 60745	-0.00000 00000 00000 00000 00008 27291

40	0.00000 00000 00000 00002 74177 51295	0.00000 00000 00000 00000 00002 06817
41	-0.00000 00000 00000 00000 91392 33194	-0.00000 00000 00000 00000 00000 51703
42	0.00000 00000 00000 00000 30464 06756	0.00000 00000 00000 00000 00000 12926
43	-0.00000 00000 00000 00000 10154 67841	-0.00000 00000 00000 00000 00000 03231
44	0.00000 00000 00000 00000 03384 89011	0.00000 00000 00000 00000 00000 00808
45	-0.00000 00000 00000 00000 01128 29603	-0.00000 00000 00000 00000 00000 00202

Thus, coefficients given by (2.3) and (2.1) reduce to

$$b_\nu(a) = \frac{\pi^{\nu+1}}{2\nu!}\int_{-\infty}^{\infty} \frac{\sinh^\nu x\cosh x\exp\left((a+\frac{1}{2})\pi\sinh x - e^{\pi\sinh x}\right)}{\sinh\left(\frac{\pi}{2}\sinh x\right)}\,dx \qquad (\nu \geq 1).$$

Similarly,

$$\alpha_\nu = \frac{\pi^{\nu+1}}{2(\nu+1)!}\int_{-\infty}^{\infty} \frac{\sinh^\nu x\cosh x\exp\left(\frac{1}{2}\pi\sinh x - e^{\pi\sinh x}\right)}{\sinh\left(\frac{\pi}{2}\sinh x\right)}\left[\pi\sinh x + \nu + 1\right]\,dx$$

for $\nu \geq 1$.

Using (2.11) in Q-arthmetic on the MICROVAX 3400 computer (machine precision $\approx 1.93 \times 10^{-34}$), we computed the coefficients $b_\nu(a)$ and $\beta_\nu(a)$ $(\nu \leq 60)$ for $a = 0$ and $a = 1$. Here, we give in Table 2.1 the coefficients $\beta_\nu(0)$ and $\beta_\nu(1)$ only for $\nu = 0(1)45$ with 30D.

3. Chebyshev expansions

Let $T_k^*(z)$ be the shifted Chebyshev polynomial of the first kind of degree k given by $T_k^*(z) = T_k(2z-1)$, $T_k(x) = \cos k\theta$ $(x = \cos\theta)$. Numerical values of the coefficients in the Chebyshev expansions for the gamma function

$$\Gamma(z+1) = \sum_{k=0}^{\infty}\gamma_k T_k^*(z) \quad \text{and} \quad \frac{1}{\Gamma(z+1)} = \sum_{k=0}^{\infty}\delta_k T_k^*(z) \qquad (0 \leq z \leq 1),$$

were given by Clenshaw [**2**] to 20D (see also Luke [**10**, p. 13]).

In this section we obtain similar expansions for the Kurepa function. Namely, we determine the coefficients c_k and d_k in the Chebyshev expansions

$$K(z+1) = \sum_{k=0}^{\infty} c_k T_k^*(z) \quad \text{and} \quad \frac{1}{K(z+1)} = \sum_{k=0}^{\infty} d_k T_k^*(z) \qquad (0 \leq z \leq 1). \quad (3.1)$$

Taking $x = 2z - 1$, we can reduce our problem to the interval $(-1, 1)$ and then use the least square approximation, with respect to the Chebyshev weight $w(x) = (1-x^2)^{-1/2}$, in order to find the coefficients c_k and d_k. Thus,

$$c_k = \frac{\left(K((x+3)/2), T_k\right)}{\|T_k\|^2}, \qquad d_k = \frac{\left(1/K((x+3)/2), T_k\right)}{\|T_k\|^2} \qquad (k \geq 0),$$

where the inner product and the norm are given by

$$(f,g)=\int_{-1}^{1} f(x)\overline{g(x)}w(x)\,dx, \qquad \|f\|^2=(f,f),$$

respectively.

TABLE 3.1: The coefficients c_k and d_k in series (3.1)

k	c_k	d_k
0	1.47404 93837 23786 91003 52009 68753	0.71933 88033 78564 80199 26920 51918
1	0.49590 32661 94158 22700 15501 19905	-0.24486 53611 27056 06741 80872 17436
2	0.02579 30869 03070 45788 53165 66164	0.02976 94905 09647 98622 36085 35374
3	0.00407 08812 31791 61386 34001 49176	-0.00498 33867 56921 33739 51852 46148
4	0.00015 73577 18751 30746 82134 28808	0.00086 57497 58509 74532 38298 56153
5	0.00002 57037 05826 92036 26292 93518	-0.00014 67971 82657 38722 42886 97797
6	0.00000 01775 04779 26512 29793 59549	0.00002 51924 09034 18335 36671 90502
7	0.00000 01478 12089 72264 27761 10704	-0.00000 43238 54800 40900 43392 52848
8	-0.00000 00057 65834 77355 97860 46608	0.00000 07414 54647 70219 98837 74381
9	0.00000 00010 46615 98880 68659 86996	-0.00000 01272 20052 30389 19464 41242
10	-0.00000 00000 83632 76388 29507 65862	0.00000 00218 27577 03559 46944 44963
11	0.00000 00000 09422 94336 75632 53320	-0.00000 00037 44928 13250 51616 37824
12	-0.00000 00000 00910 90916 80081 86592	0.00000 00006 42530 56969 56371 69126
13	0.00000 00000 00093 62274 78249 23554	-0.00000 00001 10240 75381 57048 29851
14	-0.00000 00000 00009 39687 44816 46570	0.00000 00000 18914 30308 43243 17659
15	0.00000 00000 00000 95140 34153 77730	-0.00000 00000 03245 18210 36461 92610
16	-0.00000 00000 00000 09603 61612 70033	0.00000 00000 00556 78518 95627 89374
17	0.00000 00000 00000 00970 39299 20617	-0.00000 00000 00095 52923 18136 84543
18	-0.00000 00000 00000 00098 02159 05061	0.00000 00000 00016 39022 51658 48302
19	0.00000 00000 00000 00009 90238 99413	-0.00000 00000 00002 81211 80450 47223
20	-0.00000 00000 00000 00001 00033 64229	0.00000 00000 00000 48248 31777 95406
21	0.00000 00000 00000 00000 10105 45983	-0.00000 00000 00000 08278 10261 20992
22	-0.00000 00000 00000 00000 01020 85778	0.00000 00000 00000 01420 29786 64079
23	0.00000 00000 00000 00000 00103 12756	-0.00000 00000 00000 00243 68458 86448
24	-0.00000 00000 00000 00000 00010 41800	0.00000 00000 00000 00041 80966 55298
25	0.00000 00000 00000 00000 00001 05243	-0.00000 00000 00000 00007 17340 45282
26	-0.00000 00000 00000 00000 00000 10632	0.00000 00000 00000 00001 23076 16403
27	0.00000 00000 00000 00000 00000 01074	-0.00000 00000 00000 00000 21116 53134
28	-0.00000 00000 00000 00000 00000 00108	0.00000 00000 00000 00000 03623 02400
29	0.00000 00000 00000 00000 00000 00011	-0.00000 00000 00000 00000 00621 61264
30	-0.00000 00000 00000 00000 00000 00001	0.00000 00000 00000 00000 00106 65187

Since $\|T_0\|^2=\pi$ and $\|T_k\|^2=\pi/2$ $(k\ge 1)$, we have

$$c_0=\frac{1}{\pi}\int_{-1}^{1}\frac{1}{\sqrt{1-x^2}}K\Big(\frac{x+3}{2}\Big)\,dx, \quad c_k=\frac{2}{\pi}\int_{-1}^{1}\frac{1}{\sqrt{1-x^2}}K\Big(\frac{x+3}{2}\Big)T_k(x)\,dx,$$

where $k \geq 1$. Similarly,

$$d_0 = \frac{1}{\pi}\int_{-1}^{1} \frac{1}{\sqrt{1-x^2}} \frac{1}{K\Big(\dfrac{x+3}{2}\Big)}\,dx, \quad d_k = \frac{2}{\pi}\int_{-1}^{1} \frac{1}{\sqrt{1-x^2}} \frac{T_k(x)}{K\Big(\dfrac{x+3}{2}\Big)}\,dx \quad (k \geq 1).$$

For numerical determination of these coefficients, we apply the N-point Gauss-Chebyshev quadrature formula (see [**13**])

$$\int_{-1}^{1} \frac{1}{\sqrt{1-x^2}} f(x)\,dx = \frac{\pi}{N}\sum_{k=1}^{N} f\big(x_k^{(N)}\big) + R_N(f),$$

where $x_k^{(N)}$, $k = 1, \ldots, N$, are the zeros of N-th Chebyshev polynomial $T_N(x)$, i.e., $x_k^{(N)} = \cos((2k-1)\frac{\pi}{2n}$. This formula is exact for all algebraic polynomials of degree at most $2N-1$, i.e., $R_N(\mathcal{P}_{2N-1}) \equiv 0$. To compute values of the Kurepa function we use the power series expansion from Section 2. In order to obtain numerical values to 30D of coefficients c_k and d_k for $k \leq 30$, the Gaussian formula (3.1) needs $N = 40$ and $N = 35$ points, respectively. Numerical values are displayed in Table 3.1.

REFERENCES

[1] J. Bohman and C.-E. Fröberg, *The Γ-function revisited: power series expansions and real-imaginary zero lines*, Math. Comp. **58** (1992), 315–322.

[2] C.W. Clenshaw, *Chebyshev Series for Mathematical Functions*, Nat. Phys. Lab. Math. Tables, vol. V, H.M. Stationery Office, London, 1962.

[3] H.T. Davis, *Tables of the Higher Mathematical Functions*, vol. I, Principia Press, Bloomington, IN, 1933.

[4] A. Fransén and S. Wrigge, *High-precision values of the gamma function and some related coefficients*, Math. Comp. **34** (1980), 553–566.

[5] W. Gautschi and G.V. Milovanović, *Gaussian quadrature involving Einstein and Fermi functions with an application to summation of series*, Math. Comp. **44** (1985), 177–190.

[6] A. Ivić and Ž. Mijajlović, *On Kurepa problems in number theory*, These Publications.

[7] J. Karamata, *Sur quelques problèmes posés par Ramanujan*, J. Indian Math. Soc. (N.S.) **24** (1960), 343–365.

[8] Đ. Kurepa, *On the left factorial function* $!n$, Math. Balkanica **1** (1971), 147–153.

[9] Đ. Kurepa, *Left factorial function in complex domain*, Math. Balkanica **3** (1973), 297–307.

[10] Y.L. Luke, *Mathematical Functions and Their Approximations*, Mir, Moscow, 1980 (Russian).

[11] O.I. Marichev, *Handbook of Integral Transformation of Higher Transcendental Functions: Theory and Algorithmic Tables*, Ellis Horwood Ltd., Chichester, 1983

[12] G.V. Milovanović, *Numerical Analysis*, vol. I, Naučna Knjiga, Beograd, 1991 (Serbian).

[13] G.V. Milovanović, *Numerical Analysis*, vol. II, Naučna Knjiga, Beograd, 1991 (Serbian).

[14] D.S. Mitrinović and P.M. Vasić, *Analytic Inequalities*, Springer-Verlag, Berlin – Heidelberg – New York, 1970.

[15] M. Mori, *Quadrature formulas obtained by variable transformation and DE-rule*, J. Comput. Appl. Math. **12&13** (1985), 119–130.

[16] A.P. Prudnikov, Yu.A. Brychkov, O.I. Marichev, *Intergrals and Series: Elementary Functions*, Nauka, Moscow, 1981 (Russian).

[17] D.V. Slavić, *On the left factorial function of the complex argument*, Math. Balkanica **3** (1973), 472–477.

[18] H. Takashasi and M. Mori, *Error estimation in the numerical integration of analytic functions*, Report Computer Centre University of Tokyo **3** (1970), 41–108.

Elektronski fakultet
Katedra za matematiku
18001 Niš, p.p. 73
Yugoslavia

(Received 20 09 1995)