PUBLICATIONS DE L'INSTITUT MATHÉMATIQUE
Nouvelle série, tome 84(98) (2008), 49–60 DOI: 10.2298/PIM0898049M

ORTHOGONAL POLYNOMIALS FOR THE OSCILLATORY-GEGENBAUER WEIGHT

Gradimir V. Milovanović, Aleksandar S. Cvetković, and Zvezdan M. Marjanović

Abstract. This is a continuation of our previous investigations on polynomials orthogonal with respect to the linear functional $\mathcal{L} : \mathcal{P} \to \mathbb{C}$, where $\mathcal{L} = \int_{-1}^{1} p(x)\, d\mu(x)$, $d\mu(x) = (1-x^2)^{\lambda-1/2} \exp(i\zeta x)\, dx$, and $\mathcal{P}$ is a linear space of all algebraic polynomials. Here, we prove an extension of our previous existence theorem for rational $\lambda \in (-1/2, 0]$, give some hypothesis on three-term recurrence coefficients, and derive some differential relations for our orthogonal polynomials, including the second order differential equation.

1. Introduction

In this paper we continue our investigation on orthogonality with respect to the oscillatory weight functions studied in [**7**], [**8**], and [**9**]. We are concerned with the following measure $d\mu(x) = (1-x^2)^{\lambda-1/2} \exp(i\zeta x)\, dx$, supported on the interval $[-1, 1]$, where $\lambda > -1/2$ and where $\zeta \in \mathbb{R}\backslash\{0\}$. Evidently, the corresponding weight function

$$w(x) = w^{\zeta,\lambda}(x) = (1-x^2)^{\lambda-1/2} \exp(i\zeta x)$$

depends on two real parameters λ and ζ. We investigate the questions connected with the existence of polynomials π_n orthogonal with respect to the linear functional

$$\mathcal{L}[p] = \int_{-1}^{1} p(x)\, d\mu(x), \quad p \in \mathcal{P}, \tag{1.1}$$

where $\mathcal{P}$ is the space of all algebraic polynomials. As it has been proved in [**9**] it is enough to consider only the case $\zeta > 0$, and therefore we continue our discussion only for such values of ζ. As we proved in [**9**] it can be easily inferred that the orthogonal polynomials π_n exist in the case $\lambda \in \mathbb{Q}$ with $\lambda > 0$ and ζ is positive zero of the Bessel function of the first kind $J_{\lambda-1}$. The crucial ingredient in the proof is the result which states that non-trivial zeros of Bessel functions of the first kind are

2000 *Mathematics Subject Classification*: Primary 30C10; Secondary 33C47.

The authors were supported in part by the Serbian Ministry of Science (Project: Orthogonal Systems and Applications, grant number #144004) and the Swiss National Science Foundation (SCOPES Joint Research Project No. IB7320–111079 "New Methods for Quadrature").

transcendental provided their index is rational. In this paper we extend the results from [**9**] for $-1/2 < \lambda \leqslant 0$, $\lambda \in \mathbb{Q}$, and ζ being the zero of the Bessel function $J_{\lambda-1}$.

Also, in this paper, we investigate the possibility of computation of three-term recurrence coefficients for the given sequence of orthogonal polynomials $\{\pi_n\}_{n\in\mathbb{N}_0}$, as well as some consequences of the weight function w being semiclassical. We also find the second order differential equation for the corresponding orthogonal polynomials.

2. Existence of orthogonal polynomials

According to (1.1), the moments

$$\mu_k = \mathcal{L}[x^k] = \int_{-1}^{1} x^k (1-x^2)^{\lambda-1/2} \exp(i\zeta x)\, dx, \quad k \in \mathbb{N}_0, \tag{2.1}$$

can be expressed in terms of Bessel functions J_ν of the order ν defined by

$$J_\nu(z) = \sum_{m=0}^{+\infty} \frac{(-1)^m (z/2)^{\nu+2m}}{m!\Gamma(\nu+m+1)}.$$

Using the initial conditions

$$\mu_0 = AJ_\lambda(\zeta), \quad \mu_1 = iA\left(\frac{2\lambda}{\zeta}J_\lambda(\zeta) - J_{\lambda-1}(\zeta)\right),$$
$$\mu_2 = A\left[\left(1 - \frac{2\lambda(2\lambda+1)}{\zeta^2}\right)J_\lambda(\zeta) + \frac{2\lambda+1}{\zeta}J_{\lambda-1}(\zeta)\right],$$

where $A = (2/\zeta)^\lambda \sqrt{\pi}\,\Gamma(\lambda+1/2)$, we can see that the moments (2.1) satisfy the following recurrence relation

$$\mu_{k+2} = -\frac{k+2\lambda+1}{i\zeta}\,\mu_{k+1} + \mu_k + \frac{k}{i\zeta}\,\mu_{k-1}.$$

The following result is proved in [**9**].

THEOREM 2.1. *The moments μ_k can be expressed in the form*

$$\mu_k = \frac{A}{(i\zeta)^k}\left(P_k^\lambda(\zeta)J_\lambda(\zeta) + Q_k^\lambda(\zeta)J_{\lambda-1}(\zeta)\right), \quad k \in \mathbb{N}_0, \tag{2.2}$$

where $A = (2/\zeta)^\lambda\sqrt{\pi}\,\Gamma(\lambda+1/2)$, and P_k^λ and Q_k^λ are polynomials in ζ, which satisfy the following four-term recurrence relation

$$y_{k+2} = -(k+2\lambda+1)y_{k+1} - \zeta^2 y_k - k\zeta^2 y_{k-1},$$

with the initial conditions

$$P_0^\lambda(\zeta) = 1,\ P_1^\lambda(\zeta) = -2\lambda,\ P_2^\lambda(\zeta) = 2\lambda(2\lambda+1) - \zeta^2$$

and

$$Q_0^\lambda(\zeta) = 0,\ Q_1^\lambda(\zeta) = \zeta,\ Q_2^\lambda(\zeta) = -(2\lambda+1)\zeta,$$

respectively.

Some obvious properties of the polynomials P_n^λ and Q_n^λ, $n \in \mathbb{N}_0$, are stated in the following lemma.

LEMMA 2.1. *For the degree of the polynomials P_n^λ and Q_n^λ we have*

$$\deg(P_{2n}^\lambda, P_{2n+1}^\lambda) = 2n \quad (n \in \mathbb{N}_0) \qquad \text{and} \qquad \deg(Q_{2n}^\lambda, Q_{2n-1}^\lambda) = 2n \quad (n \in \mathbb{N}).$$

The free terms in the polynomials P_n^λ and Q_n^λ are $(-1)^n(2\lambda)_n$ and zero, respectively. The free term in the polynomial $Q_n(\zeta)/\zeta$, $n \in \mathbb{N}$, equals $(-1)^{n-1}(2\lambda+1)_{n-1}$.

The leading coefficients in the polynomials P_{2n}^λ, P_{2n-1}^λ, Q_{2n+1}^λ and Q_{2n}^λ are $(-1)^n$, $(-1)^n(2n\lambda+n-1)$, $(-1)^n$ and $(-1)^n n(2\lambda+1)$, respectively.

Now we are ready to state and prove a stronger version of the existence theorem than the one given in [**9**].

THEOREM 2.2. *Let $\lambda \in \mathbb{Q}$, $\lambda > -1/2$ and let $\zeta \neq 0$ be a zero of the Bessel function $J_{\lambda-1}$. Then the sequence of monic polynomials $\{\pi_n\}_{n\in\mathbb{N}_0}$ orthogonal with respect to the linear functional* (1.1) *exists.*

PROOF. As in [**9**] our result is based upon completely algebraic facts. Consider the sequence of Hankel determinants defined by

$$\Delta_n = \begin{vmatrix} \mu_0 & \mu_1 & \cdots & \mu_{n-1} \\ \mu_1 & \mu_2 & \cdots & \mu_n \\ \vdots & \vdots & \ddots & \vdots \\ \mu_{n-1} & \mu_n & \cdots & \mu_{2n-2} \end{vmatrix}, \quad n \in \mathbb{N}.$$

The sequence of orthogonal polynomials $\{\pi_n\}_{n\in\mathbb{N}_0}$ with respect to $\mathcal{L}$ exists provided $\Delta_n \neq 0$, $n \in \mathbb{N}$ (see [**4**]). Thus, let ζ be a nontrivial zero of the Bessel function $J_{\lambda-1}$. Then we can express the moments of the linear functional $\mathcal{L}$ in the following form

$$\mu_k = \frac{A}{(i\zeta)^k} P_k^\lambda(\zeta) J_\lambda(\zeta), \quad k \in \mathbb{N}_0.$$

Obviously, from the Hankel determinant we can factor $(AJ_\lambda(\zeta))^{n+1}/(i\zeta)^{n(n+1)}$ and we are left with the sequence of Hankel determinants H_n, $n \in \mathbb{N}$, made from the new moment sequence $\mu_k' = P_k^\lambda(\zeta)$, $k \in \mathbb{N}_0$,

$$H_n = \begin{vmatrix} P_0^\lambda(\zeta) & P_1^\lambda(\zeta) & \dots & P_{n-1}^\lambda(\zeta) \\ P_1^\lambda(\zeta) & P_2^\lambda(\zeta) & \dots & P_n^\lambda(\zeta) \\ \vdots & \vdots & & \vdots \\ P_{n-1}^\lambda(\zeta) & P_n^\lambda(\zeta) & \dots & P_{2n-2}^\lambda(\zeta) \end{vmatrix}.$$

Accordingly, all H_n, $n \in \mathbb{N}$, are polynomials in ζ which have rational coefficients since $\lambda \in \mathbb{Q}$. If some $H_n = 0$, then ζ is the zero of the polynomial H_n with rational coefficients and ζ is zero of the Bessel function $J_{\lambda-1}$. According to [**13**, p. 220], based on the results by [**11**], [**12**] (see also [**6**]), non-trivial zeros of the Bessel functions J_λ, $\lambda \in \mathbb{Q}$, $-\lambda \notin \mathbb{N}$, are transcendental and cannot be the zeros of the polynomials with rational coefficients unless polynomials are identically equal to zero.

We have to prove that determinants H_n are not identically equal to zero. To prove this fact we emphasize that, according to Lemma 2.1, the free coefficient in the polynomial P_n^λ equals $(-1)^n(2\lambda)_n$, $n \in \mathbb{N}_0$, so that the free coefficient in the

polynomial H_n equals to the value of the Hankel determinant $\widetilde{H}_n$ made for the sequence of moments $\mu_k'' = (2\lambda)_k$, $k \in \mathbb{N}_0$. For $\lambda > 0$, we can easily recognize the Hankel determinants $\widetilde{H}_n$ as being the same, up to multiplicative factor $(\Gamma(2\lambda))^n$, as the Hankel determinants for the generalized Laguerre weight $x^{2\lambda-1}e^{-x}$ on $[0, +\infty)$. Hence, none of the determinants $\widetilde{H}_n$ cannot be equal according to the existence of the sequence of orthogonal polynomials with respect to the Laguerre weight function. This establishes the proof for $\lambda > 0$ (cf. [**9**]).

Actually, we can calculate the determinant $\widetilde{H}_n$ for $\lambda > 0$ in the form

$$\widetilde{H}_n = \prod_{l=0}^{n-1}(n-l)! \prod_{i=0}^{l}(2\lambda + l - i), \quad n \in \mathbb{N}.$$

We easily inspect that the same result about the values of the Hankel determinants $\widetilde{H}_n$ holds even in the case of $-1/2 < \lambda \leqslant 0$. This due to the fact that Hankel determinants $\widetilde{H}_n$ as being polynomials are analytic functions of λ as well as expressions representing their values. Inspecting values of the determinants $\widetilde{H}_n$, $n \in \mathbb{N}$, we easily conclude that $\widetilde{H}_n \neq 0$, $n \in \mathbb{N}$, for $\lambda \neq 0$. This in turn implies that $\Delta_n \neq 0$, $n \in \mathbb{N}$. Case $\lambda = 0$ requires a special attention.

Consider now the case $\lambda = 0$. As we easily inspect we have that $\widetilde{H}_n = 0$, $n \in \mathbb{N}$, for $\lambda = 0$. Consider determinants H_n, we have $H_n = (\Gamma(2\lambda))^n \widetilde{H}_n$, $n \in \mathbb{N}$. Since H_n, $n \in \mathbb{N}$, are analytic functions of λ we can compute values of H_n, $n \in \mathbb{N}$, for $\lambda = 0$, as the limiting values of the expressions on the right hand side. We have

$$\lim_{\lambda \to 0} (\Gamma(2\lambda))^n \prod_{l=0}^{n-1}(n-l)! \prod_{i=0}^{l}(2\lambda + l - i) = n! \prod_{l=1}^{n-1}(n-l)! \prod_{i=0}^{l-1}(l-i) \neq 0.$$

Thus, we have proved that if $\lambda \in \mathbb{Q}$, $\lambda > -1/2$, and ζ is the zero of the Bessel function $J_{\lambda-1}$, then the sequence of Hankel determinants

$$\Delta_n = (AJ_\lambda(\zeta))^n / (i\zeta)^{n(n-1)} H_n, \quad n \in \mathbb{N},$$

contains no zero entry. Accordingly, the sequence of orthogonal polynomials with respect to the linear functional $\mathcal{L}$ exists. □

Regarding Theorem 2.2, we suppose such parameters λ and ζ which provide the existence of orthogonal polynomials π_n with respect to the linear functional (1.1). The quasi inner-product $(p, q) := \mathcal{L}[pq]$, in our case has the property $(zp, q) = (p, zq)$, and because of that, the corresponding (monic) polynomials $\{\pi_n\}_{n\in\mathbb{N}_0}$ satisfy the fundamental three-term recurrence relation

$$\pi_{n+1}(x) = (x - i\alpha_n)\pi_n(x) - \beta_n \pi_{n-1}(x), \quad n \in \mathbb{N}, \tag{2.3}$$

with $\pi_0(x) = 1$, $\pi_{-1}(x) = 0$. The recursion coefficients α_n and β_n can be expressed in terms of Hankel determinants as (cf. [**9**]).

$$i\alpha_n = \frac{\Delta_{n+1}'}{\Delta_{n+1}} - \frac{\Delta_n'}{\Delta_n} = \frac{1}{i\zeta}\left(\frac{H_{n+1}'}{H_{n+1}} - \frac{H_n'}{H_n}\right) \tag{2.4}$$

$$\beta_n = \frac{\Delta_{n+1}\Delta_{n-1}}{\Delta_n^2} = \frac{1}{(i\zeta)^2}\frac{H_{n+1}H_{n-1}}{H_n^2}, \tag{2.5}$$

where H_n is defined before and Δ'_n is the Hankel determinant Δ_{n+1} with the penultimate column and the last row removed. The corresponding determinant H'_n is given by

$$H'_n = \begin{vmatrix} P_0^\lambda(\zeta) & P_1^\lambda(\zeta) & \dots & P_{n-2}^\lambda(\zeta) & P_n^\lambda(\zeta) \\ P_1^\lambda(\zeta) & P_2^\lambda(\zeta) & \dots & P_{n-1}^\lambda(\zeta) & P_{n+1}^\lambda(\zeta) \\ \vdots & \vdots & & \vdots & \vdots \\ P_{n-1}^\lambda(\zeta) & P_n^\lambda(\zeta) & \dots & P_{2n-3}^\lambda(\zeta) & P_{2n-1}^\lambda(\zeta) \end{vmatrix}.$$

Although β_0 can be arbitrary, as usual it is convenient to take $\beta_0 = \mu_0 = AJ_\lambda(\zeta)$.

In this case, however, the values of Hankel determinants cannot be found easily, but, it is clear that the recursion coefficients are rational functions in ζ. Using our software package [**5**] we can generate coefficients even in symbolic form for some reasonable small values of n (for $n \leqslant 2$ see Table 1).

TABLE 1. Recursion coefficients α_n and β_n for $n \leqslant 2$

n	α_n	β_n
0	$\frac{2\lambda}{\zeta}$	$\left(\frac{2}{\zeta}\right)^\lambda \sqrt{\pi}\,\Gamma\left(\lambda+\frac{1}{2}\right) J_\lambda(\zeta)$
1	$\frac{4\lambda(1+\lambda)-\zeta^2}{\zeta(2\lambda-\zeta^2)}$	$\frac{\zeta^2-2\lambda}{\zeta^2}$
2	$\frac{(7+6\lambda)\zeta^6-4\lambda(4+\lambda)(3+2\lambda)\zeta^4+32\lambda^2(3+2\lambda)\zeta^2-32\lambda^3(2+\lambda)}{2\zeta(\zeta^2-2\lambda)[\zeta^4-\lambda(5+2\lambda)\zeta^2+4\lambda^2]}$	$-\frac{2(1+2\lambda)[\zeta^4-\lambda(5+2\lambda)\zeta^2+4\lambda^2]}{\zeta^2(-2\lambda+\zeta^2)^2}$

According to symbolic calculations we can conjecture that H_n is a polynomial in ζ^2 of degree m,

$$H_n = S_m(\zeta^2) = \sum_{\nu=0}^{m} \gamma_\nu(\lambda)\zeta^{2\nu},$$

where $m = n^2/4$ for even n and $m = (n^2-1)/4$ for odd n, so that, because of (2.5),

$$\beta_{2k} = -\frac{S_{k(k+1)}(\zeta^2)S_{k(k-1)}(\zeta^2)}{\zeta^2 S_{k^2}(\zeta^2)^2}, \quad \beta_{2k+1} = -\frac{S_{(k+1)^2}(\zeta^2)S_{k^2}(\zeta^2)}{\zeta^2 S_{k(k+1)}(\zeta^2)^2}.$$

In a similar way, we can conjecture that the numerator in (2.4) is a polynomial in ζ^2 of degree $n(n+1)/2$,

$$H'_n H_{n+1} - H'_{n+1} H_n = V_{n(n+1)/2}(\zeta^2),$$

so that

$$\alpha_{2k} = \frac{V_{k(2k+1)}(\zeta^2)}{\zeta S_{k^2}(\zeta^2)S_{k(k+1)}(\zeta^2)^2}, \quad \alpha_{2k+1} = \frac{V_{(k+1)(2k+1)}(\zeta^2)}{\zeta S_{k(k+1)}(\zeta^2)S_{(k+1)^2}(\zeta^2)}.$$

Increasing n, the complexity of expressions for α_n and β_n increases quite rapidly. On the other side, using the Chebyshev algorithm, similarly as in [**7**], an efficient numerical construction of recursion coefficients can be done.

According to $\overline{\mu}_k = (-1)^k \mu_k$, $k \in \mathbb{N}_0$ (see (2.2)) we can prove:

LEMMA 2.2. *If the sequence of monic orthogonal polynomials* $\{\pi_n\}_{n\in\mathbb{N}_0}$ *exists, then* $\pi_n(z) = (-1)^n\overline{\pi_n(-\overline{z})}$.

3. Differential-difference relations

In this section we consider some differential relations for (monic) polynomials $\{\pi_n\}_{n\in\mathbb{N}_0}$ orthogonal with respect to the linear functional (1.1), i.e., to the the (quasi) inner-product product

$$(p,q) := \mathcal{L}[pq] = \int_{-1}^{1} p(x)\,q(x)\,w(x)\,dx,$$

supposing such parameters λ and ζ, which provide the existence of our orthogonal polynomials.

Our weight function $w(x) = (1-x^2)^{\lambda-1/2}e^{i\zeta x}$ satisfies the following Pearson type differential equation

$$(\phi w)' = \psi w, \quad \phi = 1-x^2, \quad \psi = -(2\lambda+1)x + i\zeta(1-x^2).$$

As before, we put $d\mu(x) = w(x)\,dx$.

THEOREM 3.1. *For every* $n \in \mathbb{N}$, *we have*

$$\phi\pi_n' + \psi\pi_n = \sum_{k=n-1}^{n+2} \theta_n^k \pi_k, \tag{3.1}$$

where

$$\begin{aligned}
\theta_n^{n+2} &= -i\zeta,\\
\theta_n^{n+1} &= \zeta(\alpha_{n+1}+\alpha_n) - (n+2\lambda+1),\\
2\theta_n^{n} &= -(2\lambda+1)i\alpha_n + i\zeta(1-\beta_{n+1}-\beta_n+\alpha_n^2),\\
\theta_n^{n-1} &= (n-1)\beta_n,
\end{aligned}$$

and α_n *and* β_n *are coefficients in the recurrence relation* (2.3).

PROOF. Since

$$\int_{-1}^{1} (\phi w\pi_n)' x^k dx = \phi w\pi_n x^k \Big|_{-1}^{1} - k\int_{-1}^{1} \phi\pi_n x^{k-1} d\mu = -k\int_{-1}^{1} \phi\pi_n x^{k-1} d\mu = 0$$

for $k < n-1$, in that case we conclude that $(\phi w\pi_n)'/w$ is orthogonal to x^k. Therefore, we have directly

$$\frac{(\phi w\pi_n)'}{w} = \phi\pi_n' + \psi\pi_n = \sum_{k=n-1}^{n+2} \theta_n^k \pi_k, \tag{3.2}$$

according to the fact that $\deg(\psi) = 2$. Since we are concerned with the monic polynomials it is readily seen that $\theta_n^{n+2} = -i\zeta$.

For θ_n^{n-1}, after multiplying (3.2) with π_{n-1} and integrating with respect to $d\mu$, we get

$$\theta_n^{n-1}\|\pi_{n-1}\|^2 = \int_{-1}^{1}(\phi w\pi_n)'\pi_{n-1}dx = -\int_{-1}^{1}\phi w\pi_n\pi'_{n-1}dx = (n-1)\|\pi_n\|^2,$$

and accordingly $\theta_n^{n-1} = (n-1)\beta_n$.

We also have

$$\begin{aligned}\theta_n^k\|\pi_k\|^2 + \theta_k^n\|\pi_n\|^2 &= \int_{-1}^{1}\left[(\phi w\pi_n)'\pi_k + (\phi w\pi_k)'\pi_n\right]dx \\ &= \int_{1}^{1}[-\phi\pi_n\pi'_k + \phi\pi'_k\pi_n + \psi\pi_k\pi_n]d\mu = \int_{1}^{1}\psi\pi_n\pi_k\,d\mu.\end{aligned}$$

The last equality for $k = n+1$ gives

$$\begin{aligned}\theta_n^{n+1} &= -\frac{\theta_{n+1}^n}{\beta_{n+1}} + \frac{1}{\|\pi_{n+1}\|^2}\int_{-1}^{1}\left[-(2\lambda+1)x + i\zeta(1-x^2)\right]\pi_n\pi_{n+1}\,d\mu \\ &= -(n+2\lambda+1) - \frac{i\zeta}{\|\pi_{n+1}\|^2}\int_{-1}^{1}x^2\pi_n\pi_{n+1}\,d\mu \\ &= -(n+2\lambda+1) + (\alpha_{n+1}+\alpha_n)\zeta,\end{aligned}$$

because of $x\pi_n = \pi_{n+1} + i\alpha_n\pi_n + \beta_n\pi_{n-1}$ and

$$\begin{aligned}\int_{-1}^{1}x^2\pi_n\pi_{n+1}\,d\mu &= \int_{-1}^{1}(\pi_{n+1} + i\alpha_n\pi_n + \beta_n\pi_{n-1})(\pi_{n+2} + i\alpha_{n+1}\pi_{n+1} + \beta_{n+1}\pi_n)d\mu \\ &= i\alpha_{n+1}\|\pi_{n+1}\|^2 + i\alpha_n\beta_n\|\pi_n\|^2.\end{aligned}$$

Similarly, for $k = n$, we get

$$\begin{aligned}2\theta_n^n &= \frac{1}{\|\pi_n\|^2}\int_{-1}^{1}\psi\pi_n^2 d\mu \\ &= \frac{1}{\|\pi_n\|^2}\left\{-(2\lambda+1)\int_{-1}^{1}x\pi_n^2 d\mu + i\zeta\left(\|\pi_n\|^2 - \int_{-1}^{1}x^2\pi_n^2 d\mu\right)\right\} \\ &= i[-(2\lambda+1)\alpha_n + \zeta(1-\beta_{n+1}-\beta_n+\alpha_n^2)],\end{aligned}$$

because of

$$\begin{aligned}\int_{-1}^{1}x^2\pi_n^2 d\mu &= \int_{-1}^{1}(\pi_{n+1} + i\alpha_n\pi_n + \beta_n\pi_{n-1})^2 d\mu \\ &= \|\pi_{n+1}\|^2 - \alpha_n^2\|\pi_n\|^2 + \beta_n^2\|\pi_{n-1}\|^2.\end{aligned}$$

Thus, the equality (3.1) is proved. □

LEMMA 3.1. *The monic polynomials orthogonal with respect to the linear functional* (1.1) *satisfy the following differential-difference equation*

$$\phi\pi'_n + \psi\pi_n = p_2^n\pi_n + q_1^n\pi_{n-1}, \quad n \in \mathbb{N},$$

where p_2^n and q_1^n are polynomials of the second and first degree, respectively.

PROOF. Starting with equation (3.1) and using the three-term recurrence relation (2.3) we get

$$\begin{aligned}\phi\pi_n' + \psi\pi_n = \left[(\theta_n^{n+2}(x - i\alpha_{n+1}) + \theta_n^{n+1})(x - i\alpha_n) - \beta_{n+1}\theta_n^{n+2} + \theta_n^n\right]\pi_n \\ + \left[-\beta_n(\theta_n^{n+2}(x - i\alpha_{n+1}) + \theta_n^{n+1}) + \theta_n^{n-1}\right]\pi_{n-1}.\end{aligned}$$

What is left to do is to read the respective polynomials. □

THEOREM 3.2. *The monic polynomials orthogonal with respect to the linear functional* (1.1) *satisfy the following equation*

$$\phi\pi_n' = p_1^n\pi_n + q_1^n\pi_{n-1}, \tag{3.3}$$

where

$$\begin{aligned}p_1^n &= -nx - \frac{i\zeta}{2}\left(1 - \beta_{n+1} + \beta_n + \alpha_n^2\right) + \frac{i\alpha_n}{2}(2n + 2\lambda + 1),\\ q_1^n &= \beta_n\left[2(n + \lambda) + i\zeta(x + i\alpha_n)\right].\end{aligned}$$

PROOF. Using expressions given in the proof of Lemma 3.1 and Theorem 3.1 we get exactly what is stated. □

THEOREM 3.3. *The polynomials p_1^n and q_1^n, which appear in* (3.3)*, satisfy the following recurrence relations*

$$\begin{aligned}p_1^{n+1} &= -q_1^n\frac{x - i\alpha_n}{\beta_n} + p_1^{n-1} + q_1^{n-1}\frac{x - i\alpha_{n-1}}{\beta_{n-1}},\\ q_1^{n+1} &= (x - i\alpha_n)\left[p_1^n + q_1^n\frac{x - i\alpha_n}{\beta_n} - p_1^{n-1} - q_1^{n-1}\frac{x - i\alpha_{n-1}}{\beta_{n-1}}\right] + \phi + q_1^{n-1}\frac{\beta_n}{\beta_{n-1}},\end{aligned}$$

respectively.

PROOF. We can prove this result using almost the same arguments from the proof in [**7**]. Namely, we have

$$\begin{aligned}\phi\pi_{n+1}' - \phi\pi_n &= (x - i\alpha_n)\phi\pi_n' - \beta_n\phi\pi_{n-1}'\\ &= (x - i\alpha_n)(p_1^n\pi_n + q_1^n\pi_{n-1}) - \beta_n(p_1^{n-1}\pi_{n-1} + q_1^{n-1}\pi_{n-2})\\ &= -\left(q_1^n\frac{x - i\alpha_n}{\beta_n} - p_1^{n-1} - q_1^{n-1}\frac{x - i\alpha_{n-1}}{\beta_{n-1}}\right)\pi_{n+1}\\ &\quad + \left\{(x - i\alpha_n)p_1^n + q_1^{n-1}\frac{\beta_n}{\beta_{n-1}}\right.\\ &\quad \left. + (x - i\alpha_n)\left(q_1^n\frac{x - i\alpha_n}{\beta_n} - p_1^{n-1} - q_1^{n-1}\frac{x - i\alpha_{n-1}}{\beta_{n-1}}\right)\right\}\pi_n,\end{aligned}$$

from where we just need to read corresponding polynomials. □

In the sequel we derive a system of nonlinear recurrence relations for the recursion coefficients α_n and β_n, which needs a few first coefficients as starting values (see Table 1).

THEOREM 3.4. *We have*

$$p_1^{n+1} + p_1^n + q_1^n \frac{x - i\alpha_n}{\beta_n} = (2\lambda - 1)x - i\zeta(1 - x^2), \quad n \in \mathbb{N}. \tag{3.4}$$

PROOF. Starting with the previous theorem we can prove that quantity

$$p_1^{n+1} + p_1^n + q_1^n \frac{x - i\alpha_n}{\beta_n},$$

is independent of n. Hence, using Table 1, we can prove that

$$p_1^2 + p_1^1 + q_1^1 \frac{x - i\alpha_1}{\beta_1} = (2\lambda - 1)x - i\zeta(1 - x^2),$$

and hence finish the proof. □

THEOREM 3.5. *The three-term recurrence coefficients satisfy the following system of nonlinear recurrence equations*

$$\begin{aligned}
\beta_{n+2} &= \beta_n + \alpha_{n+1}^2 - \alpha_n^2 - \frac{1}{\zeta}\big[\alpha_{n+1}(2n + 2\lambda + 3) - \alpha_n(2n + 2\lambda - 1)\big], \\
\alpha_{n+2} &= \frac{1}{2\beta_{n+2}}\left[2\alpha_n\beta_{n+1} + \alpha_{n+1}(2\beta_{n+1} - 2\beta_n - \beta_{n+2} + \alpha_{n+1}^2 - \alpha_n^2)\right] \\
&\quad + \frac{1}{\zeta\beta_{n+2}}\big[\alpha_{n+1}^2\big(n + \lambda + 1/2\big) - \alpha_{n+1}\alpha_n\big(n + \lambda - 1/2\big) \\
&\qquad + 2\beta_{n+2}(n + \lambda + 2) - 2\beta_{n+1}(n + \lambda) - 1\big]
\end{aligned}$$

PROOF. The first equation we prove quite easily. Namely, simply compare coefficients with x^0 in (3.4) and we have the stated equation.

For the proof of the second equation we have to use the second relation given in Theorem 3.3. In the same way, comparing coefficients with the term x^0 in this relation we obtain the result. □

Now, we can prove a result concerning zeros of orthogonal polynomials π_n.

THEOREM 3.6. *The polynomials π_{2n+1}, $n \in \mathbb{N}_0$, have only simple zeros. The even polynomials π_{2n}, $n \in \mathbb{N}$, have only simple zeros except possibly one zero of the multiplicity two which is purely imaginary.*

PROOF. It is a trivial fact that polynomials π_n and π_{n-1} cannot have zeros in common. If x is multiple zero of π_n, using equation (3.3), it must be $q_1^n(x) = 0$. However, q_1^n is a polynomial of first degree and, therefore, the multiplicity of the zero can be at most two. Using a symmetry property of polynomials, we see that if x is a zero of the polynomial π_n, then also $-\overline{x}$ is the zero of this polynomial π_n. Hence, we conclude that π_n cannot have a multiple zero outside of the imaginary axis. Finally, according to the symmetry we conclude that if polynomial of odd degree has zero on the imaginary axis it must be of odd multiplicity. Hence, all polynomials of odd degree have only simple zeros. □

We can now also state a result connected with a differential equation which solutions are polynomials orthogonal with respect to the functional (1.1).

THEOREM 3.7. *The polynomials π_n orthogonal with respect to the functional* (1.1) *satisfy the following second order differential equation*

$$\phi q_1^n \pi_n'' + (\psi q_1^n - i\zeta\beta_n\phi)\pi_n' - \left[-nq_1^n - i\zeta\beta_n p_1^n - \frac{q_1^n(\hat{p}_1^n q_1^n - p_1^n \hat{q}_1^n)}{\phi}\right]\pi_n = 0,$$

where we used the notation from Theorem (3.2) *and*

$$\hat{p}_1^n = -q_1^{n-1}/\beta_{n-1} \quad \textit{and} \quad \hat{q}_1^n = p_2^{n-1} + (x - i\alpha_{n-1})q_1^{n-1}/\beta_{n-1}.$$

The term $\hat{p}_1^n q_1^n - p_1^n \hat{q}_1^n$ must have the factor ϕ.

PROOF. Starting with the equation (3.3) and the three-term recurrence relation, we compute easily

$$\phi\pi_{n-1}' = -\frac{q_1^{n-1}}{\beta_{n-1}}\pi_n + \left(p_1^{n-1} + q_1^{n-1}\frac{x - i\alpha_{n-1}}{\beta_{n-1}}\right)\pi_{n-1} = \hat{p}_1^n\pi_n + \hat{q}_1^n\pi_{n-1}.$$

Combining this equation with the one for the polynomial π_n, i.e. with equation 3.3, we get

$$\phi\left[\frac{\phi\pi_n' - p_1^n\pi_n}{q_1^n}\right]' = \hat{p}_1^n\pi_n + \hat{q}_1^n\frac{\phi\pi_n' - p_1^n\pi_n}{q_1^n}.$$

After some calculations we obtain the following differential equation

$$\begin{aligned}\phi q_1^n\pi_n'' &+ \left[q_1^n(\phi' - p_1^n - \hat{q}_1^n) - \phi(q_1^n)'\right]\pi_n' \\ &+ \left[-(p_1^n)'q_1^n + p_1^n(q_1^n)' - q_1^n\frac{\hat{p}_1^n q_1^n - p_1^n\hat{q}_1^n}{\phi}\right]\pi_n = 0.\end{aligned}$$

If we use Theorems 3.4 and 3.2, for the term with π_n' we got exactly what is stated. Similarly we have the same for the term with π_n.

Finally, it is clear that, since all other quantities are polynomials it must be that $q_1^n(\hat{p}_1^n q_1^n - p_1^n\hat{q}_1^n)/\phi$ is polynomial, too. Obviously it cannot be that q_1^n has zeros in common with ϕ since it can have only purely imaginary zero. □

This theorem can be used to give the following result:

THEOREM 3.8. *Let n be an odd integer and let x_ν^n, $\nu = 1, \ldots, n$, be simple zeros of the polynomial π_n. Then*

$$\frac{\pi_n''(x_\nu^n)}{\pi_n'(x_\nu^n)} + \frac{\lambda + 1/2}{x_\nu^n - 1} + \frac{\lambda + 1/2}{x_\nu^n + 1} + i\zeta - \frac{1}{x_\nu^n + i\alpha_n + (2n + 2\lambda)/i\zeta} = 0, \tag{3.5}$$

for each $\nu = 1, \ldots, n$.

PROOF. We start with the differential equation for polynomials π_n, derived in the previous theorem. We divide it with $\pi_n'(x_\nu^n) \neq 0$, $\nu = 1, \ldots, n$. This equality is fulfilled due to the fact that for odd n, according to Theorem 3.6, the polynomial π_n has only simple zeros. Thus, by division and using the fact that $\pi_n(x_\nu^n) = 0$, $\nu = 1, \ldots, n$, we have

$$\left(\frac{\pi_n''}{\pi_n'} + \frac{\psi}{\phi} - \frac{i\zeta\beta_n}{q_1^n}\right)(x_\nu^n) = 0, \quad \nu = 1, \ldots, n,$$

which after small calculation becomes exactly what is stated. □

This theorem enables the computation of zeros of orthogonal polynomial π_n of the high degree using technique similar as given in [**10**].

4. Zero distribution

Here we want to present some consequences of Theorem 3.8 in the light of some new results of Aptekarev and Van Assche (see [**1**]). Namely, provided that the sequence of polynomials π_n orthogonal with respect to the functional (1.1) exists, they have proved that for $\lambda = 0$ the three term recurrence coefficients satisfy the following asymptotic properties $\alpha_n \to 0$ and $\beta_n \to 1/4$. Actually, we can use this fact to give some interesting observations connected to the zero distribution of polynomials π_n. A direct consequence of the mentioned theorem is the boundness of the zeros x_ν^n, $\nu = 1, \ldots, n$, of polynomials π_n uniformly in $n \in \mathbb{N}$ (see [**2**], [**3**]).



FIGURE 1. Zero distribution of π_n, $n = 10(10)100$, for $\lambda = 11$ and $\zeta \approx 14.475500686554541238451637655413151976304 81$ (a zero of J_{10})



FIGURE 2. Zero distribution of π_n, $n = 10(10)100$, for $\lambda = 11$ and $\zeta \approx 100.71634047409274068436610723206019123535653$ (a zero of J_{10})

According to the mentioned result and for n sufficiently large, our equation (3.5) becomes

$$\frac{\pi_n''}{\pi_n'}(x_\nu^n) + \left(\lambda + \frac{1}{2}\right)\left(\frac{1}{x_\nu^n - 1} + \frac{1}{x_\nu^n + 1}\right) + i\zeta + O(n^{-1}) \approx 0,$$

for each $\nu = 1, \dots, n$. If we compare it to the similar equation for the classical Gegenbauer weight $w^{0,\lambda}$ (see [**14**]), we notice that there is only the difference in the term $i\zeta + O(n^{-1})$. Taking the imaginary part we get

$$2\sum_{\nu\neq k}\frac{\operatorname{Im}(x_k - x_\nu)}{|x_k - x_\nu|^2} + \left(\lambda + \frac{1}{2}\right)\left(\frac{\operatorname{Im}(x_k)}{|x_\nu^n - 1|^2} + \frac{\operatorname{Im}(x_k)}{|x_k^n + 1|^2}\right) - \zeta + O(n^{-1}) \approx 0,$$

which describes a property of zeros. Namely, for a fixed n and bigger ζ, zeros are at the bigger distance from the x axis as presented in Figures 1 and 2.

References

[1] A. I. Aptekarev, W. V. Assche, *Scalar and matrix Riemann-Hilbert approach to the strong asymptotics of Pade approximants and complex orthogonal polynomials with varying weight*, J. Appox. Theory **129** (2004), 129–166.

[2] B. Beckermann, *Complex Jacobi matrices*, J. Comput. Appl. Math. **127** (2001), 17–65.

[3] B. Beckerman, M. Castro, *On the determinacy of the comoplex Jacobi operators*, Publication ANNO 446, Universite Lille, 2002.

[4] T. S. Chihara, *An Introduction to Orthogonal Polynomials*, Gordon and Breach, New York, 1978.

[5] A. S. Cvetković, G. V. Milovanović, *The Mathematica package "OrthogonalPolynomials"*, Facta Univ. Ser. Math. Inform. **19** (2004), 17–36.

[6] N. I. Fel'dman, A. B. Šidlovskiĭ, *The development and present state of the theory of transcendental numbers*, Uspehi Mat. Nauk **22**:3 (1967), (135) 3–81 (Russian) [Engl. transl. in Russian Math. Surveys **22**:3 (1967), 1–79].

[7] G. V. Milovanović, A. S. Cvetković, *Orthogonal polynomials and Gaussian quadrature rules related to oscillatory weight functions*, J. Comput. Appl. Math. **179** (2005), 263–287.

[8] G. V. Milovanović, A. S. Cvetković, *Orthogonal polynomials related to the oscillatory-Chebyshev weight function*, Bull. Cl. Sci. Math. Nat. Sci. Math. **30** (2005), 47–60.

[9] G. V. Milovanović, A. S. Cvetković, M. P. Stanić, *Orthogonal polynomial and quadrature rules realted to Oscillatory-Gegenabuer weight function* (submitted; preprint version available at: `http://gauss-megatrend.edu.rs/radovi/GVM158.pdf`).

[10] L. Pasquini, *Accurate computation of the zeros of the generalized Bessel polynomials*, Numer. Math. **86** (2000), 507–538.

[11] C. L. Siegel, *Über einige Anwendungen diophantischer Approximationen*, Abh. Preuss. Akad. Wiss., Phys.-Math. **1** (1929), 1–70 [*Gesammelte Abhandlungen*. Springer-Verlag, Berlin–New York 1966 Band I, 209–266].

[12] C. L. Siegel, *Transcendental Numbers*, Annals of Mathematics Studies 16, Princeton University Press, Princeton, N. J., 1949.

[13] A. B. Šidlovskiĭ, *Transcendental Numbers*, Nauka, Moscow, 1987 (in Russian).

[14] G. Szegő, *Orthogonal Polynomials*, Amer. Math. Soc. Colloq. Publ. 23, 4th ed., Amer. Math. Soc., Providence, R. I., 1975.

Faculty of Computer Sciences, Megatrend University
Beograd, Serbia
`gvm@megatrend-edu.net`

Faculty of Science and Mathematics, University of Niš
Niš, Serbia
`aca@elfak.ni.ac.yu`

Faculty of Electronic Engineering, University of Niš
Niš, Serbia
`zvezdan@elfak.ni.ac.yu`