PUBLICATIONS DE L'INSTITUT MATHÉMATIQUE
Nouvelle série, tome 85(99) (2009), 35–38 DOI:10.2298/PIM0999035F

# ON THREE CONJECTURES INVOLVING THE SIGNLESS LAPLACIAN SPECTRAL RADIUS OF GRAPHS

**Lihua Feng and Guihai Yu**

*Communicated by Slobodan K. Simić*

Abstract. We study the signless Laplacian spectral radius of graphs and prove three conjectures of Cvetković, Rowlinson, and Simić [*Eigenvalue bounds for the signless Laplacian*, Publ. Inst. Math., Nouv. Sér. 81(95) (2007), 11–27].

## 1. Introduction

In this paper, we consider only simple connected graphs and follow the notation of [**1**]. Let $G$ be a simple graph with vertex set $V(G)$ and edge set $E(G)$. The *adjacency matrix of* $G$ is $A(G) = (a_{ij})$, where $a_{ij} = 1$ if two vertices $i$ and $j$ are adjacent in $G$ and $a_{ij} = 0$ otherwise. The characteristic polynomial of $G$ is just $P_G(x) = \det(xI - A(G))$. Let $D(G)$ be the diagonal degree matrix of $G$. We call the matrix $L(G) = D(G) - A(G)$ the *Laplacian matrix* of $G$, and the matrix $Q(G) = D(G) + A(G)$ the *signless Laplacian matrix* or $Q$*-matrix* of $G$. We denote the largest eigenvalues of $A(G)$, $L(G)$, $Q(G)$ by $\rho(G)$, $\lambda(G)$, $\mu(G)$, respectively, and call them the adjacency spectral radius, the Laplacian spectral radius, the signless Laplacian spectral radius (or the $Q$-spectral radius) of $G$, respectively.

The study of the signless Laplacian spectral radius has recently attracted researchers' attention. In [**10**], Fan et al. studied the signless Laplacian spectral radius of bicyclic graphs with fixed order. In [**9**], the authors discussed the smallest eigenvalue of $Q(G)$ as a parameter reflecting the nonbipartiteness of the graph $G$. In [**7**], the authors studied the smallest signless Laplacian eigenvalue of non-bipartite graphs. In [**11**], the extremal graphs with maximal signless Laplacian spectral radius and fixed diameter were studied. More information about the signless Laplacian can be found in [**2**], [**3**], [**5**], [**6**]. For more information about the spectral radius of graphs, the reader can refer to [**4**].

2000 *Mathematics Subject Classification*: Primary 05C50.

Partially supported by Foundation of Education Department of Shandong Province (J07YH03), NSFSD (No.Y2006A17, Y2008A04).

In [**5**], the authors proposed the following conjectures and in this paper we confirm that they are true.

THEOREM 1.1. [**5**, Conjecture 6] *Let $G$ be a connected graph of order $n \geqslant 4$. Then*

$$\mu(G) - \frac{4m}{n} \leqslant n - 4 + \frac{4}{n}.$$

*Equality holds if and only if $G = K_{1,n-1}$.*

THEOREM 1.2. [**5**, Conjecture 7] *Let $G$ be a connected graph of order $n \geqslant 5$. Then*

$$\mu(G) - \frac{2m}{n} \leqslant n - 1.$$

*Equality holds if and only if $G = K_n$.*

THEOREM 1.3. [**5**, Conjecture 10] *Let $G$ be a connected graph of order $n \geqslant 4$. Then*

$$\mu(G) - \lambda(G) \leqslant n - 2.$$

*Equality holds if and only if $G = K_n$.*

## 2. Lemmas and results

Let $G$ be a connected graph. The degree of $u$ in $G$ is denoted by $d_u$, the average degree of $u$, denoted by $m_u$, satisfies $d_u m_u = \sum_{uv \in E} d_v$, where $E = E(G)$.

LEMMA 2.1. [**8**] *Let $G$ be a graph with $n$ vertices, and $m$ edges. Then*

$$max\{d_v + m_v \mid v \in V(G)\} \leqslant \frac{2m}{n-1} + n - 2,$$

*with equality if and only if $K_{1,n-1} \subseteq G$ or $G = K_{n-1} \cup K_1$.*

LEMMA 2.2. [**12**] *Let $M = (m_{ij})$ be an $n \times n$ irreducible nonnegative matrix with spectral radius $\rho(M)$, and let $R_i(M)$ be the $i$th row sum of $M$, i.e., $R_i(M) = \sum_{j=i}^{n} m_{ij}$. Then*

$$\min\{R_i(M) \mid \leqslant i \leqslant n\} \leqslant \rho(M) \leqslant \max\{R_i(M) \mid 1 \leqslant i \leqslant n\}.$$

*Moreover, if the row sums of $M$ are not all equal, then both above inequalities are strict.*

LEMMA 2.3. *Let $G$ be a connected graph. Then $\mu(G) \leqslant max\{d_v + m_v \mid v \in V(G)\}$, with equality holding if and only if $G$ is either semiregular bipartite or regular.*

PROOF. We consider the matrix $K = D^{-1}QD$, where the row sum corresponding to the vertex $u$ is $d_u + m_u$. From Lemma 2.2, we obtain the required upper bound for $\mu(G)$.

If equality holds, then by Lemma 2.2, for a neighbor $v$ of $u$, $d_u + m_u = d_v + m_v$, thus $\sum_{uv \in E}(d_u + m_u) = \sum_{uv \in E}(d_v + m_v)$, that is $d_u^2 + d_u m_u = d_u m_u + \sum_{uv \in E} m_v$. So we get

$$d_u^2 = \sum_{uv \in E} m_v.$$

Suppose $d_u$ is the maximum degree. Then $d_v m_v = \sum_{wv\in E} d_w \leqslant d_u d_v$, whence $m_v \leqslant d_u$ for all $v \in V(G)$. Since $d_u^2 = \sum_{uv\in E} m_v \leqslant d_u^2$, we have for any edge $uv$, $d_u = m_v$, and $d_u d_v = d_v m_v$, that is, $\sum_{wv\in E}(d_u - d_w) = 0$. Since $d_u$ is the maximum degree, we have $d_u = d_w$ whenever there exists a vertex $v$ such that $uv, vw \in E$.

If $G$ does not contain odd cycles, then $G$ is bipartite. Suppose $V = S \cup T$ is a bipartion and $u \in T$; then $v \in S$, $w \in T$. This implies that the vertices in $T$ have the same degree. Similarly, the vertices in $S$ also have the same degree. So $G$ is semiregular bipartite.

If $G$ contains odd cycles, then $G$ must be regular.

The converse is easy to check. □

LEMMA 2.4. *Let $G$ be a connected graph with $n$ vertices and $m$ edges. Then*

$$\mu(G) \leqslant \frac{2m}{n-1} + n - 2,$$

*with equality if and only if $G$ is $K_{1,n-1}$ or $K_n$.*

PROOF. By Lemmas 2.1 and 2.3, we can get the result. Note that $K_{1,n-1}$ is the only semiregular bipartite graph and $K_n$ is the only regular graph that arises in the case of equality. □

Now we can present the proof of the main results of this paper.

PROOF OF THEOREM 1.1. By Lemma 2.4, we have

$$\begin{aligned}\mu(G) - \frac{4m}{n} &\leqslant \frac{2m}{n-1} + n - 2 - \frac{4m}{n} = (n-2)\Big(1 - \frac{2m}{n(n-1)}\Big)\\ &\leqslant (n-2)\Big(1 - \frac{2(n-1)}{n(n-1)}\Big) = n - 4 + \frac{4}{n}.\end{aligned}$$

The last inequality holds since $G$ is connected and so has at least $n-1$ edges. The equality case is easy to see from Lemma 2.4. □

PROOF OF THEOREM 1.2. By Lemma 2.4, we have

$$\mu(G) - \frac{2m}{n} \leqslant \frac{2m}{n-1} + n - 2 - \frac{2m}{n} = \frac{2m}{n(n-1)} + n - 2 \leqslant 1 + n - 2 = n - 1.$$

The last inequality holds since $G$ has at most $\frac{1}{2}n(n-1)$ edges. When this bound is attained, $G = K_n$. □

PROOF OF THEOREM 1.3. Note that the sum of all the Laplacian eigenvalues is $2m$, so we have $(n-1)\lambda(G) \geqslant 2m$ and hence $\lambda(G) \geqslant \frac{2m}{n-1}$. By Lemma 2.4, we have

$$\mu(G) - \lambda(G) \leqslant \frac{2m}{n-1} + n - 2 - \frac{2m}{n-1} = n - 2.$$

The equality case is easy to see from Lemma 2.4. □

**Acknowledgments.** The authors are grateful to the referees for their comments.

## References


[1] J. A. Bondy, U. S. R. Murty, *Graph Theory with Applications*, Macmillan Press, New York, 1976.

[2] D. Cvetković, *Signless Laplacians and line graphs*, Bull. Acad. Serbe Sci. Arts. Cl. Sci. Math. Nat. Sci. Math. **131**(30) (2005), 85–92.

[3] D. Cvetković, P. Rowlinson, S. K. Simić, *Signless Laplacians of finite graphs*, Linear Algebra Appl. **423** (2007), 155–171.

[4] D. Cvetković, M. Doob, H. Sachs, *Spectra of Graphs*, Academic Press, New York, 1980.

[5] D. Cvetković, P. Rowlinson, S. K. Simić, *Eigenvalue bounds for the signless Laplacian*, Publ. Inst. Math. (Beograd) **81** (95) (2007), 11–27.

[6] D. Cvetković, *New theorems for signless Laplacians eigenvalues*, Bull. Acad. Serbe Sci. Arts, Cl. Sci. Math. Natur., Sci. Math. **137**(33) (2008), 131–146.

[7] D. M. Cardoso, D. Cvetković, P. Rowlinson, S. K. Simić, *A sharp lower bound for the least eigenvalue of the signless Laplacian of a non-bipartite graph*, Linear Algebra Appl. **429** (2008), 2770–2780.

[8] K. Ch. Das, *Maximizing the sum of the squares of the degrees of a graph*, Discrete Math. **285** (2004), 57–66.

[9] M. Desai, V. Rao, *A characterizaion of the smallest eigenvalue of a graph*, J. Graph Theory **18** (1994), 181–194.

[10] Y. Fan, B. S. Tam, J. Zhou, *Maximizing spectral radius of unoriented Laplacian matrix over bicyclic graphs of a given order*, Linear Multilinear Algebra **56** (2008), 381–397.

[11] L. Feng, G. Yu, *The signless Laplacian spectral radius of graphs with given diameter*, 2008, Utilitas Math., accepted.

[12] R. A. Horn, C. R. Johnson, *Matrix Analysis*, Cambridge University Press, New York, 1985.



School of Mathematics
Shandong Institute of Business and Technology
191 Binhaizhong Road, Yantai, Shandong, 264005
P.R. China
fenglh@163.com



(Received 26 12 2008)
(Revised 10 03 2009)