PUBLICATIONS DE L'INSTITUT MATHÉMATIQUE
Nouvelle série, tome 101(115) (2017), 121–133 https://doi.org/10.2298/PIM1715121T

ON THE NUMBER OF SUBGROUPS OF A GIVEN EXPONENT IN A FINITE ABELIAN GROUP

Marius Tărnăuceanu and László Tóth

Abstract. This paper deals with the number of subgroups of a given exponent in a finite abelian group. Explicit formulas are obtained in the case of rank two and rank three abelian groups. An asymptotic formula is also presented.

1. Introduction

One of the most important problems of combinatorial abelian group theory is to determine the number of subgroups of a finite abelian group. This topic has enjoyed a constant evolution starting with the first half of the 20th century. Since a finite abelian group is a direct product of abelian p-groups, the above counting problem can be reduced to p-groups. Formulas which give the number of subgroups of type μ of a finite p-group of type λ were established by Delsarte [**4**], Djubjuk [**5**] and Yeh [**15**]. An excellent survey on this subject together with connections to symmetric functions was written by Butler [**2**] in 1994.

Another way to find the total number of subgroups of finite abelian p-groups was described by Bhowmik [**1**] by using divisor functions of matrices. By invoking different arguments, formulas in the case of rank two p-groups were obtained by Călugăreanu [**3**], Tărnăuceanu [**11, 12**], Hampejs, Holighaus, Tóth, Wiesmeyr [**6**], Tóth [**14**] and for rank three p-groups by Hampejs, Tóth [**7**], Oh [**8**]. Note that the papers [**6, 7, 14**] include also direct formulas for the groups $\mathbb{Z}_m \times \mathbb{Z}_n$ and $\mathbb{Z}_m \times \mathbb{Z}_n \times \mathbb{Z}_r$, respectively, where $m, n, r \in \mathbb{N}^* := \{1, 2, \dots\}$ are arbitrary.

The purpose of the current paper is to count the number of subgroups of a given exponent in a finite abelian p-group. Explicit formulas are obtained for rank two and rank three p-groups. The numbers of subgroups of exponent p, respectively p^2 in an arbitrary p-group are also considered. We prove that if two finite abelian groups have the same number of subgroups of any exponent, then they are isomorphic. We also deduce compact formulas for the number of subgroups of a given exponent of the group $\mathbb{Z}_m \times \mathbb{Z}_n$, where $m, n \in \mathbb{N}^*$ are arbitrary. Furthermore,

2010 *Mathematics Subject Classification:* Primary 20K01; Secondary 20K27, 11N37.
Key words and phrases: finite Abelian group, subgroup, exponent.
Communicated by Zoran Petrović.

we obtain an exact formula for the sum of exponents of the subgroups of $\mathbb{Z}_m \times \mathbb{Z}_n$, and an asymptotic formula for the arithmetic means of exponents of the subgroups of $\mathbb{Z}_n \times \mathbb{Z}_n$.

For the proofs we use two different approaches. The first one is based on the known formula for the number of subgroups of a given type in an abelian p-group, given in terms of gaussian coefficients (Theorem 2.1). The second method, applicable only for rank two groups, uses the representation of the subgroups of $\mathbb{Z}_m \times \mathbb{Z}_n$ obtained by the second author in [**14**] (Theorem 3.1).

Most of our notation is standard and usually will not be repeated here. For basic notions and results on group theory we refer the reader to [**10**].

2. First approach

Let G be a finite abelian group of order n and $G = G_1 \times G_2 \times \cdots \times G_m$ be the primary decomposition of G, where G_i is a p_i-group ($i = 1, 2, \dots, m$). For every divisor $d = p_1^{\alpha_1} p_2^{\alpha_2} \dots p_m^{\alpha_m}$ of n we denote $\mathcal{E}_d(G) = \{H \leqslant G : \exp(H) = d\}$.

Since the subgroups H of G are of type $H = H_1 \times H_2 \times \cdots \times H_m$ with $H_i \leqslant G_i$ ($i = 1, 2, \dots, m$), we infer that

$$|\mathcal{E}_d(G)| = \prod_{i=1}^{m} |\mathcal{E}_{p_i^{\alpha_i}}(G_i)|. \tag{2.1}$$

Equality (2.1) shows that the problem of counting the number of subgroups of exponent d in G is reduced to p-groups. So, in this section we will assume that G is a finite abelian p-group, that is a group of type $\mathbb{Z}_{p^{\lambda_1}} \times \mathbb{Z}_{p^{\lambda_2}} \times \cdots \times \mathbb{Z}_{p^{\lambda_k}}$ with $\lambda_1 \geqslant \lambda_2 \geqslant \cdots \geqslant \lambda_k \geqslant 1$. In this case we will say that G is of type λ, where λ is the partition $(\lambda_1, \lambda_2, \dots, \lambda_k, 0, \dots)$, and we will denote it by G_λ.

We recall the following well–known result, which gives the number of subgroups of type μ of G_λ (see [**4, 5, 15**]).

THEOREM 2.1. *For every partition $\mu \preceq \lambda$ (i.e., $\mu_i \leqslant \lambda_i$ for every $i \in \mathbb{N}^*$) the number of subgroups of type μ in G_λ is*

$$\alpha_\lambda(\mu; p) = \prod_{i \geqslant 1} p^{(a_i - b_i) b_{i+1}} \binom{a_i - b_{i+1}}{b_i - b_{i+1}}_p,$$

where $\lambda' = (a_1, a_2, \dots, a_{\lambda_1}, 0, \dots)$, $\mu' = (b_1, b_2, \dots, b_{\mu_1}, 0, \dots)$ are the partitions conjugate to λ and μ, respectively, and

$$\binom{n}{k}_p = \frac{\prod_{i=1}^{n}(p^i - 1)}{\prod_{i=1}^{k}(p^i - 1) \prod_{i=1}^{n-k}(p^i - 1)}$$

is the gaussian binomial coefficient (it is understood that $\prod_{i=1}^{m}(p^i - 1) = 1$ for $m = 0$).

By using Theorem 2.1 a way to compute the number of subgroups of exponent p^i in G_λ can be inferred, namely

$$|\mathcal{E}_{p^i}(G_\lambda)| = \sum_{\mu \preceq \lambda,\, \mu_1 = i} \alpha_\lambda(\mu; p) \quad (i = 0, 1, \dots, \lambda_1),$$

which is a polynomial in p with integer coefficients. If $i \geqslant 1$ and $k \geqslant 2$ are arbitrary, then the polynomial $|\mathcal{E}_{p^i}(G_\lambda)|$ can not be given explicitly, but we will do this in some particular cases. Namely, we will consider the following cases: $k \in \{2,3\}$ and $i \geqslant 1$ arbitrary, $i \in \{1,2\}$ and $k \geqslant 2$ arbitrary.

We remark first that since $\sum_{H\in\mathcal{E}_{p^i}(G_\lambda)} H \leqslant G_\lambda$ and $\sum_{H\in\mathcal{E}_{p^i}(G_\lambda)} H$ is of type $\mathbb{Z}_{p^i} \times \cdots \times \mathbb{Z}_{p^i} \times \mathbb{Z}_{p^{\lambda_r}} \times \cdots \times \mathbb{Z}_{p^{\lambda_k}}$, where $r = \min\{j : \lambda_j < i\}$, we have

$$|\mathcal{E}_{p^i}(G_\lambda)| = |\mathcal{E}_{p^i}(\mathbb{Z}_{p^i} \times \cdots \times \mathbb{Z}_{p^i} \times \mathbb{Z}_{p^{\lambda_r}} \times \cdots \times \mathbb{Z}_{p^{\lambda_k}})|$$

with the convention that the direct product $\mathbb{Z}_{p^{\lambda_r}} \times \cdots \times \mathbb{Z}_{p^{\lambda_k}}$ is trivial for $i \leqslant \lambda_k$.

Our first result gives a precise expression for $|\mathcal{E}_{p^i}(G_\lambda)|$ in the case $k = 2$.

Proposition 2.1. *Let $G_\lambda = \mathbb{Z}_{p^{\lambda_1}} \times \mathbb{Z}_{p^{\lambda_2}}$ with $\lambda_1 \geqslant \lambda_2 \geqslant 1$. Then*

$$|\mathcal{E}_{p^i}(G_\lambda)| = \begin{cases} \frac{p^{i+1}+p^i-2}{p-1}, & 1 \leqslant i \leqslant \lambda_2, \\ \frac{p^{\lambda_2+1}-1}{p-1}, & \lambda_2 < i \leqslant \lambda_1. \end{cases}$$

Proof. We have $\lambda = (\lambda_1, \lambda_2, 0, \dots)$ and consequently

$$\lambda' = (2, 2, \dots, 2, 1, 1, \dots, 1, 0, \dots),$$

where the number of 2's is λ_2 and the number of 1's is $\lambda_1 - \lambda_2$. Then

$$|\mathcal{E}_{p^i}(G_\lambda)| = \sum_{j=0}^{i} N_{i,j},$$

where $N_{i,j}$ denotes the number of subgroups of type (i,j) in G_λ, or equivalently in $\mathbb{Z}_{p^i} \times \mathbb{Z}_{p^i}$ if $i \leqslant \lambda_2$ or in $\mathbb{Z}_{p^i} \times \mathbb{Z}_{p^{\lambda_2}}$ if $\lambda_2 < i \leqslant \lambda_1$.

In the first case we obtain $N_{i,j} = (p+1)p^{i-j-1}$ for $j = 0, 1, \dots, i-1$, and $N_{i,i} = 1$. Therefore

$$|\mathcal{E}_{p^i}(\mathbb{Z}_{p^i} \times \mathbb{Z}_{p^i})| = 1 + \sum_{j=0}^{i-1} (p+1)p^{i-j-1} = \frac{p^{i+1}+p^i-2}{p-1}.$$

In the second case we obtain $N_{i,j} = p^{\lambda_2 - j}$ for $j = 0, 1, \dots, \lambda_2$. Therefore

$$|\mathcal{E}_{p^i}(\mathbb{Z}_{p^i} \times \mathbb{Z}_{p^{\lambda_2}})| = \sum_{j=0}^{\lambda_2} p^{\lambda_2 - j} = \frac{p^{\lambda_2+1}-1}{p-1}.$$

This completes the proof. □

Example 2.1. We have

$$|\mathcal{E}_{p^i}(\mathbb{Z}_{p^4} \times \mathbb{Z}_{p^2})| = \begin{cases} 1, & i = 0, \\ p+2, & i = 1, \\ p^2+2p+2, & i = 2, \\ p^2+p+1, & i = 3 \text{ or } i = 4. \end{cases}$$

In particular, by summing all quantities $|\mathcal{E}_{p^i}(G_\lambda)|$ $(i = 0, 1, \dots, \lambda_1)$, we obtain the total number of subgroups of G_λ (see also [**11**, Prop. 2.9], [**12**, Th. 3.3]).

COROLLARY 2.1. *The total number of subgroups of* $G_\lambda = \mathbb{Z}_{p^{\lambda_1}} \times \mathbb{Z}_{p^{\lambda_2}}$ *where* $\lambda_1 \geqslant \lambda_2 \geqslant 1$, *is*

$$\frac{1}{(p-1)^2}\big[(\lambda_1-\lambda_2+1)p^{\lambda_2+2}-(\lambda_1-\lambda_2-1)p^{\lambda_2+1}-(\lambda_1+\lambda_2+3)p+(\lambda_1+\lambda_2+1)\big]. \tag{2.2}$$

EXAMPLE 2.2. The total number of subgroups of $\mathbb{Z}_{p^4} \times \mathbb{Z}_{p^2}$ is $3p^2 + 5p + 7$.

Now consider the case of rank three p-groups. We need the following lemma.

LEMMA 2.1. *Let* $N_{i,j,\ell}$ *denote the number of subgroups of type* (i, j, ℓ) $(i \geqslant j \geqslant \ell \geqslant 0)$ *in* $G_\lambda = \mathbb{Z}_{p^{\lambda_1}} \times \mathbb{Z}_{p^{\lambda_2}} \times \mathbb{Z}_{p^{\lambda_3}}$ *with* $\lambda_1 \geqslant \lambda_2 \geqslant \lambda_3 \geqslant 1$. *Then*

$$N_{i,j,\ell} = \begin{cases} p^{2i-2\ell-3}(p+1)(p^2+p+1), & \ell < j < i \leqslant \lambda_3, \\ p^{2(i-\ell-1)}(p^2+p+1), & \ell < j = i \leqslant \lambda_3 \text{ or } \ell = j < i \leqslant \lambda_3, \\ 1, & \ell = j = i \leqslant \lambda_3, \\ p^{\lambda_3+i-2\ell-2}(p+1)^2, & \ell < j \leqslant \lambda_3 < i \leqslant \lambda_2, \\ p^{\lambda_3+i-2j-1}(p+1), & \ell = j \leqslant \lambda_3 < i \leqslant \lambda_2, \\ p^{2\lambda_3+i-j-2\ell-1}(p+1), & \ell \leqslant \lambda_3 < j < i \leqslant \lambda_2, \\ p^{2(\lambda_3-\ell)}, & \ell \leqslant \lambda_3 < j = i \leqslant \lambda_2, \\ p^{\lambda_2+2\lambda_3-j-2\ell}, & \ell \leqslant \lambda_3 < j \leqslant \lambda_2 < i \leqslant \lambda_1, \\ p^{\lambda_2+\lambda_3-2\ell-1}(p+1), & \ell < j \leqslant \lambda_3 \leqslant \lambda_2 < i \leqslant \lambda_1, \\ p^{\lambda_2+\lambda_3-2\ell}, & \ell = j \leqslant \lambda_3 \leqslant \lambda_2 < i \leqslant \lambda_1. \end{cases}$$

PROOF. We use Theorem 2.1. We distinguish the following three cases:

I. If $i \leqslant \lambda_3$, then $N_{i,j,\ell}$ is the number of subgroups of type (i, j, ℓ) in $\mathbb{Z}_{p^i} \times \mathbb{Z}_{p^i} \times \mathbb{Z}_{p^i}$. Here we need to consider $\lambda = (i, i, i, 0, \dots)$ with $\lambda' = (3, 3, \dots, 3, 0 \dots)$, where the number of 3's is i, and $\mu = (i, j, \ell, 0, \dots)$ with

$$\mu' = (3, 3, \dots, 3, 2, 2, \dots, 2, 1, 1, \dots, 1, 0, \dots),$$

where the number of 3's is ℓ, the number of 2's is $j - \ell$, the number of 1's is $i - j$. In the subcase $\ell < j < i \leqslant \lambda_3$ we deduce

$$N_{i,j,\ell} = (p^2)^{j-\ell-1}p(p+1)(p^2)^{i-j-1}(p^2+p+1) = p^{2i-2\ell-3}(p+1)(p^2+p+1).$$

The subcases $\ell < j = i \leqslant \lambda_3$, $\ell = j < i \leqslant \lambda_3$ and $\ell = j = i \leqslant \lambda_3$ are treated similar.

II. If $\lambda_3 < i \leqslant \lambda_2$, then $N_{i,j,\ell}$ is the number of subgroups of type (i, j, ℓ) in $\mathbb{Z}_{p^i} \times \mathbb{Z}_{p^i} \times \mathbb{Z}_{p^{\lambda_3}}$. We consider $\lambda = (i, i, \lambda_3, 0, \dots)$ with $\lambda' = (3, 3, \dots, 3, 2, 2, \dots, 2, 0 \dots)$ where the number of 3's is λ_3, the number of 2's is $i-\lambda_3$, and μ, μ' like in the case I.

For example, in the subcase $\ell = j \leqslant \lambda_3 < i \leqslant \lambda_2$ we obtain

$$N_{i,j,\ell} = (p^2)^{\lambda_3-j}p^{i-\lambda_3-1}(p+1) = p^{\lambda_3+i-2j-1}(p+1).$$

III. If $\lambda_2 < i \leqslant \lambda_1$, then $N_{i,j,\ell}$ is the number of subgroups of type (i, j, ℓ) in $\mathbb{Z}_{p^i} \times \mathbb{Z}_{p^{\lambda_2}} \times \mathbb{Z}_{p^{\lambda_3}}$. We consider $\lambda = (i, \lambda_2, \lambda_3, 0, \dots)$ with

$$\lambda' = (3, 3, \dots, 3, 2, 2, \dots, 2, 1, 1, \dots, 1, 0 \dots),$$

where the number of 3's is λ_3, the number of 2's is $\lambda_2-\lambda_3$, the number of 1's is $i-\lambda_2$, and μ, μ' like in the case I. $\square$

PROPOSITION 2.2. *Let* $G_\lambda=\mathbb{Z}_{p^{\lambda_1}}\times\mathbb{Z}_{p^{\lambda_2}}\times\mathbb{Z}_{p^{\lambda_3}}$ *with* $\lambda_1\geqslant\lambda_2\geqslant\lambda_3\geqslant 1$. *Then*

$$|\mathcal{E}_{p^i}(G_\lambda)|=\begin{cases}\frac{p^{2i-1}((i+1)p^4+(i-1)p^3-p^2-(i+2)p-i)+3}{(p^2-1)(p-1)}, & 1\leqslant i\leqslant\lambda_3,\\ \frac{(\lambda_3+1)p^{\lambda_3+i}(p^2-1)(p+1)-2p^{2\lambda_3+2}+2}{(p^2-1)(p-1)}, & \lambda_3<i\leqslant\lambda_2,\\ \frac{(\lambda_3+1)p^{\lambda_2+\lambda_3+1}(p^2-1)-p^{2\lambda_3+2}+1}{(p^2-1)(p-1)}, & \lambda_2<i\leqslant\lambda_1.\end{cases}$$

PROOF. We have $|\mathcal{E}_{p^i}(G_\lambda)|=\sum_{0\leqslant\ell\leqslant j\leqslant i}N_{i,j,\ell}$ and use Lemma 2.1.

Case I. If $i\leqslant\lambda_3$, then

$$\begin{aligned}|\mathcal{E}_{p^i}(G_\lambda)|&=N_{i,i,i}+\sum_{\ell=0}^{i-1}\sum_{j=\ell+1}^{i-1}N_{i,j,\ell}+\sum_{\ell=0}^{i-1}N_{i,i,\ell}+\sum_{\ell=0}^{i-1}N_{i,\ell,\ell}\\&=1+\sum_{\ell=0}^{i-1}\sum_{j=\ell+1}^{i-1}p^{2i-2\ell-3}(p+1)(p^2+p+1)+2\sum_{\ell=0}^{i-1}p^{2(i-\ell-1)}(p^2+p+1)\\&=1+\frac{p^2+p+1}{(p^2-1)(p-1)}((i-1)p^{2i+1}-ip^{2i-1}+p)+2(p^2+p+1)\frac{p^{2i}-1}{p^2-1}\\&=\frac{p^{2i-1}((i+1)p^4+(i-1)p^3-p^2-(i+2)p-i)+3}{(p^2-1)(p-1)},\end{aligned}$$

by direct computations.

II. If $\lambda_3<i\leqslant\lambda_2$, then we have

$$\begin{aligned}|\mathcal{E}_{p^i}(G_\lambda)|&=\sum_{\ell=0}^{\lambda_3}\sum_{j=\ell+1}^{\lambda_3}N_{i,j,\ell}+\sum_{\ell=0}^{\lambda_3}\sum_{j=\lambda_3+1}^{i-1}N_{i,j,\ell}+\sum_{\ell=0}^{\lambda_3}N_{i,i,\ell}+\sum_{\ell=0}^{\lambda_3}N_{i,\ell,\ell}\\&=\sum_{\ell=0}^{\lambda_3}\sum_{j=\ell+1}^{\lambda_3}p^{\lambda_3+i-2\ell-2}(p+1)^2+\sum_{\ell=0}^{\lambda_3}\sum_{j=\lambda_3+1}^{i-1}p^{2\lambda_3+i-j-2\ell-1}(p+1)\\&\quad+\sum_{\ell=0}^{\lambda_3}p^{2(\lambda_3-\ell)}+\sum_{\ell=0}^{\lambda_3}p^{\lambda_3+i-2\ell-1}(p+1)\\&=\frac{\lambda_3p^{i-2}(p+1)^2+p^{\lambda_3}+p^i+p^{i-1}}{p^{\lambda_3}(p^2-1)}(p^{2\lambda_3+2}-1)\\&\quad-\frac{p^{i-2}(p^{2\lambda_3+2}-(\lambda_3+1)p^2+\lambda_3)}{p^{\lambda_3}(p-1)^2}+\frac{(p^{2\lambda_3+2}-1)(p^{i-\lambda_3-1}-1)}{(p-1)^2},\end{aligned}$$

which gives the above formula.

III. Finally, if $\lambda_2<i\leqslant\lambda_1$, then

$$|\mathcal{E}_{p^i}(G_\lambda)|=\sum_{\ell=0}^{\lambda_3}\sum_{j=\lambda_3+1}^{\lambda_2}N_{i,j,\ell}+\sum_{\ell=0}^{\lambda_3-1}\sum_{j=\ell+1}^{\lambda_3}N_{i,j,\ell}+\sum_{\ell=0}^{\lambda_3}N_{i,\ell,\ell}$$

$$
\begin{aligned}
&= \sum_{\ell=0}^{\lambda_3} \sum_{j=\lambda_3+1}^{\lambda_2} p^{\lambda_2+2\lambda_3-j-2\ell} + \sum_{\ell=0}^{\lambda_3-1} \sum_{j=\ell+1}^{\lambda_3} p^{\lambda_2+\lambda_3-2\ell-1}(p+1) + \sum_{\ell=0}^{\lambda_3} p^{\lambda_2+\lambda_3-2\ell} \\
&= \frac{(p^{2\lambda_3+2}-1)(p^{\lambda_2-\lambda_3}-1)}{(p^2-1)(p-1)} + \frac{p^{\lambda_2-\lambda_3+1}}{(p^2-1)(p-1)} \lambda_3 p^{2\lambda_3+2} - (\lambda_3+1)p^{2\lambda_3}+1) \\
&\quad + p^{\lambda_2-\lambda_3} \frac{p^{2\lambda_3+2}-1}{p^2-1},
\end{aligned}
$$

leading to the given result. □

Note that Proposition 2.2 is valid also in the case $\lambda_3 = 0$, when it reduces to Proposition 2.1.

Example 2.3. We have

$$
|\mathcal{E}_{p^i}(\mathbb{Z}_{p^4} \times \mathbb{Z}_{p^2} \times \mathbb{Z}_{p^2})| = \begin{cases} 1, & i = 0, \\ 2p^2 + 2p + 3, & i = 1, \\ 3p^4 + 4p^3 + 6p^2 + 3p + 3, & i = 2, \\ 3p^4 + 2p^3 + 2p^2 + p + 1, & i = 3 \text{ or } i = 4. \end{cases}
$$

Corollary 2.2. *The total number of subgroups of* $G_\lambda = \mathbb{Z}_{p^{\lambda_1}} \times \mathbb{Z}_{p^{\lambda_2}} \times \mathbb{Z}_{p^{\lambda_3}}$, *where* $\lambda_1 \geqslant \lambda_2 \geqslant \lambda_3 \geqslant 1$, *is*

$$
\frac{A}{(p^2-1)^2(p-1)},
$$

where

$$
\begin{aligned}
A = {} & (\lambda_3+1)(\lambda_1-\lambda_2+1)p^{\lambda_2+\lambda_3+5} + 2(\lambda_3+1)p^{\lambda_2+\lambda_3+4} \\
& - 2(\lambda_3+1)(\lambda_1-\lambda_2)p^{\lambda_2+\lambda_3+3} - 2(\lambda_3+1)p^{\lambda_2+\lambda_3+2} \\
& + (\lambda_3+1)(\lambda_1-\lambda_2-1)p^{\lambda_2+\lambda_3+1} - (\lambda_1+\lambda_2-\lambda_3+3)p^{2\lambda_3+4} \\
& - 2p^{2\lambda_3+3} + (\lambda_1+\lambda_2-\lambda_3-1)p^{2\lambda_3+2} \\
& + (\lambda_1+\lambda_2+\lambda_3+5)p^2 + 2p - (\lambda_1+\lambda_2+\lambda_3+1).
\end{aligned}
$$

Proof. The total number of subgroups of G_λ is

$$
\sum_{i=0}^{\lambda_1} |\mathcal{E}_{p^i}(G_\lambda)| = \sum_{0 \leqslant i \leqslant \lambda_3} |\mathcal{E}_{p^i}(G_\lambda)| + \sum_{\lambda_3 < i \leqslant \lambda_2} |\mathcal{E}_{p^i}(G_\lambda)| + \sum_{\lambda_2 < i \leqslant \lambda_1} |\mathcal{E}_{p^i}(G_\lambda)|
$$

and summing the quantities given in Proposition 2.2 we deduce the result. □

We remark that an equivalent formula to that given in Corollary 2.2 was obtained in [**8**, Cor. 2.2] by using different arguments. Corollary 2.2 is valid also in the case $\lambda_3 = 0$, when it reduces to Corollary 2.1.

Example 2.4. The total number of subgroups of $\mathbb{Z}_{p^4} \times \mathbb{Z}_{p^2} \times \mathbb{Z}_{p^2}$ is $9p^4 + 8p^3 + 12p^2 + 7p + 9$.

Next consider the number of subgroups of exponent p (that is, the number of elementary abelian subgroups) in G_λ, which equals the total number of nontrivial subgroups of $(\mathbb{Z}_p)^k$. Since $(\mathbb{Z}_p)^k$ is a k–dimensional linear space over $\mathbb{Z}_p$, the number in question is exactly the total number of nonzero subspaces, which is, as well-known, $\sum_{i=1}^k \binom{k}{i}_p$ (Galois number). So we have the following result.

PROPOSITION 2.3. *Let $G_\lambda = \mathbb{Z}_{p^{\lambda_1}} \times \mathbb{Z}_{p^{\lambda_2}} \times \cdots \times \mathbb{Z}_{p^{\lambda_k}}$ with $\lambda_1 \geqslant \lambda_2 \geqslant \cdots \geqslant \lambda_k \geqslant 1$. Then $|\mathcal{E}_p(G_\lambda)| = \sum_{r=1}^k \binom{k}{r}_p$.*

We give a direct proof of this formula based on Theorem 2.1.

PROOF. We use Theorem 2.1 in the case $\lambda = (1, 1, \dots, 1, 0, \dots)$, where the number of 1's is k, and $\mu = (1, 1, \dots, 1, 0, \dots)$, where the number of 1's is r with $1 \leqslant r \leqslant k$. Here $\lambda' = (k, 0, 0, \dots)$, $\mu' = (r, 0, 0, \dots)$ and obtain that the number of subgroups of type μ is $\binom{k}{r}_p$, while the number of subgroups of exponent 1 is exactly $\sum_{r=1}^k \binom{k}{r}_p$. □

EXAMPLE 2.5. We have for any $\lambda_1 \geqslant \lambda_2 \geqslant \lambda_3 \geqslant \lambda_4 \geqslant 1$,

$$|\mathcal{E}_p(\mathbb{Z}_{p^{\lambda_1}} \times \mathbb{Z}_{p^{\lambda_2}} \times \mathbb{Z}_{p^{\lambda_3}} \times \mathbb{Z}_{p^{\lambda_4}})| = p^4 + 3p^3 + 4p^2 + 3p + 4.$$

Concerning the number of subgroups of exponent p^2 in G_λ, we have the following formula.

PROPOSITION 2.4. *Let $G_\lambda = \mathbb{Z}_{p^{\lambda_1}} \times \mathbb{Z}_{p^{\lambda_2}} \times \cdots \times \mathbb{Z}_{p^{\lambda_k}}$ with $\lambda_1 \geqslant \lambda_2 \geqslant \cdots \geqslant \lambda_t > \lambda_{t+1} = \cdots = \lambda_k = 1$, where $0 \leqslant t \leqslant k$ is fixed ($t = 0$ if each λ_j is 1 and $t = k$ if each λ_j is $\geqslant 2$). Then*

$$|\mathcal{E}_{p^2}(G_\lambda)| = \sum_{\substack{1 \leqslant r \leqslant t \\ 0 \leqslant s \leqslant k-r}} p^{r(k-r-s)} \binom{k-r}{s}_p \binom{t}{r}_p,$$

which is zero (empty sum) for $t = 0$.

PROOF. Let $t \geqslant 1$. Use Theorem 2.1 for $\lambda = (2, 2, \dots, 2, 1, 1, \dots, 1, 0, \dots)$, where the number of 2's is t and the number of 1's is $k - t$ and

$$\mu = (2, 2, \dots, 2, 1, 1, \dots, 1, 0, \dots),$$

where the number of 2's is r and the number of 1's is s with $1 \leqslant r \leqslant t$, $0 \leqslant s \leqslant k-r$. Now $\lambda' = (k, t, 0, \dots)$, $\mu' = (r+s, r, 0, \dots)$ and obtain that the number of subgroups of type μ is $p^{r(k-r-s)}\binom{k-r}{s}_p\binom{t}{r}_p$ and the number of subgroups of exponent p^2 is deduced by summing over r and s. □

EXAMPLE 2.6. ($k = 4$, $t = 2$) We have

$$|\mathcal{E}_{p^2}(\mathbb{Z}_{p^4} \times \mathbb{Z}_{p^2} \times \mathbb{Z}_p \times \mathbb{Z}_p)| = p^5 + 5p^4 + 6p^3 + 4p^2 + 2p + 2.$$

In what follows, let $G_\lambda = \mathbb{Z}_{p^{\lambda_1}} \times \mathbb{Z}_{p^{\lambda_2}} \times \cdots \times \mathbb{Z}_{p^{\lambda_k}}$ and $G_\kappa = \mathbb{Z}_{p^{\kappa_1}} \times \mathbb{Z}_{p^{\kappa_2}} \times \cdots \times \mathbb{Z}_{p^{\kappa_\ell}}$ be two finite abelian p-groups, where $\lambda_1 \geqslant \lambda_2 \geqslant \cdots \geqslant \lambda_k \geqslant 1$ and $\kappa_1 \geqslant \kappa_2 \geqslant \cdots \geqslant \kappa_\ell \geqslant 1$. Assume that G_λ and G_κ have the same number of subgroups of exponent p^i for every i, i.e., $|\mathcal{E}_{p^i}(G_\lambda)| = |\mathcal{E}_{p^i}(G_\kappa)|$ ($i \geqslant 0$). Then

$\lambda_1 = \kappa_1$. On the other hand, since the function $f : \mathbb{N}^* \to \mathbb{N}^*$, $f(k) = \sum_{i=1}^{k} \binom{k}{i}_p$ is one–to–one, by Proposition 2.3 we infer that $k = \ell$. Clearly, if $k = 1$ one obtains $G_\lambda \cong G_\kappa$. The same thing can be also said for $k = 2$ and $k = 3$ by Propositions 2.1 and 2.2. Inspired by these remarks, we state and prove the following result.

PROPOSITION 2.5. *Two finite abelian p-groups G_λ and G_κ are isomorphic if and only if they have the same number of subgroups of exponent p^i, for every $i \geqslant 0$.*

PROOF. Let $\lambda = (\lambda_1, \lambda_2, \dots, \lambda_k, 0, \dots)$ and $\kappa = (\kappa_1, \kappa_2, \dots, \kappa_\ell, 0, \dots)$, where $\lambda_k, \kappa_\ell > 0$, be partitions such that for p-groups G_λ and G_κ one has $|\mathcal{E}_{p^i}(G_\lambda)| = |\mathcal{E}_{p^i}(G_\kappa)|$, for every $i \geqslant 0$. As noted before, $\lambda_1 = \kappa_1$ and $k = \ell$ hold true.

Let us define $\lambda_{k+1} = \kappa_{k+1} = 0$. We prove, by reverse induction on $t \leqslant k+1$ that $\lambda_t = \kappa_t$. So, let us assume that $\lambda_i = \kappa_i$ for all $k+1 \geqslant i \geqslant t+1$, and prove that $\lambda_t = \kappa_t$.

Suppose to the contrary that w.l.o.g. $\lambda_t > \kappa_t$. Since $\lambda_1 = \kappa_1$, one has $t > 1$. Consider the partitions λ^{λ_t} and κ^{λ_t} defined with

$$\lambda^{\lambda_t} = (\underbrace{\lambda_t, \dots, \lambda_t}_{t}, \gamma_{t+1}, \dots, \gamma_k, 0, \dots),$$
$$\kappa^{\lambda_t} = (\underbrace{\lambda_t, \dots, \lambda_t}_{s}, \kappa_{s+1}, \dots, \kappa_t, \gamma_{t+1}, \dots, \gamma_k, 0, \dots),$$

where $\gamma_i = \lambda_i = \kappa_i$, for $i \geqslant t+1$, and $s = \max\{j : \kappa_j \geqslant \lambda_t\}$. Note that by our assumption $s < t$, and since $\kappa_1 = \lambda_1 \geqslant \lambda_t$ also $s \geqslant 1$.

From the definition of numbers $\alpha_\omega(\mu; p)$, it is clear that for any three partitions μ, σ, τ, which satisfy $\mu \preceq \sigma \preceq \tau$, we have $\alpha_\sigma(\mu; p) \leqslant \alpha_\tau(\mu; p)$.

Using this remark, the fact that $\kappa^{\lambda_t} \precneqq \lambda^{\lambda_t}$ and $\alpha_{\lambda^{\lambda_t}}(\lambda^{\lambda_t}; p) = 1$, one has

$$\begin{aligned} 1 + |\mathcal{E}_{p^{\lambda_t}}(G_\kappa)| &= 1 + |\mathcal{E}_{p^{\lambda_t}}(G_{\kappa^{\lambda_t}})| = 1 + \sum_{\mu \preceq \kappa^{\lambda_t},\ \mu_1 = \lambda_t} \alpha_{\kappa^{\lambda_t}}(\mu; p) \\ &\leqslant \alpha_{\lambda^{\lambda_t}}(\lambda^{\lambda_t}; p) + \sum_{\mu \preceq \kappa^{\lambda_t},\ \mu_1 = \lambda_t} \alpha_{\lambda^{\lambda_t}}(\mu; p) \leqslant \sum_{\mu \preceq \lambda^{\lambda_t},\ \mu_1 = \lambda_t} \alpha_{\lambda^{\lambda_t}}(\mu; p) \\ &= |\mathcal{E}_{p^{\lambda_t}}(G_{\lambda^{\lambda_t}})| = |\mathcal{E}_{p^{\lambda_t}}(G_\lambda)|, \end{aligned}$$

a contradiction. □

COROLLARY 2.3. *Two arbitrary finite abelian groups are isomorphic if and only if they have the same number of subgroups of any exponent.*

Finally, we note that another interesting problem is to find the polynomial $|\mathcal{E}_{p^i}(G_\lambda)|$ in the case $k = 4$.

3. Second approach

We need the following result giving the representation of subgroups of the group $\mathbb{Z}_m \times \mathbb{Z}_n$. For every $m, n \in \mathbb{N}^*$ let

$$J_{m,n} := \Big\{(a, b, c, d, \ell) \in (\mathbb{N}^*)^5 : a \mid m, b \mid a, c \mid n, d \mid c, \frac{a}{b} = \frac{c}{d},$$

$$\ell \leqslant \frac{a}{b},\ \gcd\Big(\ell, \frac{a}{b}\Big) = 1\Big\}.$$

Note that here $\gcd(b,d)\cdot \operatorname{lcm}(a,c) = ad$ and $\gcd(b,d) \mid \operatorname{lcm}(a,c)$.

For $(a,b,c,d,\ell) \in J_{m,n}$ define

$$K_{a,b,c,d,\ell} := \Big\{\Big(i\frac{m}{a}, i\ell\frac{n}{c} + j\frac{n}{d}\Big) : 0 \leqslant i \leqslant a-1, 0 \leqslant j \leqslant d-1\Big\}.$$

THEOREM 3.1. [**14**, Th. 3.1] *Let $m,n \in \mathbb{N}^*$.*

i) *The map $(a,b,c,d,\ell) \mapsto K_{a,b,c,d,\ell}$ is a bijection between the set $J_{m,n}$ and the set of subgroups of $(\mathbb{Z}_m \times \mathbb{Z}_n, +)$.*

ii) *The invariant factor decomposition of the subgroup $K_{a,b,c,d,\ell}$ is*

$$K_{a,b,c,d,\ell} \simeq \mathbb{Z}_{\gcd(b,d)} \times \mathbb{Z}_{\operatorname{lcm}(a,c)}.$$

iii) *The order of the subgroup $K_{a,b,c,d,\ell}$ is ad and its exponent is $\operatorname{lcm}(a,c)$.*

Let $s_E(m,n)$ stand for the number of subgroups of exponent E of the group $\mathbb{Z}_m \times \mathbb{Z}_n$.

PROPOSITION 3.1. *For every $m,n \in \mathbb{N}^*$, $E \mid \operatorname{lcm}(m,n)$ we have*

$$s_E(m,n) = \sum_{\substack{i\mid m, j\mid n\\ \operatorname{lcm}(i,j)=E}} \gcd(i,j) \tag{3.1}$$

$$= \frac{1}{E}\sum_{\substack{i\mid m, j\mid n\\ \operatorname{lcm}(i,j)=E}} ij. \tag{3.2}$$

PROOF. According to Theorem 3.1,

$$s_E(m,n) = \sum_{\substack{a\mid m\\ b\mid a}}\sum_{\substack{c\mid n\\ d\mid c}}\sum_{\substack{a/b=c/d=e\\ \operatorname{lcm}(a,c)=E}} \phi(e),$$

where ϕ is Euler's totient function. This can be written (with $m = ax$, $a = by$, $n = cz$, $c = dt$) as

$$s_E(m,n) = \sum_{\substack{bxe=m\\ dze=n\\ e\operatorname{lcm}(b,d)=E}} \phi(e) = \sum_{\substack{ix=m\\ jz=n}}\ \sum_{\substack{be=i\\ de=j\\ e\operatorname{lcm}(b,d)=E}} \phi(e)$$

$$= \sum_{\substack{i\mid m\\ j\mid n\\ \operatorname{lcm}(i,j)=E}}\ \sum_{e\mid\gcd(i,j)} \phi(e) = \sum_{\substack{i\mid m\\ j\mid n\\ \operatorname{lcm}(i,j)=E}} \gcd(i,j),$$

which is (3.1). Formula (3.2) is its immediate consequence. □

REMARK 3.1. Proposition 2.1 is a direct consequence of the above result. The total number $s(m,n)$ of subgroups of the group $\mathbb{Z}_m \times \mathbb{Z}_n$ is (see [**6**, Th. 3], [**14**, Th. 4.1])

$$s(m,n) = \sum_{i\mid m, j\mid n} \gcd(i,j) \tag{3.3}$$

and (3.1) shows the distribution of the number of subgroups according to their exponents. Formula (2.2) can be obtained also by using (3.3).

EXAMPLE 3.1. The total number of subgroups of $\mathbb{Z}_{12}\times\mathbb{Z}_{18}$ is $s(12,18)=80$ and we have $s_1(12,18)=1$, $s_2(12,18)=4$, $s_3(12,18)=5$, $s_4(12,18)=3$, $s_6(12,18)=20$, $s_9(12,18)=4$, $s_{12}(12,18)=15$, $s_{18}(12,18)=16$, $s_{36}(12,18)=12$.

COROLLARY 3.1 ($m=n$). *For every $n\in\mathbb{N}^*$ and $E\mid n$,*

$$s_E(n,n)=\sum_{\substack{i\mid E,j\mid E\\ \gcd(E/i,E/j)=1}}\gcd(i,j), \tag{3.4}$$

which equals the number of cyclic subgroups of the group $\mathbb{Z}_E\times\mathbb{Z}_E$.

The fact that $s_E(n,n)$ equals the number of cyclic subgroups of the group $\mathbb{Z}_E\times\mathbb{Z}_E$, but without deriving formula (3.4) is [**6**, Th. 8], proved by using different arguments.

PROOF. In the case $m=n$, for every $E\mid n$ we have by (3.1),

$$s_E(n,n)=\sum_{\substack{i\mid n,j\mid n\\ \mathrm{lcm}(i,j)=E}}\gcd(i,j)=\sum_{\substack{ia=E,jb=E\\ \mathrm{lcm}(E/a,E/b)=E}}\gcd(i,j)=\sum_{\substack{ia=E,jb=E\\ \gcd(a,b)=1}}\gcd(i,j),$$

giving (3.4), which equals the number of cyclic subgroups of the group $\mathbb{Z}_E\times\mathbb{Z}_E$ by [**6**, eq. (16)]. □

PROPOSITION 3.2. *For every $m,n\in\mathbb{N}^*$ the sum of exponents of the subgroups of $\mathbb{Z}_m\times\mathbb{Z}_n$ is $\sigma(m)\sigma(n)$, where $\sigma(k)=\sum_{d\mid k}d$.*

PROOF. By (3.2) the sum of exponents of the subgroups of $\mathbb{Z}_m\times\mathbb{Z}_n$ is

$$\sum_{E\mid\mathrm{lcm}(m,n)}Es_E(m,n)=\sum_{E\mid\mathrm{lcm}(m,n)}E\cdot\frac{1}{E}\sum_{\substack{i\mid m,j\mid n\\ \mathrm{lcm}(i,j)=E}}ij=\sum_{\substack{i\mid m,j\mid n\\ \mathrm{lcm}(i,j)\mid\mathrm{lcm}(m,n)}}ij$$
$$=\sum_{i\mid m}i\sum_{j\mid n}j=\sigma(m)\sigma(n).$$ □

By Proposition 3.2, the arithmetic mean of exponents of the subgroups of $\mathbb{Z}_m\times\mathbb{Z}_n$ is $\sigma(m)\sigma(n)/s(m,n)$, where $s(m,n)$ is given by (3.3). Now consider the case $m=n$. Let $A(n)$ stand for the arithmetic mean of exponents of the subgroups of $\mathbb{Z}_n\times\mathbb{Z}_n$. We have

$$A(n)=\frac{\sigma(n)^2}{s(n)},\quad\text{where } s(n):=s(n,n). \tag{3.5}$$

Recall that a function $f:\mathbb{N}^*\to\mathbb{C}$ is said to be multiplicative if $f(nn')=f(n)f(n')$ whenever $\gcd(n,n')=1$. It is well known that the sum-of-divisors function σ is multiplicative. The function $s(n)=\sum_{i,j\mid n}\gcd(i,j)$ is also multiplicative,

as shown by the following direct proof: Let $\gcd(n,n')=1$. Then

$$s(nn') = \sum_{i,j|nn'} \gcd(i,j) = \sum_{\substack{a,b|n\\ a',b'|n'}} \gcd(aa',bb')$$
$$= \sum_{a,b|n} \gcd(a,b) \sum_{a',b'|n'} \gcd(a',b') = s(n)s(n').$$

We conclude that the function A given by (3.5) is multiplicative and and for every prime power p^ν,

$$A(p^\nu) = \frac{(p^{\nu+1}-1)^2}{p^{\nu+2}+p^{\nu+1}-(2\nu+3)p+2\nu+1},$$

cf. Corollary 2.1.

Since the exponent of every subgroup of $\mathbb{Z}_n \times \mathbb{Z}_n$ is a divisor of n, whence $\leqslant n$, we deduce that $A(n) \leqslant n$ ($n \in \mathbb{N}^*$). For the function $n \mapsto f(n) := A(n)/n \in (0,1]$ the series taken over the primes

$$\sum_p \frac{1-f(p)}{p} = \sum_p \frac{p-1}{p^2(p+3)}$$

is convergent, and it follows from a theorem of H. Delange (see, e.g., [**9**]) that the function f has a nonzero mean value M given by

$$M := \lim_{x\to\infty} \frac{1}{x} \sum_{n\leqslant x} f(n) = \prod_p \Big(1-\frac{1}{p}\Big) \sum_{\nu=0}^{\infty} \frac{f(p^\nu)}{p^\nu}$$
$$= \prod_p \Big(1-\frac{1}{p}\Big) \sum_{\nu=0}^{\infty} \frac{(p^{\nu+1}-1)^2}{p^{2\nu}(p^{\nu+2}+p^{\nu+1}-(2\nu+3)p+2\nu+1)}.$$

the products being over the primes.

We prove the following more exact result.

Proposition 3.3. *We have*

$$\sum_{n\leqslant x} A(n) = \frac{M}{2}x^2 + O(x\log^3 x). \tag{3.6}$$

Proof. Let $f(n) = \sum_{d|n} g(d)$ ($n \in \mathbb{N}^*$), that is $g = \mu * f$ in terms of the Dirichlet convolution, where μ is the Möbius function. Here $g(p^\nu) = f(p^\nu) - f(p^{\nu-1})$ for every prime power p^ν ($\nu \in \mathbb{N}^*$). Note that

$$A(p^k) = \frac{(\sum_{i=0}^k p^i)^2}{\sum_{i=0}^k (2i+1)p^{k-i}} \quad (k \geqslant 0),$$

so if we denote $S = \sum_{i=0}^{\nu-1} p^i$ and $T = \sum_{i=0}^{\nu-1}(2i+1)p^{\nu-1-i}$, we have

$$g(p^\nu) = \frac{(Sp+1)^2}{p^\nu(Tp+2\nu+1)} - \frac{S^2}{p^{\nu-1}T} = \frac{2STp+T-pS^2(2\nu+1)}{p^\nu T(Tp+2\nu+1)}.$$

Since $S \leqslant T$ and $T \leqslant (2\nu-1)S$, we have

$$\begin{aligned}|g(p^\nu)| &< \frac{\max\{2STp+T-pS^2(2\nu-1), pS^2(2\nu+1)-2STp\}}{p^\nu T(Tp+2\nu+1)}\\ &\leqslant \frac{1}{p^\nu}\max\left\{\frac{STp+T}{T(Tp+1)}, \frac{pS^2(2\nu-1)}{T^2p}\right\}\\ &\leqslant \frac{1}{p^\nu}\max\{1, 2\nu-1\} = \frac{2\nu-1}{p^\nu},\end{aligned}$$

valid for every prime power p^ν ($\nu \in \mathbb{N}^*$). Hence, $|g(n)| \leqslant \tau(n^2)/n$ for every $n \geqslant 1$, where $\tau(k)$ stands for the number of positive divisors of k.

We deduce that

$$\begin{aligned}\sum_{n\leqslant x} f(n) &= \sum_{de\leqslant x} g(d) = \sum_{d\leqslant x} g(d) \sum_{e\leqslant x/d} 1 = \sum_{d\leqslant x} g(d)(x/d + O(1))\\ &= x\sum_{d=1}^{\infty}\frac{g(d)}{d} + O\Bigl(x\sum_{d>x}\frac{|g(d)|}{d}\Bigr) + O\Bigl(\sum_{d\leqslant x}|g(d)|\Bigr)\\ &= Mx + O\Bigl(x\sum_{d>x}\frac{\tau(d^2)}{d^2}\Bigr) + O\Bigl(\sum_{d\leqslant x}\frac{\tau(d^2)}{d}\Bigr),\end{aligned}$$

where in the main term the coefficient of x is M by Euler's product formula. It is known that $\sum_{n\leqslant x}\tau(n^2) = cx\log^2 x + O(x\log x)$ with a certain constant c and partial summation shows that $\sum_{n>x}\tau(n^2)/n^2 = O((\log^2 x)/x)$, $\sum_{n\leqslant x}\tau(n^2)/n = O(\log^3 x)$. Therefore,

$$\sum_{n\leqslant x} f(n) = Mx + O(\log^3 x). \tag{3.7}$$

Now (3.6) follows from (3.7) by partial summation. □

Formula (3.6) and its proof are similar to those of [**13**, Th. 3.1.3].

Acknowledgements. The authors thank the referee for very careful reading of the manuscript, many useful comments and for the proof of Proposition 2.5.

References

1. G. Bhowmik, *Evaluation of divisor functions of matrices*, Acta Arith. **74** (1996), 155–159.
2. M. L. Butler, *Subgroup Lattices and Symmetric Functions*, Mem. Am. Math. Soc. **112**(539), 1994.
3. G. Călugăreanu, *The total number of subgroups of a finite abelian group*, Sci. Math. Jpn. **60**(1) (2004), 157–167.
4. S. Delsarte, *Fonctions de Möbius sur les groupes abeliens finis*, Annals of Math. **49** (1948), 600–609.
5. P. E. Djubjuk, *On the number of subgroups of a finite abelian group*, Izv. Akad. Nauk SSSR, Ser. Mat. **12** (1948), 351–378.
6. M. Hampejs, N. Holighaus, L. Tóth, C. Wiesmeyr, *Representing and counting the subgroups of the group $\mathbb{Z}_m \times \mathbb{Z}_n$*, J. Numbers **2014**, Article ID 491428.
7. M. Hampejs, L. Tóth, *On the subgroups of finite abelian groups of rank three*, Ann. Univ. Sci. Budap. Rolando Eötvös, Sect. Comput. **39** (2013), 111–124.

8. J.-M. Oh, *An explicit formula for the number of subgroups of a finite abelian p-group up to rank* 3, Commun. Korean Math. Soc. **28** (2013), 649–667.
9. A. G. Postnikov, *Introduction to Analytic Number Theory*, Transl. Math. Monogr. **68**, American Mathematical Society, Providence, RI, 1988.
10. M. Suzuki, *Group Theory*, vol. I, II, Springer-Verlag, Berlin, 1982, 1986.
11. M. Tărnăuceanu, *A new method of proving some classical theorems of abelian groups*, Southeast Asian Bull. Math. **31**(6) (2007), 1191–1203.
12. ———, *An arithmetic method of counting the subgroups of a finite abelian group*, Bull. Math. Soc. Sci. Math. Roum., Nouv. Sér. **53**(101) (2010), 373–386.
13. M. Tărnăuceanu, L. Tóth, *Cyclicity degrees of finite groups*, Acta Math. Hung. **145** (2015), 489–504.
14. L. Tóth, *Subgroups of finite abelian groups having rank two via Goursat's lemma*, Tatra Mt. Math. Publ. **59** (2014), 93–103.
15. Y. Yeh, *On prime power abelian groups*, Bull. Am. Math. Soc. **54** (1948), 323–327.

Faculty of Mathematics
"Al. I. Cuza" University
Iaşi
Romania
tarnauc@uaic.ro

(Received 02 07 2015)
(Revised 14 03 2016)

Department of Mathematics
University of Pécs
Pécs
Hungary
ltoth@gamma.ttk.pte.hu