PUBLICATIONS DE L'INSTITUT MATHÉMATIQUE
Nouvelle série, tome 102(116) (2017), 107–113 DOI: https://doi.org/10.2298/PIM1716107Y

A NEW THEOREM ON ABSOLUTE MATRIX SUMMABILITY OF FOURIER SERIES

Şebnem Yildiz

Abstract. We generalize a main theorem dealing with absolute weighted mean summability of Fourier series to the $|A, p_n|_k$ summability factors of Fourier series under weaker conditions. Also some new and known results are obtained.

1. Introduction

Let $\sum a_n$ be a given infinite series with partial sums (s_n). By u_n^α and t_n^α we denote the nth Cesàro means of order α, with $\alpha > -1$, of the sequence (s_n) and (na_n), respectively, that is (see [**6**])

$$u_n^\alpha = \frac{1}{A_n^\alpha}\sum_{v=0}^{n} A_{n-v}^{\alpha-1}s_v \quad \text{and} \quad t_n^\alpha = \frac{1}{A_n^\alpha}\sum_{v=0}^{n} A_{n-v}^{\alpha-1}va_v,$$

where

$$A_n^\alpha = \frac{(\alpha+1)(\alpha+2)...(\alpha+n)}{n!} = O(n^\alpha), \qquad A_{-n}^\alpha = 0 \quad \text{for} \quad n > 0.$$

The series $\sum a_n$ is said to be summable $|C,\alpha|_k$, $k \geqslant 1$, if (see [**8**, **10**])

$$\sum_{n=1}^{\infty} n^{k-1}|u_n^\alpha - u_{n-1}^\alpha|^k = \sum_{n=1}^{\infty}\frac{1}{n}|t_n^\alpha|^k < \infty.$$

If we take $\alpha = 1$, then $|C,\alpha|_k$ summability reduces to $|C,1|_k$ summability.

Let (p_n) be a sequence of positive real numbers such that

$$P_n = \sum_{v=0}^{n} p_v \to \infty \qquad \text{as} \quad n \to \infty, \quad (P_{-i} = p_{-i} = 0, \quad i \geqslant 1).$$

2010 *Mathematics Subject Classification:* 26D15; 42A24; 40F05; 40G99.

Key words and phrases: summability factors, absolute matrix summability, Fourier series, infinite series, Hölder inequality, Minkowski inequality.

Communicated by Gradimir Milovanović.

The sequence-to-sequence transformation $t_n = \frac{1}{P_n}\sum_{v=0}^{n} p_v s_v$ defines the sequence (t_n) of the Riesz mean or simply the $(\bar{N}, p_n)$ mean of the sequence (s_n) generated by the sequence of coefficients (p_n) (see [**9**]).

The series $\sum a_n$ is said to be summable $|\bar{N}, p_n|_k$, $k \geqslant 1$, if (see [**1**])

$$\sum_{n=1}^{\infty} \Big(\frac{P_n}{p_n}\Big)^{k-1} |t_n - t_{n-1}|^k < \infty.$$

In the special case when $p_n = 1$ for all values of n (resp. $k = 1$), $|\bar{N}, p_n|_k$ summability is the same as $|C, 1|_k$ (resp. $|\bar{N}, p_n|$) summability.

2. Known Results

The following theorems are dealing with $|\bar{N}, p_n|_k$ summability factors of infinite series.

THEOREM 2.1. [**2**] *Let (p_n) be a sequence of positive numbers such that*

$$P_n = O(np_n) \quad as \quad n \to \infty. \tag{2.1}$$

Let (X_n) be a positive monotonic nondecreasing sequence. If the sequences (X_n), (λ_n) and (p_n) satisfy the conditions

$$\lambda_m X_m = O(1) \quad as\ m \to \infty, \tag{2.2}$$

$$\sum_{n=1}^{m} nX_n|\Delta^2\lambda_n| = O(1) \quad as\ m \to \infty, \tag{2.3}$$

$$\sum_{n=1}^{m} \frac{p_n}{P_n}|t_n|^k = O(X_m) \quad as\ m \to \infty, \tag{2.4}$$

then the series $\sum a_n\lambda_n$ is summable $|\bar{N}, p_n|_k$, $k \geqslant 1$.

THEOREM 2.2. [**4**] *Let (X_n) be a positive monotonic nondecreasing sequence. If the sequences (X_n), (λ_n), and (p_n) satisfy the conditions* (2.1)–(2.3) *and*

$$\sum_{n=1}^{m} \frac{p_n}{P_n}\frac{|t_n|^k}{X_n^{k-1}} = O(X_m) \qquad as \quad m \to \infty, \tag{2.5}$$

then the series $\sum a_n\lambda_n$ is summable $|\bar{N}, p_n|_k$, $k \geqslant 1$.

REMARK 2.1. It should be noted that condition (2.5) is reduced to the condition (2.4), when $k = 1$. When $k > 1$, condition (2.5) is weaker than condition (2.4) but the converse is not true (see [**4**] for details).

3. An application of absolute matrix summability to Fourier series

Let $A = (a_{nv})$ be a normal matrix, i.e., a lower triangular matrix of nonzero diagonal entries. Then A defines the sequence-to-sequence transformation, mapping

the sequence $s = (s_n)$ to $As = (A_n(s))$, where $A_n(s) = \sum_{v=0}^{n} a_{nv}s_v$, $n = 0, 1, \ldots$ The series $\sum a_n$ is said to be summable $|A|_k$, $k \geqslant 1$, if (see [**13**])

$$\sum_{n=1}^{\infty} n^{k-1}|\bar{\Delta}A_n(s)|^k < \infty,$$

and it is said to be summable $|A, p_n|_k$, $k \geqslant 1$, if (see [**12**])

$$\sum_{n=1}^{\infty} \Big(\frac{P_n}{p_n}\Big)^{k-1}|\bar{\Delta}A_n(s)|^k < \infty.$$

where $\bar{\Delta}A_n(s) = A_n(s) - A_{n-1}(s)$.

If we take $p_n = 1$ for all n, then $|A, p_n|_k$ summability is the same as $|A|_k$ summability. Also, if we take $a_{nv} = \frac{p_v}{P_n}$, then $|A, p_n|_k$ summability is the same as $|\bar{N}, p_n|_k$ summability. For any sequence (λ_n) we write $\Delta^2\lambda_n = \Delta\lambda_n - \Delta\lambda_{n+1}$ and $\Delta\lambda_n = \lambda_n - \lambda_{n+1}$. A sequence (λ_n) is said to be of bounded variation, denoted by $(\lambda_n) \in \mathcal{BV}$, if $\sum_{n=1}^{\infty} |\Delta\lambda_n| < \infty$. Let $f(t)$ be a periodic function with period 2π, and Lebesgue integrable over $(-\pi, \pi)$. Write

$$f(x) \sim \frac{1}{2}a_0 + \sum_{n=1}^{\infty}(a_n \cos nx + b_n \sin nx) = \sum_{n=0}^{\infty} C_n(x),$$

$\phi(t) = \frac{1}{2}[f(x+t) + f(x-t)]$, and $\phi_\alpha(t) = \frac{\alpha}{t^\alpha}\int_0^t (t-u)^{\alpha-1}\phi(u)\,du$ $(\alpha > 0)$.

It is well known that if $\phi(t) \in \mathcal{BV}(0, \pi)$, then $t_n(x) = O(1)$, where $t_n(x)$ is the $(C, 1)$ mean of the sequence $(nC_n(x))$ (see [**7**]).

Many works have been done dealing with absolute summability factors of Fourier series (see [**3**–**5**, **11**]). Among them, in [**4**], Bor has proved the following theorem dealing with the Fourier series.

THEOREM 3.1. *If $\phi_1(t) \in \mathcal{BV}(0, \pi)$, (X_n) is a positive monotonic nondecreasing sequence, the sequences (p_n), (λ_n) satisfy conditions* (2.1)–(2.3) *and*

$$\sum_{n=1}^{m} \frac{p_n}{P_n}\frac{|t_n(x)|^k}{X_n^{k-1}} = O(X_m) \qquad as \quad m \to \infty,$$

then the series $\sum C_n(x)\lambda_n$ is summable $|\bar{N}, p_n|_k$, $k \geqslant 1$.

If we take $p_n = 1$ for all values of n, then we obtain a new result dealing with $|C, 1|_k$ summability factors of Fourier series.

4. Main Results

We generalize Theorem 3.1 for $|A, p_n|_k$ summability factors of Fourier series. Before stating the main theorem, we must first introduce some further notations.

With a normal matrix $A = (a_{nv})$, we associate two lower semimatrices $\bar{A} = (\bar{a}_{nv})$ and $\hat{A} = (\hat{a}_{nv})$ where $\bar{a}_{nv} = \sum_{i=v}^{n} a_{ni}$, $n, v = 0, 1, \ldots$ and $\hat{a}_{00} = \bar{a}_{00} = a_{00}$,

$\hat{a}_{nv} = \bar{a}_{nv} - \bar{a}_{n-1,v}$, $n = 1, 2, \ldots$ We note that $\bar{A}$ and $\hat{A}$ are the well-known matrices of series-to-sequence and series-to-series transformations, respectively. So, we have

$$A_n(s) = \sum_{v=0}^{n} a_{nv} s_v = \sum_{v=0}^{n} \bar{a}_{nv} a_v \quad \text{and} \quad \bar{\Delta} A_n(s) = \sum_{v=0}^{n} \hat{a}_{nv} a_v. \tag{4.1}$$

THEOREM 4.1. *Let $k \geqslant 1$ and $A = (a_{nv})$ be a positive normal matrix such that*

$$\bar{a}_{n0} = 1, \ n = 0, 1, \ldots, \ \ a_{n-1,v} \geqslant a_{nv}, \ \textit{for } n \geqslant v+1,$$
$$a_{nn} = O\big(p_n/P_n\big), \ \ \hat{a}_{n,v+1} = O(v|\Delta_v(\hat{a}_{nv}|).$$

If all the conditions of Theorem 3.1 *are satisfied, then the series $\sum C_n(x)\lambda_n$ is summable $|A, p_n|_k$, $k \geqslant 1$.*

If we take $a_{nv} = \frac{p_v}{P_n}$, then we get Theorem 3.1. We need the following lemma for the proof of our theorem.

LEMMA 4.1. [**2**] *Under the conditions of Theorem* 2.2 *we have*

$$nX_n|\Delta\lambda_n| = O(1) \ \textit{as } n \to \infty, \quad \textit{and} \quad \sum_{n=1}^{\infty} X_n|\Delta\lambda_n| < \infty.$$

5. Proof of Theorem 4.1

Let $(I_n(x))$ denote the A-transform of the series $\sum_{n=1}^{\infty} C_n(x)\lambda_n$. Then, by (4.1), we have $\bar{\Delta} I_n(x) = \sum_{v=1}^{n} \hat{a}_{nv} C_v(x)\lambda_v$. Applying Abel's transformation to this sum, we get

$$\begin{aligned}
\bar{\Delta} I_n(x) &= \sum_{v=1}^{n} \hat{a}_{nv} C_v(x)\lambda_v \frac{v}{v} = \sum_{v=1}^{n-1} \Delta_v\Big(\frac{\hat{a}_{nv}\lambda_v}{v}\Big) \sum_{r=1}^{v} rC_r(x) + \frac{\hat{a}_{nn}\lambda_n}{n} \sum_{r=1}^{n} rC_r(x) \\
&= \sum_{v=1}^{n-1} \Delta_v\Big(\frac{\hat{a}_{nv}\lambda_v}{v}\Big)(v+1)t_v(x) + \hat{a}_{nn}\lambda_n \frac{n+1}{n} t_n(x) \\
&= \sum_{v=1}^{n-1} \Delta_v(\hat{a}_{nv})\lambda_v t_v(x)\frac{v+1}{v} + \sum_{v=1}^{n-1} \hat{a}_{n,v+1}\Delta\lambda_v t_v(x)\frac{v+1}{v} \\
&\quad + \sum_{v=1}^{n-1} \hat{a}_{n,v+1}\lambda_{v+1}\frac{t_v(x)}{v} + a_{nn}\lambda_n t_n(x)\frac{n+1}{n} \\
&= I_{n,1}(x) + I_{n,2}(x) + I_{n,3}(x) + I_{n,4}(x).
\end{aligned}$$

To complete the proof of Theorem 4.1, by Minkowski's inequality, it is sufficient to show that

$$\sum_{n=1}^{\infty} \Big(\frac{P_n}{p_n}\Big)^{k-1} |I_{n,r}(x)|^k < \infty, \quad \text{for} \quad r = 1, 2, 3, 4.$$

First, by applying Hölder's inequality with indices k and k', where $k > 1$ and $\frac{1}{k} + \frac{1}{k'} = 1$, we have

$$\begin{aligned}
\sum_{n=2}^{m+1}\Big(\frac{P_n}{p_n}\Big)^{k-1}|I_{n,1}(x)|^k &\leqslant \sum_{n=2}^{m+1}\Big(\frac{P_n}{p_n}\Big)^{k-1}\Big\{\sum_{v=1}^{n-1}|\frac{v+1}{v}||\Delta_v(\hat{a}_{nv})||\lambda_v||t_v(x)|\Big\}^k\\
&= O(1)\sum_{n=2}^{m+1}\Big(\frac{P_n}{p_n}\Big)^{k-1}\sum_{v=1}^{n-1}|\Delta_v(\hat{a}_{nv})||\lambda_v|^k|t_v(x)|^k\\
&\qquad\times\Big\{\sum_{v=1}^{n-1}|\Delta_v(\hat{a}_{nv})|\Big\}^{k-1}\\
&= O(1)\sum_{n=2}^{m+1}\Big(\frac{P_n}{p_n}\Big)^{k-1}a_{nn}^{k-1}\Big\{\sum_{v=1}^{n-1}|\Delta_v(\hat{a}_{nv})||\lambda_v|^k|t_v(x)|^k\Big\}\\
&= O(1)\sum_{v=1}^{m}|\lambda_v|^{k-1}|\lambda_v||t_v(x)|^k\sum_{n=v+1}^{m+1}|\Delta_v(\hat{a}_{nv})|\\
&= O(1)\sum_{v=1}^{m}\frac{1}{X_v^{k-1}}|\lambda_v||t_v(x)|^k\frac{p_v}{P_v}\\
&= O(1)\sum_{v=1}^{m-1}\Delta|\lambda_v|\sum_{r=1}^{v}\frac{p_r}{P_r}\frac{|t_r(x)|^k}{X_r^{k-1}}+O(1)|\lambda_m|\sum_{v=1}^{m}\frac{p_v}{P_v}\frac{|t_v(x)|^k}{X_v^{k-1}}\\
&= O(1)\sum_{v=1}^{m-1}|\Delta\lambda_v|X_v+O(1)|\lambda_m|X_m = O(1)\quad\text{as}\ \ m\to\infty,
\end{aligned}$$

by virtue of the hypotheses of Theorem 4.1 and Lemma 4.1. Now, using Hölder's inequality we have

$$\begin{aligned}
\sum_{n=2}^{m+1}\Big(\frac{P_n}{p_n}\Big)^{k-1}|I_{n,2}(x)|^k &\leqslant \sum_{n=2}^{m+1}\Big(\frac{P_n}{p_n}\Big)^{k-1}\Big\{\sum_{v=1}^{n-1}|\frac{v+1}{v}||\hat{a}_{n,v+1}||\Delta\lambda_v||t_v(x)|\Big\}^k\\
&= O(1)\sum_{n=2}^{m+1}\Big(\frac{P_n}{p_n}\Big)^{k-1}\Big\{\sum_{v=1}^{n-1}|\hat{a}_{n,v+1}||\Delta\lambda_v||t_v(x)|\Big\}^k\\
&= O(1)\sum_{n=2}^{m+1}\Big(\frac{P_n}{p_n}\Big)^{k-1}\sum_{v=1}^{n-1}(v|\Delta\lambda_v|)^k|\Delta_v(\hat{a}_{nv})||t_v(x)|^k\\
&\qquad\times\Big\{\sum_{v=1}^{n-1}|\Delta_v(\hat{a}_{nv})|\Big\}^{k-1}\\
&= O(1)\sum_{n=2}^{m+1}\Big(\frac{P_n}{p_n}\Big)^{k-1}a_{nn}^{k-1}\sum_{v=1}^{n-1}(v|\Delta\lambda_v|)^k|\Delta_v(\hat{a}_{nv})||t_v(x)|^k\\
&= O(1)\sum_{v=1}^{m}(v|\Delta\lambda_v|)^{k-1}(v|\Delta\lambda_v|)|t_v(x)|^k\sum_{n=v+1}^{m+1}|\Delta_v(\hat{a}_{nv})|
\end{aligned}$$

$$
\begin{aligned}
&= O(1)\sum_{v=1}^{m}\frac{p_v}{P_v}\frac{1}{X_v^{k-1}}|t_v(x)|^k(v|\Delta\lambda_v|)\\
&= O(1)\sum_{v=1}^{m-1}\Delta(v|\Delta\lambda_v|)\sum_{r=1}^{v}\frac{p_r}{P_r}\frac{1}{X_r^{k-1}}|t_r(x)|^k\\
&\quad + O(1)m|\Delta\lambda_m|\sum_{v=1}^{m}\frac{p_v}{P_v}\frac{1}{X_v^{k-1}}|t_v(x)|^k\\
&= O(1)\sum_{v=1}^{m-1}|\Delta(v|\Delta\lambda_v|)|X_v + O(1)m|\Delta\lambda_m|X_m\\
&= O(1)\sum_{v=1}^{m-1}vX_v|\Delta^2\lambda_v| + O(1)\sum_{v=1}^{m-1}X_v|\Delta\lambda_v| + O(1)m|\Delta\lambda_m|X_m = O(1)
\end{aligned}
$$

as $m \to \infty$, by virtue of the hypotheses of Theorem 4.1 and Lemma 4.1. Again, we have that

$$
\begin{aligned}
\sum_{n=2}^{m+1}\Big(\frac{P_n}{p_n}\Big)^{k-1}|I_{n,3}(x)|^k &= \sum_{n=2}^{m+1}\Big(\frac{P_n}{p_n}\Big)^{k-1}\Big|\sum_{v=1}^{n-1}\hat{a}_{n,v+1}\lambda_{v+1}\frac{t_v(x)}{v}\Big|^k\\
&\leqslant \sum_{n=2}^{m+1}\Big(\frac{P_n}{p_n}\Big)^{k-1}\Big\{\sum_{v=1}^{n-1}|\hat{a}_{n,v+1}||\lambda_{v+1}|\frac{|t_v(x)|}{v}\Big\}^k\\
&= O(1)\sum_{n=2}^{m+1}\Big(\frac{P_n}{p_n}\Big)^{k-1}\Big\{\sum_{v=1}^{n-1}|\Delta_v(\hat{a}_{nv})||\lambda_{v+1}||t_v(x)|\Big\}^k\\
&= O(1)\sum_{n=2}^{m+1}\Big(\frac{P_n}{p_n}\Big)^{k-1}\sum_{v=1}^{n-1}|\Delta_v(\hat{a}_{nv})||\lambda_{v+1}|^k|t_v(x)|^k\\
&\qquad\qquad \times\Big\{\sum_{v=1}^{n-1}|\Delta_v(\hat{a}_{nv})|\Big\}^{k-1}\\
&= O(1)\sum_{n=2}^{m+1}\Big(\frac{P_n}{p_n}\Big)^{k-1}a_{nn}^{k-1}\sum_{v=1}^{n-1}|\Delta_v(\hat{a}_{nv})||\lambda_{v+1}|^k|t_v(x)|^k\\
&= O(1)\sum_{v=1}^{m}|\lambda_{v+1}|^k|t_v(x)|^k\sum_{n=v+1}^{m+1}|\Delta_v(\hat{a}_{nv})|\\
&= O(1)\sum_{v=1}^{m}\frac{p_v}{P_v}|t_v(x)|^k|\lambda_{v+1}|^{k-1}|\lambda_{v+1}|\\
&= O(1)\sum_{v=1}^{m}\frac{1}{X_v^{k-1}}|\lambda_{v+1}||t_v(x)|^k\frac{p_v}{P_v} = O(1) \quad \text{as } m \to \infty,
\end{aligned}
$$

by virtue of the hypotheses of Theorem 4.1 and Lemma 4.1. Finally, as in $T_{n,1}$, we have that

$$\begin{aligned}\sum_{n=1}^{m}\Big(\frac{P_n}{p_n}\Big)^{k-1}|I_{n,4}(x)|^k &= O(1)\sum_{n=1}^{m}\Big(\frac{P_n}{p_n}\Big)^{k-1}a_{nn}^k|\lambda_n|^k|t_n(x)|^k\\ &= O(1)\sum_{n=1}^{m}\frac{p_n}{P_n}|\lambda_n|^{k-1}|\lambda_n||t_n(x)|^k\\ &= O(1)\sum_{n=1}^{m}\frac{1}{X_n^{k-1}}|\lambda_n||t_n(x)|^k\frac{p_n}{P_n} = O(1) \quad \text{as } m\to\infty,\end{aligned}$$

by virtue of hypotheses of the Theorem 4.1 and Lemma 4.1. This completes the proof of Theorem 4.1.

If we take $a_{nv} = \frac{p_v}{P_n}$ in Theorem 4.1, then we get Theorem 3.1 and if we take $p_n = 1$ for all values of n in Theorem 4.1, then we get a new result dealing with the $|A|_k$ summability method. Also, if we take $a_{nv} = \frac{p_v}{P_n}$ and $p_n = 1$ for all values of n in Theorem 4.1, then we get a result concerning the $|C,1|_k$ summability methods.

References

1. H. Bor, *On two summability methods*, Math. Proc. Camb. Philos. Soc. **97** (1985), 147–149.
2. ———, *On absolute summability factors*, Proc. Am. Math. Soc. **118** (1993), 71–75.
3. ———, *Some new results on infinite series and Fourier series*, Positivity **19** (2015), 467–473.
4. ———, *On absolute weighted mean summability of infinite series and Fourier series*, Filomat **30**(10) (2016), 2803–2807.
5. H. Bor, D. S. Yu, P. Zhou, *On local property of absolute summability of factored Fourier series*, Filomat **28**(8) (2014), 1675–1686.
6. E. Cesàro, *Sur la multiplication des sèries*, Bull. Sci. Math. **14** (1890), 114–120.
7. K. K. Chen, *Functions of bounded variation and the Cesaro means of Fourier series*, Acad. Sinica Sci. Record **1** (1945), 283–289.
8. T. M. Flett, *On an extension of absolute summability and some theorems of Littlewood and Paley*, Proc. Lond. Math. Soc. **7** (1957), 113–141.
9. G. H. Hardy, *Divergent Series*, Oxford Univ. Press, Oxford, (1949).
10. E. Kogbetliantz, *Sur lès series absolument sommables par la methode des moyennes arithmetiques*, Bull. Sci. Math. **49** (1925), 234–256.
11. H. S. Özarslan, Ş. Yıldız, *A new study on the absolute summability factors of Fourier series*, J. Math. Anal. **7** (2016), 31–36.
12. W. T. Sulaiman, *Inclusion theorems for absolute matrix summability methods of an infinite series.* IV, Indian J. Pure Appl. Math. **34**(11) (2003), 1547–1557.
13. N. Tanović-Miller, *On strong summability*, Glas. Mat., III. Ser. **14** (34) (1979), 87–97.

Department of Mathematics
Ahi Evran University
Kırşehir
Turkey
sebnemyildiz@ahievran.edu.tr,
sebnem.yildiz82@gmail.com

(Received 16 07 2016)