PUBLICATIONS DE L'INSTITUT MATHÉMATIQUE
Nouvelle série, tome 103(117) (2018), 191–198 DOI: https://doi.org/10.2298/PIM1817191P

RELATIONS BETWEEN KERNELS AND IMAGES OF REDUCED POWERS FOR SOME RIGHT $\mathcal{A}_p$-MODULES

Theodore Popelensky

Abstract. We investigate the right action of the mod p Steenrod algebra $\mathcal{A}_p$ on the homology $H_*(L^{\wedge s}, \mathbb{Z}_p)$ where $L = B\mathbb{Z}_p$ is the lens space. Following ideas of Ault and Singer we investigate the relation between intersection of kernels of the reduced powers P^{p^i} and Bockstein element β and the intersection of images of $P^{p^{i+1}-1}$ and of β. Namely one can check that $\bigcap_{i=0}^{k} \operatorname{im} P^{p^{i+1}-1} \subset \bigcap_{i=0}^{k} \ker P^{p^i}$ and $\bigcap_{i=0}^{k} \operatorname{im} P^{p^{i+1}-1} \cap \operatorname{im} \beta \subset \bigcap_{i=0}^{k} \ker P^{p^i} \cap \ker \beta$. We generalize Ault's homotopy systems to $p > 2$ and examine when the reverse inclusions are true.

Introduction

Two natural questions arise when the action of the Steenrod algebra $\mathcal{A}_p$ on the cohomology $H^*(X, \mathbb{Z}_p)$ of a space X is considered. One is to describe the *annihilating ideal* $I_X \subset \mathcal{A}_p$ which is defined as $I_X = \{\phi \in \mathcal{A}_p \mid \phi(H^*(X, \mathbb{Z}_p)) = 0\}$. The problem appeared to be difficult even for $p = 2$ and $X = \mathbb{R}P^\infty$, see [**1**].

The second question is to describe a minimal generating set of the $\mathcal{A}_p$-module $H^*(X, \mathbb{Z}_p)$ and the set of elements of the form ϕx, where $\phi \in \mathcal{A}_p$, $\deg \phi > 0$, $x \in H^*(X, \mathbb{Z}_p)$ (or the space spanned by such elements, this is the so called hit problem).

This question is far from being solved even for $X = \prod_{i=1}^{k} \mathbb{R}P^\infty$, though cases $k \leqslant 4$ are completely described in the papers [**4, 5, 6, 7**]

In this note we address the problem which is dual to the second question. Consider the conjugate action of the Steenrod algebra $\mathcal{A}_p$ on the $\mathbb{Z}_p$-homology of a space X. We say that $x \in H_*(X, \mathbb{Z}_p)$ is *P-annihilated* iff $xP^m = 0$ for all $m > 0$ and *βP-annihilated* iff x is P-annihilated and $x\beta = 0$. The problem is to describe the subspace of all P-anihilated or βP-annihilated elements. Following ideas of Ault and Singer [**2, 3**], we study subspaces of *partially* annihilated elements ($\Delta_M(k)$ and $\Delta_M^\beta(k)$, see below). It appears that there is a structure (homotopy system, see Definitions 2.1 and 2.2) which relates these spaces with the spaces of so called

2010 *Mathematics Subject Classification:* 55S10; 55R40; 57T25.
Key words and phrases: Steenrod algebra, reduced powers.
This work was funded by Russian Science Foundation 16-11-10069.

spike images ($I_M(k)$ and $I_M^\beta(k)$, see below). The main results are Theorem 2.1 and Corollary 2.1. Also we show that such homotopy systems exist for $\tilde{\Lambda}$ (see Section 3 and Theorem 4.1). Here $\tilde{\Lambda}$ is a bigraded space which is given by (1.2), see below.

Throughout the paper all the (co)homologies are considered with $\mathbb{Z}_p$ coefficients.

The author is grateful to the referees for careful reading of the manuscript and for finding inconsistencies in the preliminary version of it.

1. Lens space and the $\mathcal{A}_p$-module $\tilde{\Lambda}$.

Denote by L the classifying space $B\mathbb{Z}_p$. The cohomology ring structure is described by the isomorphism $H^*(L) \equiv \mathbb{Z}_p[x] \otimes \Lambda_{\mathbb{Z}_p}(y)$, where $\deg x = 2$ and $\deg y = 1$. The action of the Steenrod algebra $\mathcal{A}_p$ on $H^*(L)$ is given by the following relations:

$$P^k x^m = \binom{m}{k} x^{k(p-1)+m}, \qquad \beta x^m = 0,$$
$$P^k x^m y = \binom{m}{k} x^{k(p-1)+m} y, \quad \beta x^m y = x^{m+1}.$$

There is the conjugate right action of the algebra $\mathcal{A}_p$ on the reduced homology of L. Denote by $[a]$, $a \geqslant 1$, the generator of $\tilde{H}_{2a}(L)$ and by $[\hat{a}]$, $a \geqslant 0$, the generator of $\tilde{H}_{2a+1}(L)$. The action of $\mathcal{A}_p$ on these elements is described by the formulas:

$$\begin{aligned} P^k &= \binom{a-(p-1)k}{k}[a-(p-1)k], \qquad [a]\beta = [\widehat{a-1}], \\ [\hat{a}]P^k &= \binom{a-(p-1)k}{k}[a-\widehat{(p-1)}k], \qquad [\hat{a}]\beta = 0. \end{aligned} \tag{1.1}$$

Here we write a homomorphism to the right from its argument.

Consider the bigraded algebra $\tilde{\Lambda} = \{\tilde{\Lambda}_{*,*}\}$ which is defined by

$$\tilde{\Lambda}_{n,k} = \begin{cases} \mathbb{Z}_p, & \text{if } n = k = 0; \\ \tilde{H}_k(L^{\wedge n}, \mathbb{Z}_p), & \text{if } n \geqslant 1; \\ 0, & \text{othervise.} \end{cases} \tag{1.2}$$

This algebra is a very natural module on which the right action of $\mathcal{A}_p$ should be studied.

2. Homotopy system

For a right $\mathcal{A}_p$-module M define the following subspaces:

$$\Delta_M(k) = \bigcap_{i=0}^{k} \ker P^{p^i}, \qquad I_M(k) = \bigcap_{i=0}^{k} \operatorname{im} P^{p^{i+1}-1},$$
$$\Delta_M^\beta(k) = \Delta_M(k) \cap \ker\beta, \quad I_M^\beta(k) = I_M(k) \cap \operatorname{im}\beta.$$

The subspaces $\Delta_M(k)$ and $\Delta_M^\beta(k)$ are subspaces of partially annihilated elements. The subspaces $I_M(k)$ and $I_M^\beta(k)$ are subspaces of simultaneous spike images. For $M = \tilde{\Lambda}$, denote these modules by $\Delta(k)$, $\Delta^\beta(k)$, $I(k)$, $I^\beta(k)$.

LEMMA 2.1. *For an $\mathcal{A}_p$-module M and $k \geqslant 0$ one has $I_M(k) \subset \Delta_M(k)$ and $I_M^\beta(k) \subset \Delta_M^\beta(k)$.*

REMARK 2.1. *Generally $I_M(k) \not\subset \Delta_M^\beta(k)$.*

PROOF. The statements easily follow from the relations $P^{p^{m+1}-1}P^{p^m} = 0$ and $\beta\beta = 0$. The last one is a particular case of the general relation $P^{pm-1}P^m = 0$. To check it, note that the Adem relation applies to $P^{pm-1}P^m$:

$$P^{pm-1}P^m = \sum_{j=0}^{m-1} (-1)^{pm-1+j} \binom{(p-1)(m-j)-1}{pm-1-pj} P^{(p+1)m-1-j}P^j.$$

The inequality $(p-1)(m-j)-1 < pm-1-pj$ follows easily from the restriction $j \leqslant m-1$. Hence all the binomial coefficients in the last sum vanish. □

In some cases one can prove the reverse inclusions $\Delta_M(k) \subset I_M(k)$ and $\Delta_M^\beta(k) \subset I_M^\beta(k)$. Now we describe an algebraic structure which is responsible for these inclusions.

DEFINITION 2.1. For a right $\mathcal{A}_p$-module M define a *k-th order homotopy system* to be a *null space* $N \subset M$ and a collection of $\mathbb{Z}_p$-homomorphisms $Q_m : M \to M$ for $0 \leqslant m \leqslant k$ such that

(i) $Q_m(N) \subset N$ for $0 \leqslant m \leqslant k$;

(ii) for $1 \leqslant m \leqslant k$ and $l < p^m$ the following diagram comutes

$$\begin{array}{ccc} N & \xrightarrow{P^l} & M \\ {\scriptstyle Q_m}\downarrow & & \downarrow{\scriptstyle Q_m} \\ N & \xrightarrow{P^l} & M \end{array}$$

(iii) $(x)(P^{p^m}Q_m - Q_m P^{p^m}) = x$ for any $x \in N$.

DEFINITION 2.2. For a right $\mathcal{A}_p$-module M define a *k-th order β-homotopy system* to be a k-th order homotopy system $\{Q_m\}$ with null space $N \subset M$, which is equipped with a $\mathbb{Z}_p$-homomorphism $\alpha : M \to M$ such that

(iv) $\alpha(N) \subset N$ and for any $x \in N$ one has $x\alpha\beta + x\beta\alpha = x$.

For a nonnegative integer s, denote by $0 \leqslant d_j(s) < p$ the coefficients of the expansion $s = \sum d_j(s)p^j$ in the base p. Recall Lucas's theorem which states that for any nonnegative integers x and y one has

$$\binom{x}{y} \equiv \prod_j \binom{d_j(x)}{d_j(x)} \mod p.$$

LEMMA 2.2. *For $1 \leqslant a \leqslant p-1$ one has*

$$(P^{p^0})^a (P^{p^1})^a \dots (P^{p^{n-1}})^a (P^{p^n})^a = cP^{a(p^0+p^1+\dots+p^{n-1}+p^n)},$$

for some nonzero $c \in \mathbb{Z}_p$.

PROOF. First of all we show that for any a such that $1 \leqslant a \leqslant p-1$ one has $(P^1)^a = cP^a$ for some nonzero $c \in \mathbb{Z}_p$. For $a = 1$ the statement is trivial. Assume the statement is true for $a = b$, where $b < p-1$. Then $(P^1)^{b+1} = (P^1)^b P^1 = cP^b P^1$, where $c \neq 0$. Apply the Adem relation to $P^b P^1$:

$$P^b P^1 = (-1)^b \binom{p-2}{b} P^{b+1},$$

hence $(P^1)^{b+1} = c(-1)^b \binom{p-2}{b} P^{b+1}$. One easily checks that the binomial coefficient $\binom{p-2}{b}$ is nonzero in $\mathbb{Z}_p$.

Now we proceed by induction to prove that for any integers a and b such that $1 \leqslant a \leqslant p-1$ and $0 \leqslant b < a$ one has

$$P^{a(p^0+p^1+\cdots+p^m)+bp^{m+1}} P^{p^{m+1}} = dP^{a(p^0+p^1+\cdots+p^m)+(b+1)p^{m+1}}$$

for some nonzero $d \in \mathbb{Z}_p$.

Denote $a(p^0 + p^1 + \cdots + p^m) + bp^{m+1}$ by N. Apply the Adem relation to the product $P^N P^{p^{m+1}}$:

$$P^N P^{p^{m+1}} = \sum_{t=0}^{[N/p]} (-1)^{N+t} \binom{(p-1)(p^{m+1}-t)-1}{N-pt} P^{N+p^{m+1}-t} P^t.$$

The coefficient for $t = 0$, by Lucas's theorem, is equal to $(-1)^N \binom{p-2}{b} \binom{p-1}{a}^{m+1}$ and hence is not zero.

It is left to prove that for $t > 0$, the binomial coefficient vanishes. To apply Lucas's theorem, it is enough to find j such that $d_j((p-1)(p^{m+1}-t)-1) < d_j(N-pt)$. Fix $0 < t < [N/p]$. For simplicity let $a_j = d_j(t)$. Choose the smallest j such that $a_j \neq 0$.

Case 1. If $1 \leqslant a_j \leqslant a$ then $d_j(N-pt) = a$ while $d_j((p-1)(p^{m+1}-t)-1) = a_j - 1$. But $a_j - 1 < a$.

Case 2. If $a_{j+1} \geqslant a_j > a$ then $d_{j+1}(N-pt) = p + a - a_j$, while $d_{j+1}((p-1)(p^{m+1}-t)-1) = a_{j+1} - a_j$. Note that $a_{j+1} - a_j < p + a - a_j$.

Case 3. Assume for some k one has $a_{k+1} > a_k \geqslant a$ and $a_k \leqslant a_{k-1} \leqslant \cdots \leqslant a_{j+2} \leqslant a_{j+1} < a_j$. Then $d_{k+1}(N-pt) = p - 1 + a - a_k$ and $d_{k+1}((p-1)(p^{m+1}-t)-1) = -1 + a_{k+1} - a_k$. Note that $p - 1 + a - a_k > -1 + a_{k+1} - a_k$.

Case 4. Assume for some k one has $a_{k+1} < a \leqslant a_k \leqslant a_{k-1} \leqslant \cdots \leqslant a_{j+2} \leqslant a_{j+1} < a_j$. Then $d_{k+1}(N-pt) = p - 1 + a - a_k$ and $d_{k+1}((p-1)(p^{m+1}-t)-1) = p - 1 + a_{k+1} - a_k$. Note that $p - 1 + a - a_k > p - 1 + a_{k+1} - a_k$. □

THEOREM 2.1. *Suppose that for a right $\mathcal{A}_p$-module M there exists a k-th order homotopy system $\{Q_m\}$ with a null space N. Then $\Delta_M(k) \cap N = I_M(k) \cap N$. Moreover, if $x \in \Delta_M(k) \cap N$ then $y = xQ_m^{p-1} Q_{m-1}^{p-1} \dots Q_0^{p-1}$ satisfies $yP^{p^{m+1}-1} = cx$ for some nonzero $c \in \mathbb{Z}_p$.*

PROOF. Denote for simplicity P^{p^m} by P_m. Using standard technique for calculations with commutators involving creation-annihilation operators we prove for $j = 0, \dots, m-1$ that

$$xQ_m^{p-1} Q_{m-1}^{p-1} \dots Q_j^{p-1} P_j^{p-1} = -xQ_m^{p-1} Q_{m-1}^{p-1} \dots Q_{j+1}^{p-1}.$$

The relation

$$xQ_m^{p-1}P_m^{p-1} = -x$$

is proved in the same way. Denote $xQ_m^{p-1}Q_{m-1}^{p-1}\dots Q_{j+1}^{p-1}$ by W. Since all Q_n's left the subspace N invariant and P_j commutes with $Q_m, \dots, Q_{j+1}$ on the subspace N one has $WP_j = 0$. Note that

$$[Q_j^n, P_j^n] = \sum_{s+t=n-1} P_j^s [Q_j^n, P_j] P_j^t.$$

Apply this relation, for $n = p-1$, to the product $W(Q_j)^{p-1}(P_j)^{p-1}$:

$$WQ_j^{p-1}P_j^{p-1} = WP_j^{p-1}Q_j^{p-1} + \sum_{s+t=p-2} WP_j^s [Q_j^{p-1}, P_j] P_j^t.$$

On the right-hand side, all the summands vanish except $W[Q_j^{p-1}, P_j]P_j^{p-2}$. Now use the equality

$$[Q_j^{p-1}, P_j] = \sum_{s+t=p-2} Q_j^s [Q_j, P_j] Q_j^t$$

and rewrite the product $W[Q_j^{p-1}, P_j]P_j^{p-2}$ as

$$W[Q_j^{p-1}, P_j]P_j^{p-2} = \sum_{s+t=p-2} WQ_j^s [Q_j, P_j] Q_j^t P_j^{p-2}.$$

Since $WQ_j^s \in N$, one has $WQ_j^s[Q_j, P_j] = -WQ_j^s$. Finally,

$$W[Q_j^{p-1}, P_j]P_j^{p-2} = \sum_{s+t=p-2} -WQ_j^s Q_j^t P_j^{p-2} = -(p-1)WQ_j^{p-2}P_j^{p-2}.$$

Proceeding in the same way, we obtain the equality

$$WQ_j^{p-1}P_j^{p-1} = (-1)^{p-1}(p-1)!W = -W.$$

Here we use simple part of the Wilson theorem which states that for a prime p one has $(p-1)! + 1 \equiv 0 \mod p$.

By Lemma 2.2, $P^{p^{m+1}-1} = d\, P_0^{p-1} \dots P_m^{p-1}$ for some nonzero $d \in \mathbb{Z}_p$. Then

$$\begin{aligned} yP^{p^{m+1}-1} &= d\, xQ_m^{p-1} \dots Q_1^{p-1}Q_0^{p-1}P_0^{p-1}P_1^{p-1} \dots P_m^{p-1} \\ &= -d\, xQ_m^{p-1} \dots Q_1^{p-1}P_1^{p-1} \dots P_m^{p-1} \\ &= \dots = (-1)^m d\, xQ_m^{p-1}P_m^{p-1} = (-1)^{m+1} d\, x. \end{aligned}$$ □

Corollary 2.1. *Suppose that for a right $\mathcal{A}_p$-module M there exists a k-th order β-homotopy system $\{Q_m\}$ with a null space N and a $\mathbb{Z}_p$-homomorphism α. Then $\Delta_M^\beta(k) \cap N = I_M^\beta(k) \cap N$. Moreover, if $x \in \Delta_M^\beta(k) \cap N$, then $y = x\alpha$ belongs to N and satisfies $y\beta = x$.*

Proof. From $x\beta\alpha + x\alpha\beta = x$ and $x \in \ker\beta$ it follows that $y\beta = x\alpha\beta = x$. From Definition 2.2 one has $y \in N$. □

3. Shift maps in the homology of the smash-product power of the lens space

In this section we consider examples of homotopy systems for certain $\mathcal{A}_p$-modules. The reduced homology of $L^{\wedge s}$ is a vector space with the basis which consists of the elements $[u_1, \dots, u_s]$ of degree $\sum \deg u_j$, where every u_j is a, $a \geqslant 1$, or $\hat{a}$, $a \geqslant 0$, and $\deg a = 2a$, $\deg \hat{a} = 2a+1$. Now we describe homotopy systems which appear naturally for $\tilde{H}_*(L^{\wedge s})$.

Define the shift map $\Psi_i^l : \tilde{\Lambda}_{s,*} \to \tilde{\Lambda}_{s,*+l}$ by formulas $[u_1, \dots, u_i, \dots, u_s]\Psi_i^l = [u_1, \dots, u_i+l, \dots, u_s]$. Here u_j+l denotes a_j+l if $u_j = a_j$ and $\widehat{a_j+l}$ if $u_j = \hat{a}_j$. Also define $[u_1, \dots, a_i, \dots, u_s]\alpha_i = 0$ and $[u_1, \dots, \hat{a}_i, \dots, u_s]\alpha_i = [u_1, \dots, a_i+1, \dots, u_s]$. Clearly $(\Psi_i^{p^m})^k = \Psi_i^{kp^m}$.

Hereinafter for $s = 1$ we denote $\Psi^l = \Psi_1^l$ and $\alpha = \alpha_1$.

Lemma 3.1. *We have*
(a) $[a]\Psi^{kp^m}P^n = [a]P^n\Psi^{kp^m}$ *for* $a \geqslant (p-1)n$, $p^m > n$, $1 \leqslant k \leqslant p-1$;
(b) $[\hat{a}]\Psi^{kp^m}P^n = [\hat{a}]P^n\Psi^{kp^m}$ *for* $a \geqslant (p-1)n$, $p^m > n$, $1 \leqslant k \leqslant p-1$;
(c) $[a]\Psi^{kp^m}\beta = [a]\beta\Psi^{kp^m} \neq 0$;
(d) $[\hat{a}]\Psi^{kp^m}\beta = 0 = [\hat{a}]\beta\Psi^{kp^m}$.

Proof. By formulas (1.1), one has

$$[a]\Psi^{kp^m}P^n = \binom{a+kp^m-(p-1)n}{n}[a+kp^m-(p-1)n],$$
$$[a]P^n\Psi^{kp^m} = \binom{a-(p-1)n}{n}[a+kp^m-(p-1)n].$$

Two binomial coefficients coincide under the assumptions of Lemma. Other statements are proved in the same way. □

Lemma 3.2. *We have*
(a) $[a]P^{p^m}\Psi^{(p-1)p^m} - [a]\Psi^{(p-1)p^m}P^{p^m} = [a]$ *for* $m \geqslant 0$ *and* $a \geqslant (p-1)p^m$;
(b) $[\hat{a}]P^{p^m}\Psi^{(p-1)p^m} - [\hat{a}]\Psi^{(p-1)p^m}P^{p^m} = [\hat{a}]$ *for* $m \geqslant 0$ *and* $a \geqslant (p-1)p^m$;
(c) $[a]\alpha\beta + [a]\beta\alpha = [a]$ *for* $a \geqslant 1$;
(d) $[\hat{a}]\alpha\beta + [\hat{a}]\beta\alpha = [\hat{a}]$ *for* $a \geqslant 0$.

Proof. By straightforward computation, one can get

$$[a]P^{p^m}\Psi^{(p-1)p^m} = \binom{a-(p-1)p^m}{p^m}[a], \quad [a]\Psi^{(p-1)p^m}P^{p^m} = \binom{a}{p^m}[a].$$

Assume $a = \sum b_k p^k$, where $0 \leqslant b_k < p$. If $b_m = p-1$ then $\binom{a}{p^m} = p-1$ and $\binom{a-(p-1)p^m}{p^m} = 0$. If $b_m < p-1$ then $\binom{a}{p^m} = b_m$ and $\binom{a-(p-1)p^m}{p^m} = b_m+1$. The proof of (b) is similar, and the statements (b) and (c) are clear. □

Lemma 3.3. *Assume* $x = [u_1, \dots, u_s]$, *where each* u_j *is* a_j *or* $\hat{a}_j$. *If* $p^m > n$, $a_i \geqslant (p-1)n$, $1 \leqslant k \leqslant p-1$, *then one has* $x\Psi_i^{kp^m}P^n = xP^n\Psi_i^{kp^m}$.

PROOF. Without loss of generality one can assume $i = 1$. The case $s = 1$ was considered in Lemma 3.1, so assume $s > 1$. Let $y = [u_2, \dots, u_s]$. Then $x = [u_1] \times y$, and by Lemma 3.1 one has

$$\begin{aligned}([u_1] \times y)P^n\Psi_1^{kp^m} &= \sum_l ([u_1]P^l \times yP^{n-l})\Psi_1^{kp^m} \\ = \sum_l (-1)^{k \deg y}(xP^l\Psi_1^{kp^m} \times yP^{n-l}) &= \sum_l (-1)^{k \deg y}([u_1]\Psi_1^{kp^m}P^l \times yP^{n-l}) \\ &= (-1)^{k \deg y}([u_1]\Psi_1^{kp^m} \times y)P^n = x\Psi_1^{kp^m}P^n.\end{aligned}$$

Here we have used the standard sign convention. □

LEMMA 3.4. *Assume $x = [u_1, \dots, u_s]$, where each u_j is a_j or $\hat{a}_j$. If $a_i \geqslant (p-1)p^m$, then $xP^{p^m}\Psi_i^{(p-1)p^m} - x\Psi_i^{(p-1)p^m}P^{p^m} = x$.*

PROOF. Again one can consider only $i = 1$. For $s = 1$ the statement coincides with Lemma 3.2, so assume $s > 1$. Let $y = [u_2, \dots, u_s]$; then $x = [u_1] \times y$. By Lemma 3.1, one has

$$\begin{aligned}(3.1)\qquad &([u_1] \times y)P^{p^m}\Psi_1^{(p-1)p^m} \\ &\quad = \sum_{l=0}^{p^m-1} ([u_1]P^l \times yP^{p^m-l})\Psi_1^{(p-1)p^m} + ([u_1]P^{p^m} \times y)\Psi_1^{(p-1)p^m} \\ &\quad = \sum_{l=0}^{p^m-1} ([u_1]P^l\Psi_1^{(p-1)p^m} \times yP^{p^m-l}) + ([u_1]P^{p^m}\Psi_1^{(p-1)p^m} \times y);\end{aligned}$$

$$\begin{aligned}(3.2)\qquad &([u_1] \times y)\Psi_1^{(p-1)p^m}P^{p^m} \\ &\quad = \sum_{l=0}^{p^m-1} ([u_1]\Psi_1^{(p-1)p^m}P^l \times yP^{p^m-l}) + ([u_1]\Psi_1^{(p-1)p^m}P^{p^m} \times y).\end{aligned}$$

Take the difference of (3.1) and (3.2) and note that the sums on the right-hand sides cancel by Lemma 3.1. The difference of the last summands in (3.1) and (3.2) is equal to x by Lemma 3.2. □

LEMMA 3.5. *Let $x = [u_1, \dots, u_s]$. Then $x\beta\alpha_i + x\alpha_i\beta = x$.*

PROOF. Again one can consider only $i = 1$. For $s = 1$ the statement coincides with Lemma 3.2(b,c), so assume $s > 1$. Let $y = [u_2, \dots, u_s]$. Then $x = [u_1] \times y$, and by Lemma 3.1 one has

$$\begin{aligned}(3.3)\qquad ([u_1] \times y)\beta\alpha_1 &= ((-1)^{\deg y}[u_1]\beta \times y + [u_1] \times y\beta)\alpha_1 \\ &= [u_1]\beta\alpha_1 \times y + (-1)^{\deg y\beta}[u_1]\alpha_1 \times y\beta.\end{aligned}$$

$$\begin{aligned}(3.4)\qquad ([u_1] \times y)\alpha_1\beta &= (-1)^{\deg y}([u_1]\alpha_1 \times y)\beta \\ &= [u_1]\alpha_1\beta \times y + (-1)^{\deg y}[u_1]\alpha_1 \times y\beta.\end{aligned}$$

Summing (3.3) and (3.4), we obtain

$$(x)\beta\alpha_1 + (x)\alpha_1\beta = [u_1]\beta\alpha_1 \times y + [u_1]\alpha_1\beta \times y = ([u_1]\beta\alpha_1 + [u_1]\alpha_1\beta) \times y$$

which by Lemma 3.2(b,c) is equal to $[u_1] \times y = x$. □

4. Examples

For some $i = 1, \dots, s$ denote by $N_s(i,k)$ the subspace in $\tilde{\Lambda}_{s,*} = \tilde{H}_*(L^{\wedge s})$ spanned by $[u_1, \dots, u_s]$, where $u_i = a_i$ or $u_i = \hat{a}_i$ and $a_i \geqslant (p-1)p^k$.

LEMMA 4.1. *For $s \geqslant 1$ and $k \geqslant 0$ the collection of maps $\{Q_m = \Psi_i^{(p-1)p^m} \mid m \leqslant k\}$ forms a k-th homotopy system for $\tilde{\Lambda}_{s,*}$ with the null space $N_s(i,k)$.*

PROOF. The subspace $N_s(i,k)$ is stable under the action of Q_m. The properties (ii) and (iii) from Definition 2.1 are checked in Lemmas 3.3 and 3.3. The property (iv) from Definition 2.2 is checked in Lemma 3.5. □

THEOREM 4.1. *Fix $s \geqslant 1$ and $k \geqslant 0$. Assume $x \in N_s(i,k)$ for all $1 \leqslant i \leqslant s$. Then $x \in \Delta(k)$ iff $x \in I(k)$ and $x \in \Delta^\beta(k)$ iff $x \in I^\beta(k)$.*

PROOF. The statement follows immediately from Lemma 4.1, Theorem 2.1, and Corollary 2.1. □

References

1. V. Giambalvo, F. P. Peterson, *The annihilator ideal of the action of the Steenrod algebra on $H^*(\mathbb{R}P^\infty)$*, Topology Appl. **65** (1995), 105–122
2. S. Ault, *Relations among the kernels and images of Steenrod squares acting on right $\mathcal{A}$-modules*, J. Pure Appl. Algebra **216** (2012), 1428–1437.
3. S. V. Ault, W. M. Singer, *On the homology of elementary abelian groups as modules over the Steenrod algebra* J. Pure Appl. Algebra **215** (2011), 2847–2852
4. J. M. Boardman, *Modular representations on the homology of powers of real projective space*; in: M. C. Tangora (Ed.), *Algebraic Topology: Oaxtepec 1991*, Contemp. Math. **146** (1993), pp. 49–70
5. M. Kameko, *Generators of the cohomology of BV_3*, J. Math. Kyoto Univ. **38** (3) (1998), 587–593
6. ———, *Generators of the cohomology of BV_4*, Toyama University, 2003 (preprint).
7. N. Sum, *On the hit problem for the polynomial algebra in four variables*, University of Quynhon, Vietnam, 2007 (preprint).

Moscow State Lomonosov University
Department of Mechanics and Mathematics
Moscow, Russia
popelens@mech.math.msu.su