

The Rational qKZ Equation and Shifted Non-Symmetric Jack Polynomials^{*}

Saburo KAKEI[†], Michitomo NISHIZAWA[‡], Yoshihisa SAITO[§] and Yoshihiro TAKEYAMA[¶]

[†] Department of Mathematics, College of Science, Rikkyo University,
Nishi-Ikebukuro, Toshima-ku, Tokyo 171-8501, Japan
E-mail: kakei@rikkyo.ac.jp

[‡] Department of Mathematics, Faculty of Education, Hirosaki University,
1 Bunkyo-cho, Hirosaki, Aomori 036-8560, Japan
E-mail: mnishi@cc.hirosaki-u.ac.jp

[§] Graduate school of Mathematical Sciences, University of Tokyo, Tokyo 153-8914, Japan
E-mail: yoshihisa@ms.u-tokyo.ac.jp

[¶] Department of Mathematics, Graduate School of Pure and Applied Sciences,
University of Tsukuba, Tsukuba, Ibaraki 305-8571, Japan
E-mail: takeyama@math.tsukuba.ac.jp

Received October 15, 2008, in final form January 15, 2009; Published online January 27, 2009
doi:[10.3842/SIGMA.2009.010](https://doi.org/10.3842/SIGMA.2009.010)

Abstract. We construct special solutions to the rational quantum Knizhnik–Zamolodchikov equation associated with the Lie algebra gl_N . The main ingredient is a special class of the shifted non-symmetric Jack polynomials. It may be regarded as a shifted version of the singular polynomials studied by Dunkl. We prove that our solutions contain those obtained as a scaling limit of matrix elements of the vertex operators of level one.

Key words: qKZ equation; shifted Jack polynomial; degenerate double affine Hecke algebra

2000 Mathematics Subject Classification: 39A13; 33C52; 81R50

1 Introduction

The quantum Knizhnik–Zamolodchikov (qKZ) equation, derived by Frenkel and Reshetikhin [9], is a system of difference equations satisfied by matrix elements of vertex operators in the representation theory of the quantum affine algebras. In this paper we consider the rational version of the qKZ equation. Let $V = \bigoplus_{\epsilon=1}^N \mathbb{C}v_{\epsilon}$ be the vector representation of the Lie algebra gl_N . The rational qKZ equation is the following system of difference equations for an unknown function $G(z_1, \dots, z_n)$ taking values in $V^{\otimes n}$:

$$G(z_1, \dots, z_m - \kappa, \dots, z_n) = R_{m,m-1}(z_m - z_{m-1} - \kappa) \cdots R_{m,1}(z_m - z_1 - \kappa) \\ \times \left(\prod_{j=1}^N p_j^{h_j} \right)_m R_{m,n}(z_m - z_n) \cdots R_{m,m+1}(z_m - z_{m+1}) G(z_1, \dots, z_m, \dots, z_n)$$

for $m = 1, \dots, n$. Here h_i ($1 \leq i \leq N$) is a basis of the Cartan subalgebra, p_1, \dots, p_N and κ are nonzero parameters, and $R(z) \in \text{End}(V^{\otimes 2})$ is the rational R -matrix (see (8) below). The lower indices of the operators signify the components of $V^{\otimes n}$ on which they act. The value $-(N + \kappa)$ is called the level of the equation.

^{*}This paper is a contribution to the Special Issue on Dunkl Operators and Related Topics. The full collection is available at <http://www.emis.de/journals/SIGMA/DunklOperators.html>

In [12] special solutions to the qKZ equation are constructed. Let us recall the construction briefly. Consider the coefficients in an expansion of the unknown function with respect to the standard basis of $V^{\otimes n}$. In [12] a sufficient condition is given so that they form a solution to the qKZ equation. It is described as an eigenvalue problem for the q -Dunkl–Cherednik operators and the Demazure–Lusztig operators. The problem can be solved by using the non-symmetric Macdonald polynomials [1, 13], which are by definition eigenfunctions of the q -Dunkl–Cherednik operators. Once one obtains a solution to the eigenvalue problem, the coefficients are generated from it through the action of the Demazure–Lusztig operators. Thus we obtain special solutions to the qKZ equation from the non-symmetric Macdonald polynomials. In this construction the polynomial representation of the affine Hecke algebra is a main tool.

In this article we apply the method of [12] to the rational qKZ equation. The main tool is the degenerate double affine Hecke algebra $\tilde{\mathbf{H}}_\kappa$. It is a trigonometric degeneration of the double affine Hecke algebra introduced by Cherednik [2]. Usually we consider the polynomial representation of $\tilde{\mathbf{H}}_\kappa$ given by the differential operators called the Dunkl–Cherednik operators. They form a commuting family of differential operators and the polynomial eigenfunctions are the non-symmetric Jack polynomials [14]. In our construction of solutions to the rational qKZ equation, we consider another polynomial representation given in terms of commuting difference operators Y_i ($1 \leq i \leq n$), which are difference analogues of the Dunkl operators

$$\mathcal{D}_i := \kappa \frac{\partial}{\partial x_i} + \sum_{j(\neq i)} \frac{1}{x_i - x_j} (1 - \sigma_{ij}),$$

where σ_{ij} is the transposition of variables $x_i \leftrightarrow x_j$. The parameter κ will play the role of the difference step in the rational qKZ equation. The operator Y_i has an expansion $Y_i = 1 - \mathcal{D}_i + \tilde{Y}_i$, where \tilde{Y}_i decreases the total degree by at least two. We can construct another family of commuting difference operators in the same way as the construction of the Dunkl–Cherednik operators from the Dunkl operators. Then the corresponding polynomial eigenfunctions are the shifted non-symmetric Jack polynomials, which are originally obtained by Knop [11] as a classical limit of the non-symmetric quantum Capelli polynomials.

In our construction of special solutions to the rational qKZ equation we need a polynomial eigenfunction of the operators Y_i ($1 \leq i \leq n$). Then, from the expansion $Y_i = 1 - \mathcal{D}_i + \tilde{Y}_i$, the eigenvalue should be one, and the leading degree part of a polynomial eigenfunction should belong to $\cap_{i=1}^n \ker \mathcal{D}_i$ (note that any eigenvalue of the Dunkl operator is equal to zero). A nonzero polynomial in the joint kernel is called a singular polynomial, which is deeply studied by Dunkl [3, 4]. It is proved that constants are only singular polynomials for generic κ , and specific values of κ , called singular values, such that there exists a non-constant singular polynomial are completely determined by Dunkl, de Jeu and Opdam in more general setting associated with finite reflection groups [5]. When κ is equal to a singular value, the space of singular polynomials becomes a non-trivial $\mathbb{C}\mathfrak{S}_n$ -module. In [3, 4] Dunkl determined its irreducible components and proved that all the multiplicities are equal to one. Each irreducible component has a basis consisting of non-symmetric Jack polynomials which are well-defined at the singular value and singular. In this paper we consider the shifted case. We call a polynomial eigenfunction of Y_i ($1 \leq i \leq n$) a *shifted singular polynomial*. In Theorem 4 below we prove that the space of shifted singular polynomials is isomorphic to that of singular polynomials as $\mathbb{C}\mathfrak{S}_n$ -modules, and it has a basis consisting of shifted non-symmetric Jack polynomials. This fact follows from the inversion formula [11], which relates the non-symmetric Jack polynomials and the shifted ones. We can get a solution to the rational qKZ equation from a shifted singular polynomial which satisfies some symmetry conditions (see (10) below).

Now we mention the result [6] by Di Francesco and Zinn-Justin. They constructed a class of polynomial solutions to the qKZ equation and observed that their scaling limit give “extended

Joseph polynomials". As mentioned in [6], the scaling limit of their solutions solves the rational qKZ equation of level one. It had been used in the construction of a new integral formula for solutions to the rational qKZ equation of level zero in [16]. We prove that our solutions contain the special solutions above of level one. In our framework, the extended Joseph polynomials appearing in the solutions to the rational qKZ equation are shifted non-symmetric Jack polynomials which are singular. The leading degree part of the extended Joseph polynomials is nothing but the Joseph polynomial [10] which is a singular non-symmetric Jack polynomial. Thus the inversion formula for (shifted) non-symmetric Jack polynomials at a singular value may be regarded as a map which bridges some data of cohomology of orbital varieties (Joseph polynomials) and their equivariant counterpart (extended Joseph polynomials).

The rest of the paper is organized as follows. In Section 2 we summarize the definitions and properties of (shifted) non-symmetric Jack polynomials from a viewpoint of the representation theory of the degenerate double affine Hecke algebra. In Section 3 we give the rational qKZ equation and a corresponding eigenvalue problem. In Section 4 we construct polynomial solutions to the eigenvalue problem, and hence we obtain special solutions to the rational qKZ equation. The key is the isomorphism as $\mathbb{C}\mathfrak{S}_n$ -modules between the space of singular polynomials and that of shifted singular polynomials. At the very last we prove an explicit formula for some singular (shifted) non-symmetric Jack polynomials.

2 Shifted non-symmetric Jack polynomials

We introduce some notation. We call an element of $\mathbb{Z}_{\geq 0}^n$ a *composition*. In the following we fix n , and denote by Λ the set of compositions. A composition $\lambda = (\lambda_1, \dots, \lambda_n)$ is called *dominant* if $\lambda_1 \geq \dots \geq \lambda_n$. The symmetric group \mathfrak{S}_n acts on Λ by $\sigma\lambda := (\lambda_{\sigma^{-1}(1)}, \dots, \lambda_{\sigma^{-1}(n)})$. For $\lambda \in \Lambda$, denote by λ^+ the unique dominant element in the orbit $\mathfrak{S}_n\lambda$, and by w_λ^+ the shortest element of \mathfrak{S}_n such that $w_\lambda^+\lambda^+ = \lambda$. We define the dominance order \leq on Λ by $\lambda \leq \mu \Leftrightarrow \sum_{i=1}^k \lambda_i \leq \sum_{i=1}^k \mu_i$ ($\forall k$).

2.1 Non-symmetric Jack polynomials

First let us recall the definition of the degenerate double affine Hecke algebra (or the trigonometric Cherednik algebra) following [15]¹. Let κ be an indeterminate and denote the coefficient field by $\mathbb{F} = \mathbb{C}(\kappa)$.

Definition 1. The *degenerate double affine Hecke algebra* $\tilde{\mathbf{H}}_\kappa$ is the unital associative \mathbb{F} -algebra generated by

$$x_i^{\pm 1} \quad (1 \leq i \leq n), \quad s_i \quad (1 \leq i < n), \quad u_i \quad (1 \leq i \leq n)$$

satisfying the following relations:

$$\begin{aligned} [x_i, x_j] &= 0, & [u_i, u_j] &= 0, \\ s_i^2 &= 1, & s_i s_{i+1} s_i &= s_{i+1} s_i s_{i+1}, & s_i s_j &= s_j s_i & (|i - j| > 1), \\ s_i x_i s_i &= x_{i+1}, & s_i x_j &= x_j s_i & (j \neq i, i + 1), \\ s_i u_i &= u_{i+1} s_i + 1, & s_i u_j &= u_j s_i & (j \neq i, i + 1), \\ [u_i, x_j] &= \begin{cases} -x_i s_{ji} & (j < i), \\ \kappa x_j + x_j \sum_{1 \leq k < j} s_{kj} + \sum_{j < k \leq n} s_{jk} x_j & (j = i), \\ -x_j s_{ij} & (i < j). \end{cases} \end{aligned}$$

Here $s_{ij} = (s_i \cdots s_{j-1})(s_{j-2} \cdots s_i)$ for $i < j$.

¹We change the notation in [15] as $x_i \rightarrow x_{n+1-i}$, $s_i \rightarrow s_{n-i}$, $u_i \rightarrow u_{n+1-i}$.

Note that $\widetilde{\mathbf{H}}_\kappa$ contains the group algebra of the symmetric group $\langle s_1, \dots, s_{n-1} \rangle \simeq \mathbb{F}\mathfrak{S}_n$.

Denote by $\widetilde{\mathbf{H}}_\kappa^+$ the subalgebra generated by x_i, u_i ($1 \leq i \leq n$) and s_i ($1 \leq i < n$). The algebra $\widetilde{\mathbf{H}}_\kappa^+$ and its polynomial representation are crucial tools in the theory of Jack polynomials. Let $\mathbb{F}[x] = \mathbb{F}[x_1, \dots, x_n]$ be the polynomial ring. Denote by $\sigma_{i,j}$ the transposition of variables $x_i \leftrightarrow x_j$. Then the polynomial representation $\phi : \widetilde{\mathbf{H}}_\kappa^+ \rightarrow \text{End } \mathbb{F}[x]$ is defined by

$$\phi(x_i) = x_i, \quad \phi(s_i) = \sigma_{i,i+1}, \quad \phi(u_i) = x_i \mathcal{D}_i + \sum_{i < k \leq n} \sigma_{ik}, \quad (1)$$

where \mathcal{D}_i is the *Dunkl operator*

$$\mathcal{D}_i := \kappa \frac{\partial}{\partial x_i} + \sum_{j(\neq i)} \frac{1}{x_i - x_j} (1 - \sigma_{ij}). \quad (2)$$

The operators $\phi(u_i)$ ($1 \leq i \leq n$) are called the *Dunkl–Cherednik operators*.

The *non-symmetric Jack polynomials* ξ_λ ($\lambda \in \Lambda$) are uniquely defined by the following conditions:

$$\phi(u_i)\xi_\lambda = c_i(\lambda)\xi_\lambda \quad (1 \leq i \leq n), \quad \xi_\lambda(x) = x^\lambda + \sum_{\mu < \lambda} c_{\lambda\mu} x^\mu \quad (c_{\lambda\mu} \in \mathbb{F}).$$

Here $x^\lambda = x_1^{\lambda_1} \cdots x_n^{\lambda_n}$, and $c(\lambda) = (c_1(\lambda), \dots, c_n(\lambda)) \in \mathbb{F}^n$ is defined by

$$c(\lambda) := \kappa\lambda + w_\lambda^+ \rho,$$

where $\rho := (n-1, \dots, 1, 0)$.

2.2 Shifted non-symmetric Jack polynomials

The shifted non-symmetric Jack polynomials are defined by the vanishing conditions. Hereafter we use the variables $z = (z_1, \dots, z_n)$ for the shifted non-symmetric Jack polynomials, and denote by $\mathbb{F}[z] = \mathbb{F}[z_1, \dots, z_n]$ the ring of polynomials in z_1, \dots, z_n .

Theorem 1 ([11, Theorem 6.2]). *For $\lambda \in \Lambda$ there exists a polynomial $E_\lambda(z) \in \mathbb{F}[z]$ uniquely determined by the following conditions:*

$$\begin{aligned} E_\lambda(c(\mu)) &= 0 \quad \text{for any } \mu \in \Lambda \text{ such that } |\mu| \leq |\lambda|, \mu \neq \lambda, \\ E_\lambda(z) &= \sum_{|\nu| \leq |\lambda|} d_{\lambda\nu} z^\nu \quad (d_{\lambda\nu} \in \mathbb{F}, d_{\lambda\lambda} = 1). \end{aligned}$$

Here $|\lambda| := \sum_{i=1}^n \lambda_i$. We call E_λ ($\lambda \in \Lambda$) the *shifted non-symmetric Jack polynomials*.

Remark 1. The shifted non-symmetric Jack polynomials defined in Theorem 1 are slightly different from the original ones given by Knop [11] as follows. In [11] the parameter is denoted by r . Let $\widetilde{\mathbb{F}} := \mathbb{C}(r)$ be the corresponding coefficient field. Define a \mathbb{C} -algebra isomorphism $\Psi : \mathbb{F}[z] \rightarrow \widetilde{\mathbb{F}}[z]$ by $\Psi(\kappa) = 1/r$ and $\Psi(z_i) = n-1 + z_i/r$. Then the polynomials \widetilde{E}_λ defined in [11, Theorem 6.2] are related to E_λ in Theorem 1 by $\Psi(E_\lambda) = r^{-|\lambda|} \widetilde{E}_\lambda$.

The shifted non-symmetric Jack polynomials are also characterized as joint eigenfunctions of some commuting difference operators. To construct the operators, we introduce another polynomial representation of $\widetilde{\mathbf{H}}_\kappa$. Define the operators T_i ($1 \leq i < n$) and ω acting on $\mathbb{F}[z]$ by

$$T_i := \sigma_{i,i+1} + \frac{1}{z_i - z_{i+1}} (1 - \sigma_{i,i+1}), \quad (3)$$

$$(\omega f)(z_1, \dots, z_n) := f(z_n - \kappa, z_1, \dots, z_{n-1}). \quad (4)$$

Here $\sigma_{i,i+1}$ is the transposition of variables $z_i \leftrightarrow z_{i+1}$. Set

$$Y_i := T_i \cdots T_{n-1} \omega T_1 \cdots T_{i-1} \quad (1 \leq i \leq n). \quad (5)$$

Proposition 1. *There exists an \mathbb{F} -algebra homomorphism $\psi : \widetilde{\mathbf{H}}_\kappa \rightarrow \text{End } \mathbb{F}[z]$ such that*

$$\psi(x_i) = Y_i, \quad \psi(s_i) = T_i, \quad \psi(u_i) = z_i.$$

Proof. It can be checked by direct calculation using $\omega T_i = T_{i-1} \omega$ ($1 < i < n$). \blacksquare

Note that the generator x_i acts as a difference operator and u_i as the multiplication of the variable z_i on $\mathbb{F}[z]$. It is opposite to the action ϕ (1).

Remark 2. The usual polynomial representation ϕ is induced from the one-dimensional representation of the subalgebra generated by s_i ($1 \leq i < n$) and u_i ($1 \leq i \leq n$). The representation ψ in Proposition 1 is constructed from an opposite side in the following sense. Consider the subalgebra generated by s_i ($1 \leq i < n$) and x_i ($1 \leq i \leq n$). It is isomorphic to the group algebra of the extended affine Weyl group $\widetilde{\mathfrak{S}}_n$. Then the induced module of the trivial $\mathbb{C}\widetilde{\mathfrak{S}}_n$ -module is isomorphic to $\mathbb{F}[u_1, \dots, u_n]$ as an \mathbb{F} -vector space. Change the variables $u_i \rightarrow z_i$ and we obtain the polynomial representation ψ .

Proposition 2. *Define $v_i \in \widetilde{\mathbf{H}}_\kappa$ ($1 \leq i \leq n$) by*

$$v_i := u_i(1 - x_i) + \sum_{i < k \leq n} s_{ik} x_i.$$

Then the following relations hold:

$$[v_i, v_j] = 0, \quad s_i v_i = v_{i+1} s_i + 1, \quad s_i v_j = v_j s_i \quad (j \neq i, i+1).$$

Proof. Set $\widehat{u}_i := u_i - v_i = (u_i - \sum_{i < k \leq n} s_{ik}) x_i$. Then we can check that $[\widehat{u}_i, \widehat{u}_j] = 0$ and $[v_j, \widehat{u}_i] = [v_i, \widehat{u}_j] = -\widehat{u}_j s_{ij}$ for $i < j$. Thus we get the first relation. The others can be proved by easy calculation. \blacksquare

Thus we obtain commuting difference operators $\psi(v_i)$ ($1 \leq i \leq n$) acting on $\mathbb{F}[z]$. The shifted non-symmetric Jack polynomials are joint eigenfunctions of them.

Theorem 2 ([11, Theorem 6.6]). *The shifted non-symmetric Jack polynomials E_λ ($\lambda \in \Lambda$) are uniquely characterized by the following properties:*

$$\begin{aligned} \psi(v_i) E_\lambda &= c_i(\lambda) E_\lambda \quad (1 \leq i \leq n), \\ E_\lambda(z) &= \sum_{|\nu| \leq |\lambda|} d_{\lambda\nu} z^\nu \quad (d_{\lambda\nu} \in \mathbb{F}, d_{\lambda\lambda} = 1). \end{aligned}$$

Remark 3. The operators $\psi(v_i)$ ($1 \leq i \leq n$) are related to $\widetilde{\Xi}_i$'s in [11, Theorem 6.6] by $\Psi^{-1} \widetilde{\Xi}_i \Psi = (\psi(v_i) - (n-1))/\kappa$, where Ψ is the map defined in Remark 1.

Here we briefly explain why the eigenvalues of $\psi(v_i)$ ($1 \leq i \leq n$) are the same as those of the Dunkl–Cherednik operators $\phi(u_i)$ ($1 \leq i \leq n$). Since the operator $T_i - \sigma_i$ decreases the total degree by one, we have an expansion $Y_i = 1 - \mathcal{D}_i + \widetilde{Y}_i$, where \mathcal{D}_i is the Dunkl operator (2), and \widetilde{Y}_i is an operator which decreases the total degree by at least two. Thus we see that $\psi(v_i) - \phi(u_i)$ is an operator which strictly decreases the total degree. Hence $\psi(v_i)$ is triangular with respect to the order on monomials in $\mathbb{F}[z]$ induced from the dominance order \leq and the total degree, and its eigenvalues are equal to those of $\phi(u_i)$.

Proposition 3. *Any eigenvalue of the operators Y_i ($1 \leq i \leq n$) on $\mathbb{F}[z]$ is equal to one.*

Proof. From the argument above, the operator $Y_i - 1 = -\mathcal{D}_i + \widetilde{Y}_i$ decreases the total degree by at least one. This implies the proposition. \blacksquare

2.3 The inversion formula

The $\tilde{\mathbf{H}}_\kappa$ -module $\mathbb{F}[z]$ is naturally an $\tilde{\mathbf{H}}_\kappa^+$ -module by restriction. However we consider another $\tilde{\mathbf{H}}_\kappa^+$ -module structure defined through the following homomorphism.

Proposition 4 ([7]). *Set $\hat{u}_i := u_i - v_i = (u_i - \sum_{i < k \leq n} s_{ik})x_i$ ($1 \leq i \leq n$). Then there exists an \mathbb{F} -algebra homomorphism τ satisfying*

$$\tau(x_i) = \hat{u}_i, \quad \tau(s_i) = s_i, \quad \tau(u_i) = v_i.$$

Hereafter we regard $\mathbb{F}[z]$ as an $\tilde{\mathbf{H}}_\kappa^+$ -module through the map $\psi \circ \tau$. To clarify it we write ${}^\tau\mathbb{F}[z]$ instead of $\mathbb{F}[z]$ when we consider the $\tilde{\mathbf{H}}_\kappa^+$ -module structure. On the other hand we regard $\mathbb{F}[x]$ as an $\tilde{\mathbf{H}}_\kappa^+$ -module by the restriction $\phi|_{\tilde{\mathbf{H}}_\kappa^+}$.

Proposition 5. *The map*

$$\Phi : \mathbb{F}[x] \longrightarrow {}^\tau\mathbb{F}[z], \quad f(x_1, \dots, x_n) \mapsto \psi(f(\hat{u}_1, \dots, \hat{u}_n))1 \quad (6)$$

is an isomorphism between $\tilde{\mathbf{H}}_\kappa^+$ -modules. The inverse Φ^{-1} is a map taking the leading degree part and changing the variables $z_i \rightarrow x_i$.

Proof. The intertwining property follows from $\phi(s_i)1 = 1$, $\phi(u_i)1 = n - i$ in $\mathbb{F}[x]$, and $\psi(\tau(s_i))1 = \psi(s_i)1 = 1$ and $\psi(\tau(u_i))1 = \psi(v_i)1 = n - i$ in ${}^\tau\mathbb{F}[z]$. The operator $\psi(\hat{u}_i) - z_i$ decreases the total degree of polynomials in z , and hence $\Phi(x^\lambda) = z^\lambda + (\text{lower degree terms})$ for any $\lambda \in \Lambda$. Therefore Φ is an isomorphism. \blacksquare

Corollary 1 ([11, Theorem 6.9]). $\Phi(\xi_\lambda) = E_\lambda$ for any $\lambda \in \Lambda$. Hence the leading degree part of E_λ is equal to ξ_λ .

Proof. Apply Φ to the both hand sides of $\phi(u_i)\xi_\lambda = c_i(\lambda)\xi_\lambda$. Then we see that $\Phi(\xi_\lambda)$ is an eigenfunction of $\psi(v_i)$ with the eigenvalue $c_i(\lambda)$. The coefficient of z^λ in $\Phi(\xi_\lambda)$ is equal to one. This implies that $\Phi(\xi_\lambda) = E_\lambda$. \blacksquare

Remark 4. The operators $\psi(\hat{u}_i)$ ($1 \leq i \leq n$) are related to \tilde{Z}_i 's in [11, Theorem 6.9] by $\Psi^{-1}\tilde{Z}_i\Psi = \psi(\hat{u}_i)/\kappa$, where Ψ is the map defined in Remark 1.

The equality $\Phi(\xi_\lambda) = E_\lambda$ is called the *inversion formula*.

The non-symmetric Jack polynomials ξ_λ may have poles when κ is a non-positive rational number. From the inversion formula, we see that the shifted one E_λ also has the same property. We say that ξ_λ (or E_λ) is *well-defined at $\kappa = \kappa_0 \in \mathbb{Q}_{\leq 0}$* if it does not have a pole at $\kappa = \kappa_0$.

Corollary 2. *For $\lambda \in \Lambda$, ξ_λ is well-defined at $\kappa = \kappa_0 \in \mathbb{Q}_{\leq 0}$ if and only if E_λ is well-defined.*

In Section 4.2 we specialize κ to some constant $\kappa = \kappa_0 \in \mathbb{Q}^\times$. Then the \mathbb{C} -algebra $\tilde{\mathbf{H}}_{\kappa_0}^+$ acts on $\mathbb{C}[x]$ and ${}^\tau\mathbb{C}[z]$ through the maps ϕ and $\psi \circ \tau$, respectively. In this situation the map

$$\Phi : \mathbb{C}[x] \longrightarrow {}^\tau\mathbb{C}[z], \quad f(x_1, \dots, x_n) \mapsto \psi(f(\hat{u}_1, \dots, \hat{u}_n))1 \quad (7)$$

is still an isomorphism between $\tilde{\mathbf{H}}_{\kappa_0}^+$ -modules.

3 The rational qKZ equation

3.1 The rational qKZ equation

Let $V = \bigoplus_{\epsilon=1}^N \mathbb{C}v_\epsilon$ be the N -dimensional vector space. We regard V as the vector representation of the Lie algebra gl_N . Denote by h_i ($1 \leq i \leq N$) a basis of the Cartan subalgebra, which acts on V by $h_i v_\epsilon = \delta_{i,\epsilon} v_\epsilon$.

The rational R -matrix $R(z) \in \text{End}(V^{\otimes 2})$ is defined by

$$R(z) = \frac{z + P}{z + 1}, \quad (8)$$

where P is the transposition: $P(u \otimes v) = v \otimes u$.

Fix a parameter $\kappa \in \mathbb{C}^\times$. The *rational quantum Knizhnik–Zamolodchikov (rational qKZ) equation* is the following system of difference equations for an unknown function $G(z_1, \dots, z_n)$ taking values in $V^{\otimes n}$:

$$\begin{aligned} G(z_1, \dots, z_m - \kappa, \dots, z_n) &= R_{m,m-1}(z_m - z_{m-1} - \kappa) \cdots R_{m,1}(z_m - z_1 - \kappa) \\ &\times \left(\prod_{j=1}^N p_j^{h_j} \right)_m R_{m,n}(z_m - z_n) \cdots R_{m,m+1}(z_m - z_{m+1}) G(z_1, \dots, z_m, \dots, z_n) \end{aligned}$$

for $m = 1, \dots, n$. Here p_1, \dots, p_N are nonzero parameters, and the lower indices of the operators signify the components of $V^{\otimes n}$ on which they act. The value $-(N + \kappa)$ is called the *level* of the equation.

3.2 Eigenvalue problem associated with the rational qKZ equation

Hereafter we assume that

$$n \geq N \geq 2.$$

Fix positive integers d_1, \dots, d_N satisfying $\sum_{j=1}^N d_j = n$, and define an n -tuple δ by $\delta = (1^{d_1}, \dots, N^{d_N})$. Consider the following eigenvalue problem for a function $F_\delta = F_\delta(z_1, \dots, z_n)$:

$$Y_i F_\delta = \chi_i F_\delta \quad (1 \leq i \leq n, \chi_i \in \mathbb{C}^\times), \quad (9)$$

$$T_i F_\delta = -F_\delta \quad \text{if } \delta_i = \delta_{i+1}. \quad (10)$$

Here Y_i and T_i are given by (5) and (3), respectively. Note that the consistency of (9) and (10) implies that $\chi_i = \chi_{i+1}$ if $\delta_i = \delta_{i+1}$. Hence there are N independent parameters $\chi_{d_1+\dots+d_j}$ ($1 \leq j \leq N$).

Once one solves the eigenvalue problem above, a solution to the rational qKZ equation can be constructed as follows. Suppose that F_δ is a solution to the eigenvalue problem (9) and (10). Set $I_{d_1, \dots, d_N} := \mathfrak{S}_n \delta \subset \{1, \dots, N\}^n$. For $\epsilon = (\epsilon_1, \dots, \epsilon_n) \in I_{d_1, \dots, d_N}$, take a shortest element $w_\epsilon \in \mathfrak{S}_n$ such that $w_\epsilon \epsilon = \delta$. Let $w_\epsilon = s_{i_1} \cdots s_{i_\ell}$ be a reduced expression, and set $T_{w_\epsilon} := T_{i_1} \cdots T_{i_\ell}$. This does not depend on a choice of the reduced expression. Now define a function $F_\epsilon(z_1, \dots, z_n)$ by

$$F_\epsilon := (-1)^{\ell(w_\epsilon)} T_{w_\epsilon} F_\delta,$$

where $\ell(w)$ is the length of $w \in \mathfrak{S}_n$. Set

$$F(z_1, \dots, z_n) := \sum_{\epsilon \in I_{d_1, \dots, d_N}} F_\epsilon(z_1, \dots, z_n) v_{\epsilon_1} \otimes \cdots \otimes v_{\epsilon_n}.$$

Let $K(z_1, \dots, z_n)$ be a function satisfying

$$\frac{K(\dots, z_m - \kappa, \dots)}{K(\dots, z_m, \dots)} = \prod_{j=1}^{m-1} \frac{z_j - z_m + \kappa + 1}{z_j - z_m + \kappa - 1} \prod_{j=m+1}^n \frac{z_m - z_j - 1}{z_m - z_j + 1} \quad (11)$$

for $m = 1, \dots, n$. For example, take

$$K(z_1, \dots, z_n) = \prod_{1 \leq i < j \leq n} \frac{\Gamma\left(-\frac{z_i - z_j - 1}{\kappa}\right)}{\Gamma\left(-\frac{z_i - z_j + 1}{\kappa}\right)}. \quad (12)$$

Theorem 3. Set $G(z_1, \dots, z_n) := K(z_1, \dots, z_n)F(z_1, \dots, z_n)$. Then G is a solution of the rational qKZ equation with $p_j = \chi_{d_1 + \dots + d_j}$ ($1 \leq j \leq N$).

Proof. In the same way as the proof of Theorem 3.6 in [12], we can see that the following relations hold:

$$\begin{aligned} T_i F_{\dots, \epsilon_i, \epsilon_{i+1}, \dots} &= -F_{\dots, \epsilon_{i+1}, \epsilon_i, \dots}, \\ \omega F_{\epsilon_n, \epsilon_1, \dots, \epsilon_{n-1}} &= (-1)^{n-1} \chi_{d_1 + \dots + d_{\epsilon_n}} F_{\epsilon_1, \dots, \epsilon_n}, \end{aligned}$$

where ω is defined by (4). By setting $p_j = \chi_{d_1 + \dots + d_j}$, the two relations above are equivalent to

$$\begin{aligned} \frac{z_i - z_{i+1} + 1}{z_{i+1} - z_i + 1} P_{i, i+1} R_{i, i+1}(z_i - z_{i+1}) F(\dots, z_i, z_{i+1}, \dots) &= F(\dots, z_{i+1}, z_i, \dots), \\ P_{n-1, n} \cdots P_{1, 2} F(z_n - \kappa, z_1, \dots, z_{n-1}) &= (-1)^{n-1} \left(\prod_{j=1}^N p_j^{h_j} \right)_n F(z_1, \dots, z_n). \end{aligned}$$

Combining them and (11), we see that $G = KF$ is a solution to the rational qKZ equation. ■

4 Special solutions to the rational qKZ equation

Now we construct a class of polynomial solutions to the eigenvalue problem (9) and (10). They create solutions to the rational qKZ equation as discussed in the previous subsection (see Theorem 3).

4.1 Shifted singular polynomials

First let us consider the equation (9). We want to obtain polynomial solutions. Then the eigenvalues χ_i ($1 \leq i \leq n$) should be one from Proposition 3. As seen below the polynomial eigenfunctions can be regarded as a shifted version of singular polynomials.

A polynomial $f(x_1, \dots, x_n) \in \mathbb{C}[x]$ is called a *singular polynomial* if $f \in \cap_{i=1}^n \ker \mathcal{D}_i$ [3, 4, 5]. Denote by \mathcal{S} the subspace of $\mathbb{C}[x]$ consisting of singular polynomials. Recall that $\mathbb{C}[x]$ has $\tilde{\mathbf{H}}_\kappa$ -module structure defined by the map ϕ (1). From the commutation relations $\sigma_i \mathcal{D}_i = \mathcal{D}_{i+1} \sigma_i$ and $\sigma_i \mathcal{D}_j = \mathcal{D}_j \sigma_i$ ($j \neq i, i+1$), the maps $\phi(s_i)$ ($1 \leq i < n$) preserve \mathcal{S} . Thus \mathcal{S} is a $\mathbb{C}\mathfrak{S}_n$ -module.

Now let us consider the shifted version.

Definition 2. We call $g(z_1, \dots, z_n) \in \mathbb{C}[z]$ a *shifted singular polynomial* if it satisfies $Y_i g = g$ for all $1 \leq i \leq n$.

Denote by $\tilde{\mathcal{S}}$ the subspace of $\mathbb{C}[z]$ consisting of shifted singular polynomials. From the relations $T_i Y_i = Y_{i+1} T_i$ and $T_i Y_j = Y_j T_i$ ($j \neq i, i+1$), which follow from Proposition 1, the maps T_i ($1 \leq i < n$) preserve $\tilde{\mathcal{S}}$. Hence $\tilde{\mathcal{S}}$ is also a $\mathbb{C}\mathfrak{S}_n$ -module. It is naturally obtained from the $\tilde{\mathbf{H}}_\kappa^+$ -module structure on ${}^\tau \mathbb{C}[z]$ (note that $\tau(s_i) = s_i$ for all i).

Theorem 4. *The map Φ (7) gives a $\mathbb{C}\mathfrak{S}_n$ -module isomorphism $\Phi|_{\mathcal{S}} : \mathcal{S} \simeq \tilde{\mathcal{S}}$.*

Proof. Recall that Φ^{-1} is the map taking the leading degree part and changing the variables $z_i \rightarrow x_i$. Then we easily see $\Phi^{-1}(\tilde{\mathcal{S}}) \subset \mathcal{S}$ from the expansion $Y_i = 1 - \mathcal{D}_i + \tilde{Y}_i$ given in the argument preceding to Proposition 3.

Let us prove $\Phi(\mathcal{S}) \subset \tilde{\mathcal{S}}$. Suppose that $f \in \mathbb{C}[x]$ be a singular polynomial. Then we have $x_i \mathcal{D}_i f = 0$ ($1 \leq i \leq n$). From $x_i \mathcal{D}_i = \phi(u_i - \sum_{i < k \leq n} s_{ik})$ and the intertwining property of Φ , we have

$$\begin{aligned} 0 &= \Phi(x_i \mathcal{D}_i f) = \psi \left(\tau \left(u_i - \sum_{i < k \leq n} s_{ik} \right) \right) \Phi(f) \\ &= \psi(v_i - \sum_{i < k \leq n} s_{ik}) \Phi(f) = \left(z_i - \sum_{i < k \leq n} T_{ik} \right) (1 - Y_i) \Phi(f), \end{aligned}$$

where $T_{ik} = \psi(s_{ik})$. Since the operator $z_i - \sum_{i < k \leq n} T_{ik}$ is injective, we get $Y_i \Phi(f) = \Phi(f)$ for $1 \leq i \leq n$. ■

If κ is generic, we have $\mathcal{S} = \mathbb{C}$, that is, constants are only singular polynomials. A specific parameter value $\kappa = \kappa_0$ is called a *singular value* if there exists a non-constant singular polynomial at $\kappa = \kappa_0$. In [5] it is proved that the set of singular values is equal to $\{-l/m \mid m \in \mathbb{Z}_{>0}, l = 2, \dots, n \text{ and } m \notin l\mathbb{Z}\}^2$. From Corollary 1, Corollary 2 and Theorem 4, we find

Corollary 3. *Let κ_0 be a singular value. Suppose that λ is a composition such that ξ_λ is well-defined and singular at $\kappa = \kappa_0$. Then E_λ is also well-defined and shifted singular.*

In [3, 4] Dunkl obtained an explicit description of \mathcal{S} as a $\mathbb{C}\mathfrak{S}_n$ -module. At a singular value $\kappa = \kappa_0$, the module \mathcal{S} is multiplicity-free and the set of the irreducible components is completely determined from κ_0 and n . Each irreducible component has a basis consisting of non-symmetric Jack polynomials which are well-defined and singular. From Theorem 4 and Corollary 3, the $\mathbb{C}\mathfrak{S}_n$ -module structure of $\tilde{\mathcal{S}}$ is the same as \mathcal{S} . In the shifted case, $\tilde{\mathcal{S}}$ has a basis consisting of shifted non-symmetric Jack polynomials.

Now we give a limiting procedure from non-symmetric Macdonald polynomials to shifted singular ones. Let us recall the definition of non-symmetric Macdonald polynomials. Let q and $t^{1/2}$ be indeterminates and denote the coefficient field by $\mathbb{K} = \mathbb{C}(q, t^{1/2})$. Consider the following operators \hat{T}_i ($1 \leq i < n$) and $\hat{\omega}$ acting on the polynomial ring $\mathbb{K}[X_1, \dots, X_n]$:

$$\hat{T}_i := t^{1/2} \sigma_i + \frac{t^{1/2} - t^{-1/2}}{X_i/X_{i+1} - 1} (\sigma_i - 1), \quad (\hat{\omega}f)(X_1, \dots, X_n) := f(qX_n, X_1, \dots, X_{n-1}),$$

where σ_i is the transposition of variables $X_i \leftrightarrow X_{i+1}$. The operators \hat{T}_i ($1 \leq i < n$) are called the *Demazure–Lusztig operators*. Then the *q-Dunkl–Cherednik operators* \hat{Y}_i ($1 \leq i \leq n$) are defined by

$$\hat{Y}_i := \hat{T}_i \cdots \hat{T}_{n-1} \hat{\omega} \hat{T}_1^{-1} \cdots \hat{T}_{i-1}^{-1}.$$

The *non-symmetric Macdonald polynomials* $E_\lambda^{q,t}$ ($\lambda \in \Lambda$) are defined by the following properties:

$$\hat{Y}_i E_\lambda^{q,t} = t^{(w_\lambda^+ \rho)_i} q^{\lambda_i} E_\lambda^{q,t} \quad (1 \leq i \leq n), \quad E_\lambda^{q,t}(X) = X^\lambda + \sum_{\mu < \lambda} a_{\mu\lambda} X^\mu \quad (a_{\mu\lambda} \in \mathbb{K}).$$

Here $X^\lambda := X_1^{\lambda_1} \cdots X_n^{\lambda_n}$.

²Note that the parameter κ in [3, 4] is equal to $1/\kappa$ in this article. See the definition of the Dunkl operator (2).

Introduce a small parameter ϵ and set

$$X_j = e^{\epsilon z_j}, \quad t^{1/2} = e^{-\epsilon/2}, \quad q = e^{-\kappa\epsilon}. \quad (13)$$

Then we find $\widehat{T}_i^{\pm 1} \rightarrow T_i$ and $\widehat{\omega} \rightarrow \omega$ in the limit $\epsilon \rightarrow 0$, and hence $\widehat{Y}_i \rightarrow Y_i$. Suppose that $E_\lambda^{q,t}$ has the Laurent expansion at $\epsilon = 0$ in the form

$$E_\lambda^{q,t}(X)|_{q=t^\kappa} = \epsilon^\alpha (g_\lambda(z) + o(1))$$

for some $\alpha \in \mathbb{Z}$ and nonzero $g_\lambda(z) \in \mathbb{F}[z]$. Then we obtain $Y_i g_\lambda = g_\lambda$ for $1 \leq i \leq n$, that is, g_λ is shifted singular. From the discussion preceding to Corollary 3, g_λ can become non-constant only if κ is a singular value. In Proposition 7 below, we give an example of λ and a singular value κ_0 such that g_λ is non-constant at $\kappa = \kappa_0$.

4.2 Construction of special solutions

To construct a solution to the rational qKZ equation, we need a shifted singular polynomial satisfying the condition (10).

Let r and k be positive integers such that $r \geq 2$, $k \leq n$ and $\gcd(r-1, k+1) = 1$. Hereafter we set³

$$\kappa = -\frac{k+1}{r-1}. \quad (14)$$

Take three integers s, m and l satisfying

$$n = (s+m)k + s + l, \quad s \geq 0, \quad m \geq 1, \quad 0 \leq l < k.$$

For $a \geq 1$, we set

$$\lambda[r, s, a] := ((r-1)(s+a), (r-1)(s+a-1), \dots, (r-1)(s+1), 0).$$

For $a = 0$, set $\lambda[r, s, 0] = (0)$. Now we define the composition $\lambda(k, s, m, l, r) \in \Lambda$ by

$$\lambda(k, s, m, l, r) := (0^{(k+1)s}, \lambda[r, s, m-1]^{k-l}, \lambda[r, s, m]^l).$$

Next we take an N -tuple (d_1, \dots, d_N) satisfying the following conditions. For $1 \leq i \leq (k+1)s$, we have $d_i = 1$. The rest $(d_{(k+1)s+1}, \dots, d_N)$ should be a subdivision of $(m^{k-l}, (m+1)^l)$, that is, $d_i > 0$ and

$$d_{i_j} + \dots + d_{i_{j+1}-1} = \begin{cases} m & (1 \leq j \leq k-l), \\ m+1 & (k-l < j \leq k) \end{cases}$$

for some $i_1 = (k+1)s + 1 < i_2 < \dots < i_k < i_{k+1} = N + 1$.

Now set $\delta = (1^{d_1}, \dots, N^{d_N})$. Then we find

Proposition 6. $E_{\lambda(k,s,m,l,r)}$ is well-defined at (14) and satisfies (9) and (10) with $\chi_i = 1$ ($1 \leq i \leq n$). Therefore the shifted non-symmetric Jack polynomial $E_{\lambda(k,s,m,l,r)}$ at (14) creates a solution to the rational qKZ equation with $p_j = 1$ ($1 \leq j \leq N$) of level $\frac{k+1}{r-1} - N$.

³The parametrization (14) is motivated by the condition of (k, r) -admissibility employed in the construction of special solutions to the (trigonometric) qKZ equation [12]. It is originally appeared in [8].

Proof. In [3, Theorem 5.7] it is proved that $\xi_{\lambda(k,s,m,l,r)}$ is well-defined and singular at (14) (note that if $n = k$ then $(s, m, l) = (0, 1, 0)$ and $E_{\lambda(k,0,1,0,r)} = E_{0^k} = 1$ is trivially well-defined and singular). Hence, from Corollary 3, $E_{\lambda(k,s,m,l,r)}$ is also well-defined and satisfies (9).

To prove that $E_{\lambda(k,s,m,l,r)}$ satisfies (10), we need the formula for the action of T_i [7]:

$$T_i E_\mu = \begin{cases} a_i(\mu)E_\mu + b_i(\mu)E_{s_i\mu} & (\mu_i > \mu_{i+1}), \\ E_\mu & (\mu_i = \mu_{i+1}), \\ a_i(\mu)E_\mu + E_{s_i\mu} & (\mu_i < \mu_{i+1}). \end{cases} \quad (15)$$

Here $a_i(\mu)$ and $b_i(\mu)$ are defined by

$$a_i(\mu) := \frac{1}{c_i(\mu) - c_{i+1}(\mu)}, \quad b_i(\mu) := 1 - a_i(\mu)^2.$$

The formula (15) can be obtained from the corresponding formula for the non-symmetric Jack polynomials and the isomorphism Φ (6).

Now let us check (10). We abbreviate $\lambda(k, s, m, l, r)$ to λ in the rest of the proof. If $\delta_i = \delta_{i+1}$, we have $\lambda_i > \lambda_{i+1}$, $a_i(\lambda) = -1$ and $b_i(\lambda) = 0$. Hence we get $T_i E_\lambda = -E_\lambda$ from (15). ■

4.3 Factorized formula for the case of $(k, r, s) = (N, 2, 0)$

In [16] a new integral formula for solutions of the rational qKZ equation of level zero is constructed. In the construction we use as an ingredient a special solution of level one given as follows. We consider the case of $n = Nm$ for some $m \geq 1$, and $d_1 = \dots = d_N = m$. Then the special solution is given in the form $G = KF$, where $K = K(z_1, \dots, z_n)$ is the function defined by (12), and

$$F_\delta = \prod_{\substack{1 \leq i < j \leq n \\ \delta_i = \delta_j}} (z_i - z_j - 1).$$

It is obtained as a limit of a matrix element of the vertex operators associated with the quantum affine algebra $U_q(\widehat{sl}_N)$. It is also found by Di Francesco and Zinn-Justin in [6].

The special solution above is contained in the polynomial solutions given in Proposition 6 as follows.

Proposition 7. *Let $(k, r) = (N, 2)$. For $n = Nm + l$ ($m \geq 1, 0 \leq l < N$), consider*

$$\lambda(N, 0, m, l, 2) = ((m-1, \dots, 1, 0)^{N-l}, (m, \dots, 1, 0)^l).$$

Then we have

$$\xi_{\lambda(N,0,m,l,2)}(x_1, \dots, x_n) \Big|_{\kappa=-(N+1)} = \prod_{\substack{1 \leq i < j \leq n \\ \delta_i = \delta_j}} (x_i - x_j), \quad (16)$$

$$E_{\lambda(N,0,m,l,2)}(z_1, \dots, z_n) \Big|_{\kappa=-(N+1)} = \prod_{\substack{1 \leq i < j \leq n \\ \delta_i = \delta_j}} (z_i - z_j - 1), \quad (17)$$

where $\delta = (1^m, \dots, (N-l)^m, (N-l+1)^{m+1}, \dots, N^{m+1})$.

Proof. Hereafter we set $\kappa = -(N+1)$ and abbreviate $\lambda_0 = \lambda(N, 0, m, l, 2)$. Denote the right hand sides of (16) and (17) by $\tilde{\xi}(x)$ and $\tilde{E}(z)$, respectively.

In [12, Proposition 5.4] the following formula is proved:

$$E_{\lambda_0}^{q,t}(X) \Big|_{q=t^{-(N+1)}} = \prod_{\substack{1 \leq i < j \leq n \\ \delta_i = \delta_j}} (X_i - t^{-1}X_j),$$

where $E_{\lambda_0}^{q,t}$ is the non-symmetric Macdonald polynomial. Applying the substitution (13), we see that the right hand side above has the expansion

$$\epsilon^{(N-l)\binom{m}{2}+l\binom{m+1}{2}}(\tilde{E}(z) + o(1))$$

at $\epsilon = 0$. From the argument in the end of Section 4.1, \tilde{E} is shifted singular, and hence $\tilde{\xi} = \Phi^{-1}(\tilde{E})$ is singular. Hence the Specht module $\tilde{F} := \mathbb{C}\mathfrak{S}_n \cdot \tilde{\xi}$ of isotype (N^m, l) consists of singular polynomials. From the result of [3, 4] (see Section 6 of [4]), \tilde{F} has the basis $\{\xi_\mu \mid \mu \in \Lambda^{\text{rev}}(\lambda_0^+)\}$, where $\Lambda^{\text{rev}}(\lambda_0^+)$ is the set of reverse lattice permutations of λ_0^+ . Since λ_0 is minimal in $\Lambda^{\text{rev}}(\lambda_0^+)$ with respect to the dominance order and $\tilde{\xi}(x) = x^{\lambda_0} + (\text{lower order terms})$, we obtain $\tilde{\xi} = \xi_{\lambda_0}$. Sending the both hand sides by Φ , we get $\tilde{E} = E_{\lambda_0}$. ■

Acknowledgements

Research of SK is supported by Grant-in-Aid for Scientific Research (C) No. 19540228. Research of MN is supported by Grant-in-Aid for Young Scientists (B) No. 18749001. Research of YS is supported by Grant-in-Aid for Scientific Research (C) No. 20540009. Research of YT is supported by Grant-in-Aid for Young Scientists (B) No. 20740088.

The authors thank Takeshi Suzuki for helpful discussions. They are also deeply grateful to Charles Dunkl for valuable comments and pointing out an error in Section 4.2 of the earlier draft. YT thanks Takeshi Ikeda, Masahiro Kasatani and Hiroshi Naruse for discussions.

References

- [1] Cherednik I., Nonsymmetric Macdonald polynomials, *Int. Math. Res. Not.* **1995** (1995), no. 10, 483–515.
- [2] Cherednik I., Double affine Hecke algebras, Knizhnik–Zamolodchikov equations, and Macdonald’s operators, *Int. Math. Res. Not.* **1992** (1992), no. 9, 171–180.
- [3] Dunkl C.F., Singular polynomials for the symmetric groups, *Int. Math. Res. Not.* **2004** (2004), no. 67, 3607–3635, [math.RT/0403277](#).
- [4] Dunkl C.F., Singular polynomials and modules for the symmetric groups, *Int. Math. Res. Not.* **2005** (2005), no. 39, 2409–2436, [math.RT/0501494](#).
- [5] Dunkl C.F., de Jeu M.F.E., Opdam E.M., Singular polynomials for finite reflection groups, *Trans. Amer. Math. Soc.* **346** (1994), 237–256.
- [6] Di Francesco P., Zinn-Justin P., Quantum Knizhnik–Zamolodchikov equation, generalized Razumov–Stroganov sum rules and extended Joseph polynomials, *J. Phys. A: Math. Gen.* **38** (2005), L815–L822, [math-ph/0508059](#).
- [7] Forrester P., Log-gases and Random matrices, in progress.
- [8] Feigin B., Jimbo M., Miwa T., Mukhin E., A differential ideal of symmetric polynomials spanned by Jack polynomials at $\beta = -(r-1)/(k+1)$, *Int. Math. Res. Not.* **2002** (2002), no. 23, 1223–1237, [math.QA/0112127](#).
- [9] Frenkel I.B., Reshetikhin N.Yu., Quantum affine algebras and holonomic difference equations, *Comm. Math. Phys.* **146** (1992), 1–60.
- [10] Joseph A., On the variety of a highest weight module, *J. Algebra* **88** (1984), 238–278.
- [11] Knop F., Symmetric and non-symmetric quantum Capelli polynomials, *Comment. Math. Helv.* **72** (1997), 84–100, [q-alg/9603028](#).
- [12] Kasatani M., Takeyama Y., The quantum Knizhnik–Zamolodchikov equation and non-symmetric Macdonald polynomials, *Funkcial. Ekvac.* **50** (2007), 491–509, [math.QA/0608773](#).
- [13] Macdonald I.G., Affine Hecke algebras and orthogonal polynomials, *Astérisque* **237** (1996), 189–207.
- [14] Opdam E.M., Harmonic analysis for certain representations of graded Hecke algebras, *Acta Math.* **175** (1995), 75–121.
- [15] Suzuki T., Rational and trigonometric degeneration of the double affine Hecke algebra of type A, *Int. Math. Res. Not.* **2005** (2005), no. 37, 2249–2262, [math.RT/0502534](#).
- [16] Takeyama Y., Form factors of $SU(N)$ invariant Thirring model, *Publ. Res. Inst. Math. Sci.* **39** (2003), 59–116, [math-ph/0112025](#).