ISSN 1472-2739 (on-line) 1472-2747 (printed)

Algebraic & Geometric Topology
Volume 3 (2003) 905–920
Published: 29 September 2003

On the slice genus of links

Vincent Florens
Patrick M. Gilmer

Abstract We define Casson-Gordon σ-invariants for links and give a lower bound of the slice genus of a link in terms of these invariants. We study as an example a family of two component links of genus h and show that their slice genus is h, whereas the Murasugi-Tristram inequality does not obstruct this link from bounding an annulus in the 4-ball.

AMS Classification 57M25; 57M27

Keywords Casson-Gordon invariants, link signatures

1 Introduction

A knot in S^3 is slice if it bounds a smooth 2-disk in the 4-ball B^4. Levine showed [Le] that a slice knot is algebraically slice, i.e. any Seifert form of a slice knot is metabolic. In this case, the Tristram-Levine signatures at the prime power order roots of unity of a slice knot must be zero. Levine showed also that the converse holds in high odd dimensions, i.e. any algebraically slice knot is slice. This is false in dimension 3: Casson and Gordon [CG1, CG2, G] showed that certain two-bridge knots in S^3, which are algebraically slice, are not slice knots. For this purpose, they defined several knot and 3-manifold invariants, closely related to the Tristram-Levine signatures of associated links. Further methods to calculate these invariants were developed by Gilmer [Gi3, Gi4], Litherland [Li], Gilmer-Livingston [GL], and Naik [N]. Lines [L] also computed some of these invariants for some fibered knots, which are algebraically slice but not slice. The slice genus of a link is the minimal genus for a smooth oriented connected surface properly embedded in B^4 with boundary the given link.

The Murasugi-Tristram inequality (see Theorem 2.1 below) gives a lower bound on the slice genus of a link in terms of the link's Tristram-Levine signatures and related nullity invariants. The second author [Gi1] used Casson-Gordon invariants to give another lower bound on the slice genus of a knot. In particular he gave examples of algebraically slice knots whose slice genus is arbitrarily large. We apply these methods to restrict the slice genus of a link.

We study as an example a family of two component links, which have genus h Seifert surfaces. Using Theorem 4.1, we show that these links cannot bound a smoothly embedded surface in B^4 with genus lower than h, while the Murasugi-Tristram inequality does not show this. In fact there are some links with the same Seifert form that bound annuli in B^4. We work in the smooth category.

The second author was partially supported by NSF-DMS-0203486.

2 Preliminaries

2.1 The Tristram-Levine signatures

Let L be an oriented link in S^3, with μ components, and θ_S be the Seifert pairing corresponding to a connected Seifert surface S of the link. For any complex number λ with $|\lambda| = 1$, one considers the hermitian form $\theta_S^\lambda := (1 - \lambda)\theta_S + (1 - \overline{\lambda})(\theta_S)^T$. The Tristram signature $\sigma_L(\lambda)$ and nullity $n_L(\lambda)$ of L are defined as the signature and nullity of θ_S^λ. Levine defined these same signatures for knots [Le]. The Alexander polynomial of L is $\Delta_L(t) := \mathrm{Det}(\theta_S - t(\theta_S)^T)$. As is well-known, σ_L is a locally constant map on the complement in S^1 of the roots of Δ_L and n_L is zero on this complement. If $\Delta_L = 0$, it is still true that the signature and nullity are locally constant functions on the complement of some finite collection of points.

The Murasugi-Tristram inequality allows one to estimate the slice genus of L, in terms of the values of $\sigma_L(\lambda)$ and $n_L(\lambda)$.

Theorem 2.1 [M, T] *Suppose that L is the boundary of a properly embedded connected oriented surface F of genus g in B^4. Then, if λ is a prime power order root of unity, we have*

$$|\sigma_L(\lambda)| + n_L(\lambda) \leq 2g + \mu - 1.$$

2.2 The Casson-Gordon σ-invariant

In this section, for the reader convenience, we review the definition and some of the properties of the simplest kind of Casson-Gordon invariant. It is a reformulation of the Atiyah-Singer α-invariant.

Let M be an oriented compact three manifold and $\chi\colon H_1(M) \to \mathbb{C}^*$ be a character of finite order. For some $q \in \mathbb{N}^*$, the image of χ is contained a cyclic subgroup of order q generated by $\alpha = e^{2i\pi/q}$. As $\mathrm{Hom}(H_1(M), C_q) = [M, B(C_q)]$, it follows that χ induces q-fold covering of M, denoted $\widetilde{M}$, with a canonical deck transformation. We will denote this transformation also by α. If χ maps onto C_q, the canonical deck transformation sends x to the other endpoint of the arc

that begins at x and covers a loop representing an element of $(\chi)^{-1}(\alpha)$.

As the bordism group $\Omega_3(B(C_q)) = C_q$, we may conclude that n disjoint copies of M , for some integer n, bounds bound a compact 4-manifold W over $B(C_q)$. Note n can be taken to be q. Let $\widetilde{W}$ be the induced covering with the deck transformation, denoted also by α, that restricts to α on the boundary. This induces a $\mathbb{Z}[C_q]$- module structure on $C_*(\widetilde{W})$, where the multiplication by $\alpha \in \mathbb{Z}[C_q]$ corresponds to the action of α on $\widetilde{W}$.

The cyclotomic field $\mathbb{Q}(C_q)$ is a natural $\mathbb{Z}[C_q]$-module and the twisted homology $H_*^t(W; \mathbb{Q}(C_q))$ is defined as the homology of

$$C_*(\widetilde{W}) \otimes_{\mathbb{Z}[C_q]} \mathbb{Q}(C_q).$$

Since $\mathbb{Q}(C_q)$ is flat over $\mathbb{Z}[C_q]$, we get an isomorphism

$$H_*^t(W; \mathbb{Q}(C_q)) \simeq H_*(\widetilde{W}) \otimes_{\mathbb{Z}[C_q]} \mathbb{Q}(C_q).$$

Similarly, the twisted homology $H_*^t(M; \mathbb{Q}(C_q))$ is defined as the homology of

$$C_*(\widetilde{M}) \otimes_{\mathbb{Z}[C_q]} \mathbb{Q}(C_q).$$

Let $\widetilde{\phi}$ be the intersection form on $H_2(\widetilde{W}; \mathbb{Q})$ and define

$$\phi_\chi(W)\colon\ H_2^t(W; \mathbb{Q}(C_q)) \times H_2^t(W; \mathbb{Q}(C_q)) \to \mathbb{Q}(C_q)$$

so that, for all a, b in $\mathbb{Q}(C_q)$ and x, y in $H_2(\widetilde{W})$,

$$\phi_\chi(W)(x \otimes a, y \otimes b) = \overline{a} b \sum_{i=1}^{q} \widetilde{\phi}(x, \alpha^i y) \overline{\alpha}^i,$$

where $a \to \bar{a}$ denotes the involution on $\mathbb{Q}(C_q)$ induced by complex conjugation.

Definition 2.2 The Casson-Gordon σ-invariant of (M, χ) and the related nullity are

$$\sigma(M, \chi) := \frac{1}{n}\big(\mathrm{Sign}(\phi_\chi(W)) - \mathrm{Sign}(W)\big)$$

$$\eta(M, \chi) := \dim\, H_1^t(M; \mathbb{Q}(C_q)).$$

If U is a closed 4-manifold and $\chi\colon H_1(U) \to C_q$ we may define $\phi_\chi(U)$ as above. One has that modulo torsion the bordism group $\Omega_4(B(C_q))$ is generated by the constant map from $CP(2)$ to $B(C_q)$. If χ is trivial, one has that $\mathrm{Sign}(\phi_\chi(U)) = \mathrm{Sign}(U)$. Since both signatures are invariant under cobordism, one has in general that $\mathrm{Sign}(\phi_\chi(U)) = \mathrm{Sign}(U)$. The independence of $\sigma(M,\chi)$ from the choice of W and n follows from this and Novikov additivity. One may see directly that these invariants do not depend on the choice of q. In this way Casson and Gordon argued that $\sigma(M,\chi)$ is an invariant. Alternatively one may use the Atiyah-Singer G-Signature theorem and Novikov additivity [AS].

We now describe a way to compute $\sigma(M,\chi)$ for a given surgery presentation of (M,χ).

Definition 2.3 Let K be an oriented knot in S^3. Let A be an embedded annulus such that $\partial A = K \cup K'$ with $lk(K,K') = f$. A *p-cable on K with twist f* is defined to be the union of oriented parallel copies of K lying in A such that the number of copies with the same orientation minus the number with opposite orientation is equal to p.

Let us suppose that M is obtained by surgery on a framed link $L = L_1 \cup \cdots \cup L_\mu$ with framings $f_1, \dots, f_\mu$. One shows that the linking matrix Λ of L with framings in the diagonal is a presentation matrix of $H_1(M)$ and a character on $H_1(M)$ is determined by $\alpha^{p_i} = \chi(m_{L_i}) \in C_q$ where m_{L_i} denotes the class of the meridian of L_i. Let $\vec{p} = (p_1, \dots, p_\mu)$. We use the following generalization of a formula in [CG2, Lemma (3.1)], where all p_i are assumed to be 1, that is given in [Gi2, Theorem(3.6)].

Proposition 2.4 *Suppose χ maps onto C_q. Let L' with μ' components be the link obtained from L by replacing each component by a non-empty algebraic p_i-cable with twist f_i along this component. Then, if $\lambda = e^{2ir\pi/q}$, for $(r,q) = 1$, one has*

$$\sigma(M,\chi^r) = \sigma_{L'}(\lambda) - \mathrm{Sign}(\Lambda) + 2\frac{r(q-r)}{q^2}\vec{p}^\top \Lambda \vec{p},$$

$$\eta(M,\chi^r) = \eta_{L'}(\lambda) - \mu' + \mu.$$

The following proposition collects some easy additivity properties of the σ-invariant and the nullity under the connected sum.

Proposition 2.5 *Suppose that M_1, M_2 are connected. Then, for all $\chi_i \in H^1(M_i; C_q)$, $i = 1,2$, we have*

$$\sigma(M_1 \# M_2, \chi_1 \oplus \chi_2) = \sigma(M_1,\chi_1) + \sigma(M_2,\chi_2).$$

If both χ_i are non-trivial, then

$$\eta(M_1 \# M_2, \chi_1 \oplus \chi_2) = \eta(M_1, \chi_1) + \eta(M_2, \chi_2) + 1.$$

If one χ_i is trivial, then

$$\eta(M_1 \# M_2, \chi_1 \oplus \chi_2) = \eta(M_1, \chi_1) + \eta(M_2, \chi_2).$$

Proposition 2.6 *For all $\chi \in H_1(S^1 \times S^2; C_q)$, we have*

$$\sigma(S^1 \times S^2, \chi) = 0$$

If $\chi \neq 0$, then $\eta(S^1 \times S^2, \chi) = 0$. If $\chi = 0$, then $\eta(S^1 \times S^2, \chi) = 1$.

Proposition 2.6 for non-trivial χ can be proved for example by the use of Proposition 2.4, since $S^1 \times S^2$ is obtained by surgery on the unknot framed 0. However it is simplest to derive this result directly from the definitions.

2.3 The Casson-Gordon τ-invariant

In this section, we recall the definition and some of the properties of the Casson-Gordon τ-invariant. Let C_∞ denote a multiplicative infinite cyclic group generated by t. For $\chi^+ \colon H_1(M) \to C_q \oplus C_\infty$, we denote $\bar{\chi} \colon H_1(M) \to C_q$ the character obtained by composing χ^+ with projection on the first factor. The character χ^+ induces a $C_q \times C_\infty$-covering $\widetilde{M}_\infty$ of M.

Since the bordism group $\Omega_3(B(C_q \times C_\infty)) = C_q$, bounds a compact 4-manifold W over $B(C_q \times C_\infty)$ Again n can be taken from to be q.

If we identify $\mathbb{Z}[C_q \times C_\infty]$ with the Laurent polynomial ring $\mathbb{Z}[C_q][t, t^{-1}]$, the field $\mathbb{Q}(C_q)(t)$ of rational functions over the cyclotomic field $\mathbb{Q}(C_q)$ is a flat $\mathbb{Z}[C_q \times C_\infty]$-module. We consider the chain complex $C_*(\widetilde{W}_\infty)$ as a $\mathbb{Z}[C_q \times C_\infty]$-module given by the deck transformation of the covering. Since W is compact, the vector space $H_2^t(W; \mathbb{Q}(C_q)(t)) \simeq H_2(\widetilde{W}_\infty) \otimes_{\mathbb{Z}[C_q][t,t^{-1}]} \mathbb{Q}(C_q)(t)$ is finite dimensional.

We let J denote the involution on $\mathbb{Q}(C_q)(t)$ that is linear over $\mathbb{Q}$ sends t^i to t^{-i} and α^i to α^{-i}. As in [G], one defines a hermitian form, with respect to J,

$$\phi_{\chi^+} \colon H_2^t(W; \mathbb{Q}(C_q)(t)) \times H_2^t(W; \mathbb{Q}(C_q)(t)) \to \mathbb{Q}(C_q)(t),$$

such that

$$\phi_{\chi^+}(x \otimes a, y \otimes b) = J(a) \cdot b \cdot \sum_{i \in \mathbb{Z}} \sum_{j=1}^{q} \widetilde{\phi^+}(x, t^i \alpha^j y) \overline{\alpha}^j t^{-i}.$$

Here $\widetilde{\phi^+}$ denotes the ordinary intersection form on $\widetilde{W}_\infty$. Let $\mathcal{W}(\mathbb{Q}(C_q)(t))$ be the Witt group of non-singular hermitian forms on finite dimensional $\mathbb{Q}(C_q)(t)$ vector spaces. Let us consider $H_2^t(W;\mathbb{Q}(C_q)(t))/(\mathrm{Radical}(\phi_{\chi^+}))$. The induced form on it represents an element in $\mathcal{W}$ $(\mathbb{Q}(C_q)(t))$, which we denote $w(W)$. Furthermore, the ordinary intersection form on $H_2(W;\mathbb{Q})$ represents an element of $\mathcal{W}(\mathbb{Q})$. Let $w_0(W)$ be the image of this element in $\mathcal{W}(\mathbb{Q}(C_q)(t))$.

Definition 2.7 The Casson-Gordon τ-invariant of (M,χ^+) is

$$\tau(M,\chi^+) := \frac{1}{n}\big(w(W) - w_0(W)\big) \in \mathcal{W}(\mathbb{Q}(C_q)(t)) \otimes \mathbb{Q}.$$

Suppose that nM bounds another compact 4-manifold W' over $B(C_q \times C_\infty)$. Form the closed compact manifold over $B(C_q \times C_\infty)$, $U := W \cup W'$ by gluing along the boundary. By Novikov additivity, we get $w(U) - w_0(U) = \big(w(W) - w_0(W)\big) - \big(w(W') - w_0(W')\big)$. Using [CF], the bordism group $\Omega_4(B(C_q \times C_\infty))$, modulo torsion, is generated by $CP(2)$, with the constant map to $B(C_q \times C_\infty)$. We have that $w(CP(2)) = w_0(CP(2))$. Since $w(U)$, and $w_0(U)$ only depend on the bordism class of U over $B(C_q \times C_\infty)$, it follows that $w(U) = w_0(U)$ and $\tau(M,\chi^+)$ is independent of the choice of W. Using the above techniques, one may check $\tau(M,\chi^+)$ is independent of n.

If $A \in \mathcal{W}(\mathbb{Q}(C_q)(t))$, let $A(t)$ be a matrix representative for A. The entries of $A(t)$ are Laurent polynomials with coefficients in $\mathbb{Q}(C_q)$. If λ is in $S^1 \subset \mathbb{C}$, then $A(\lambda)$ is hermitian and has a well defined signature $\sigma_\lambda(A)$. One can view $\sigma_\lambda(A)$ as a locally constant map on the complement of the set of the zeros of $\det A(\lambda)$. As in [CG1], we re-define $\sigma_\lambda(A)$ at each point of discontinuity as the average of the one-sided limits at the point.

We have the following estimate [Gi3, Equation (3.1)].

Proposition 2.8 *Let $\chi^+\colon H_1(M) \to C_q \oplus C_\infty$ and $\bar{\chi}\colon H_1(M) \to C_q$ be χ^+ followed by the projection to C_q. We have*

$$|\sigma_1\big(\tau(M,\chi^+)\big) - \sigma(M,\bar{\chi})| \leq \eta(M,\bar{\chi}).$$

2.4 Linking forms

Let M be a rational homology 3-sphere with linking form

$$l\colon H_1(M) \times H_1(M) \to \mathbb{Q}/\mathbb{Z}.$$

We have that l is non-singular, that is the adjoint of l is an isomorphism $\iota\colon H_1(M) \to \mathrm{Hom}(H_1(M),\mathbb{Q}/\mathbb{Z})$. Let $H_1(M)^*$ denote $\mathrm{Hom}(H_1(M),\mathbb{C}^*)$. Let

ν denote the map $\mathbb{Q}/\mathbb{Z} \to \mathbb{C}^*$ that sends $\frac{a}{b}$ to $e^{\frac{2\pi i a}{b}}$. So we have an isomorphism $\jmath\colon H_1(M) \to H_1(M)^*$ given by $x \mapsto \nu \circ \iota(x)$. Let $\beta\colon H_1(M)^* \times H_1(M)^* \to \mathbb{Q}/\mathbb{Z}$ be the dual form defined by $\beta(\jmath x, \jmath y) = -l(x,y)$.

Definition 2.9 The form β is metabolic with metabolizer H if there exists a subgroup H of $H_1(M)^*$ such that $H^\perp = H$.

Lemma 2.10 [Gi1] *If M bounds a spin 4-manifold W then $\beta = \beta_1 \oplus \beta_2$ where β_2 is metabolic and β_1 has an even presentation with rank dim $H_2(W;\mathbb{Q})$ and signature Sign(W). Moreover, the set of characters that extend to $H_1(W)$ forms a metabolizer for β_2.*

2.5 Link invariants

Let $L = L_1 \cup \cdots \cup L_\mu$ be an oriented link in S^3. Let N_2 be the two-fold covering of S^3 branched along L and β_L be the linking form on $H_1(N_2)^*$, see previous section.

We suppose that the Alexander polynomial of L satisfies

$$\Delta_L(-1) \neq 0.$$

Hence, N_2 is a rational homology sphere. Note that if $\Delta_L(-1) \neq 1$, then $H_1(N_2;\mathbb{Z})$ is non-trivial.

Definition 2.11 For all characters χ in $H_1(N_2)^*$, the Casson-Gordon σ-invariant of L and the related nullity are (see Definition 2.2):

$$\sigma(L,\chi) := \sigma(N_2,\chi),$$

$$\eta(L,\chi) := \eta(N_2,\chi).$$

Remark 2.12 If L is a knot, then Definition 2.11 coincides with $\sigma(L,\chi)$ defined in [CG1, p.183].

3 Framed link descriptions

In this section, we study the Casson-Gordon τ-invariants of the two-fold cover M_2 of the manifold M_0 described below.

Let $S^3 - T(L)$ be the complement in S^3 of an open tubular neighborhood of L in S^3 and P be a planar surface with μ boundary components.

Let S be a Seifert surface for L and γ_i for $i = 1, \dots, \mu$ be the curves where S intersects the boundary of $S^3 - T(L)$. We define M_0 as the result of gluing $P \times S^1$ to $S^3 - T(L)$, where $P \times 1$ is glued along the curves γ_i. Let $*$ be a point in the boundary of P.

A recipe for drawing a framed link description for M_0 is given in the proof of Proposition 3.1.

Proposition 3.1

$$H_1(M_0) \simeq \mathbb{Z} \oplus \mathbb{Z}^{\mu-1} \simeq \langle m \rangle \oplus \ \mathbb{Z}^{\mu-1},$$

where m denotes the class of $* \times S^1$ in $P \times S^1$.

Proof Form a 4-manifold X by gluing $P \times D^2$ to D^4 along S^3 in such a way that the total framing on L agrees with the Seifert surface S. The boundary of this 4-manifold is M_0. We can get a surgery description of M_0 in the following way: pick $\mu - 1$ paths of S joining up the components of L in a chain. Deleting open neighborhoods of these paths in S gives a Seifert surface for a knot L' obtained by doing a fusion of L along bands that are neighborhoods of the original paths. Put a circle with a dot around each of these bands (representing a 4-dimensional 1-handle in Kirby's [K] notation), and the framing zero on L'. This describes a handlebody decomposition of X.

One can then get a standard framed link description of M_0 by replacing the circle with dots with unknots $T_1, \dots, T_{\mu-1}$ framed zero. This changes the 4-manifold but not the boundary. Note also that $lk(T_i, T_j) = 0$ and $lk(T_i, L') = 0$ for all $i = 1, \dots, \mu - 1$. Hence $H_1(M_0) \simeq \mathbb{Z}^\mu$ and m represents one of the generators. □

We now consider an infinite cyclic covering M_∞ of M_0, defined by a character $H_1(M_0) \to C_\infty = \langle t \rangle$ that sends m to t and the other generators to zero. Let us denote by M_2 the intermediate two-fold covering obtained by composing this character with the quotient map $C_\infty \to C_2$ sending t to -1. Let m_2 denote the loop in M_2 given by the inverse image of m. A recipe for drawing a framed link description for M_2 is given in the proof of Remark 3.3.

Proposition 3.2 *There is an isomorphism between $H_1(N_2)$ and the torsion subgroup of $H_1(M_2)$, which only depends on L. Moreover*

$$H_1(M_2) \simeq H_1(N_2) \oplus \mathbb{Z}^\mu \simeq H_1(N_2) \oplus \langle m_2 \rangle \oplus \mathbb{Z}^{\mu-1}.$$

Proof Let R be the result of gluing $P \times D^2$ to $S^3 \times I$ along $L \times 1 \subset S^3 \times 1$ using the framing given by the Seifert surface. Thus R is the result of adding $\mu - 1$ 1-handles to $S^3 \times I$ and then one 2-handle along L', as in the proof above. Then X in the proof above can be obtained by gluing D^4 to R along $S^3 \times 0$. Since D^2 is the double branched cover of itself along the origin, $P \times D^2$ is the double branched cover of itself along $P \times 0$. Let R_2 denote the double branched cover of R that is obtained by gluing $P \times D^2$ to $N_2 \times I$ along a neighborhood of the lift of $L \times 1 \subset S^3 \times 1$. We have that $\partial R_2 = -N_2 \sqcup M_2$, where R_2 is the result of adding $\mu - 1$ 1-handles to $N_2 \times I$ and then one 2-handle along the lift L'. Moreover this lift of L' is null-homologous in N_2. It follows that $H_1(R_2)$ is isomorphic to $H_1(N_2) \oplus \mathbb{Z}^{\mu-1}$, with the inclusion of N_2 into R_2 inducing an isomorphism i_N of $H_1(N_2)$ to the torsion subgroup of $H_1(R_2)$. Turning this handle decomposition upside down we have that R_2 is the result of adding to $M_2 \times I$ one 2-handle along a neighborhood of m_2 and then $\mu - 1$ 3-handles. It follows that $H_1(R_2) \oplus \mathbb{Z} = H_1(R_2) \oplus \langle m_2 \rangle$ is isomorphic to $H_1(M_2)$ with the inclusion of M_2 in R_2 inducing an isomorphism i_M of the torsion subgroup $H_1(M_2)$ to the torsion subgroup of $H_1(R_2)$. Thus $(i_M)^{-1} \circ i_N$ is an isomorphism from $H_1(N_2)$ to the torsion subgroup of $H_1(M_2)$ and this isomorphism is constructed without any arbitrary choices. □

Remark 3.3 We could have proved Proposition 3.1 in a similar way to the proof of Proposition 3.2. We could have also proved Proposition 3.2 (except for the isomorphism only depending on L) in a similar way to the proof of Proposition 3.1 as follows. We can find a surgery description of M_2 from a surgery description of N_2. The procedure of how to visualize a lift of L and the surface S in N_2 is given in [AK]. One considers the lifts of the paths chosen in the proof of Proposition 3.1, on the lift of S. One then fuses the components of the lift of L along these paths, obtaining a lift of L'. The surgery description of M_2 is obtained by adding to the surgery description of N_2 the lift of L' with zero framing together with $\mu - 1$ more unknotted zero-framed components encircling each fusion. The linking matrix of this link is a direct sum of that of N_2 and a $\mu \times \mu$ zero matrix.

Let i_T denote the inclusion of the torsion subgroup of $H_1(M_2)$ into $H_1(M_2)$, and let $\psi\colon H_1(N_2) \to H_1(M_2)$ denote the monomorphism given by $i_T \circ (i_M)^{-1} \circ i_N$.

Theorem 3.4 *Let $\chi^+\colon H_1(M_2) \to C_q \oplus C_\infty$. Let $\chi\colon H_1(N_2) \to C_q$ be $\chi^+ \circ \psi$ composed with the projection to C_q. We have that:*

$$|\sigma_1(\tau(M_2,\chi^+)) - \sigma(L,\chi)| \leq \eta(L,\chi) + \mu.$$

Remark 3.5 If L is a knot, then $\tau(M_2,\chi^+)$ coincides with $\tau(L,\chi)$ defined in [CG1, p.189].

Proof of Theorem 3.4 We use the surgery description of M_2 given in Remark 3.3. Let P be given by the surgery description of M_2 but with the component corresponding to L' deleted. Hence,

$$P = N_2 \sharp_{(\mu-1)} S^1 \times S^2.$$

$\chi+$ induces some character χ' on $H_1(P)$.

According to Section 2.3, we let $\overline{\chi} \in H^1(M_2;C_q)$ and $\overline{\chi}' \in H^1(P;C_q)$ denote the characters χ^+ and χ' followed by the projection $C_q \oplus C_\infty \to C_q$. Using Propositions 2.5 and 2.6, one has that

$$\sigma(P,\overline{\chi}') = \sigma(L,\chi) \text{ and } \eta(P,\overline{\chi}') = \eta(L,\chi) + \mu - 1.$$

Moreover, since M_2 is obtained by surgery on L' in P, it follows from [Gi3, Proposition (3.3)] that

$$|\sigma(P,\overline{\chi}') - \sigma(M_2,\overline{\chi})| + |\eta(M_2,\overline{\chi}) - \eta(P,\overline{\chi}')| \leq 1 \text{ or}$$

$$|\sigma(L,\chi) - \sigma(M_2,\overline{\chi})| + |\eta(M_2,\overline{\chi}) - \eta(L,\chi) - \mu + 1| \leq 1.$$

Thus

$$|\sigma(L,\chi) - \sigma(M_2,\overline{\chi})| \leq \eta(L,\chi) + \mu - \eta(M_2,\overline{\chi}).$$

Finally, one gets, by Theorem 2.8,

$$\begin{aligned}|\sigma_1(\tau(M_2,\chi^+)) - \sigma(L,\chi)| &\leq |\sigma_1(\tau(M_2,\chi^+)) - \sigma(M_2,\overline{\chi})| + |\sigma(M_2,\overline{\chi}) - \sigma(L,\chi)| \\ &\leq \eta(M_2,\overline{\chi}) + \eta(L,\chi) + \mu - \eta(M_2,\overline{\chi}) = \eta(L,\chi) + \mu.\end{aligned}$$

□

4 The slice genus of links

See Section 2.5 for notations.

Theorem 4.1 *Suppose L is the boundary of a connected oriented properly embedded surface F of genus g in B^4, and that $\Delta_L(-1) \neq 0$. Then, β_L can be written as a direct sum $\beta_1 \oplus \beta_2$ such that the following two conditions hold:*

1) *β_1 has an even presentation of rank $2g+\mu-1$ and signature $\sigma_L(-1)$, and β_2 is metabolic.*

2) *There is a metabolizer for β_2 such that for all characters χ of prime power order in this metabolizer,*

$$|\sigma(L,\chi)+\sigma_L(-1)| \leq \eta(L,\chi)+4g+3\mu-2.$$

Proof We let $b_i(X)$ denote the ith Betti number of a space X. We have $b_1(F)=2g+\mu-1$.

Let W_0', with boundary M_0', be the complement of an open tubular neighborhood of F in B^4. By the Thom isomorphism, excision, and the long exact sequence of the pair (B^4, W_0'), W_0' has the homology of S^1 wedge $b_1(F)$ 2-spheres. Let W_2' with boundary M_2' be the two-fold covering of W_0'. Note that if F is planar, $M_0'=M_0$, and $M_2'=M_2$ (see Section 3).

Let V_2 be the two-fold covering of B^4 with branched set F. Note that V_2 is spin as $w_2(V_2)$ is the pull-up of a class in $H^2(B^4,\mathbb{Z}_2)$, by [Gi5, Theorem 7], for instance. The boundary of V_2 is N_2. As in [Gi1], one calculates that $b_2(V_2)=2g+\mu-1$. One has $\mathrm{Sign}(V_2)=\sigma_L(-1)$ by [V].

By Lemma 2.10, β_L can be written as a direct sum $\beta_1\oplus\beta_2$ as in condition 1) above, such that the characters on $H_1(N_2)$ that extend to $H_1(V_2)$ form a metabolizer H for β_2. We now suppose $\chi\in H$ and show that Condition 2) holds for χ.

We also let χ denote an extension of χ to $H_1(V_2)$ with image some cyclic group C_q where q is a power of a prime integer (possibly larger than those corresponding to the character on $H_1(N_2)$). Of course $\chi\in H^1(V_2,C_q)$ restricted to W_2' extends χ restricted to M_2'. We simply denote all these restrictions by χ.

Let W_∞' denote the infinite cyclic cover of W_0'. Note that W_2' is a quotient of this covering space. χ induces a C_q-covering of V_2 and thus of W_2'. If we pull the C_q-covering of W_2' up to W_∞', we obtain $\widetilde{W}_\infty'$, a $C_q\times C_\infty$-covering of W_2'. If we identify properly $F\times S^1$ in M_2', this covering restricted to $F\times S^1$ is given by a character $H_1(F\times S^1)\simeq H_1(F)\oplus H_1(S^1)\to C_q\times C_\infty$ that maps $H_1(F)$ to zero in C_∞, $H_1(S^1)$ to zero in C_q and isomorphically onto C_∞. For this note: since $\mathrm{Hom}(H_1(F),\mathbb{Z})=H^1(F)=[F,S^1]$, we may define diffeomorphisms of $F\times S^1$ that induce the identity on the second factor of $H_1(F\times S^1)\approx H_1(F)\oplus\mathbb{Z}$, and send $(x,0)\in H_1(F)\oplus\mathbb{Z}$, to $(x,f(x))\in H_1(F)\oplus\mathbb{Z}$, for any $f\in\mathrm{Hom}(H_1(F),\mathbb{Z})$.

As in [Gi1], choose inductively a collection of g disjoint curves in the kernel of χ that form a metabolizer for the intersection form on $H_1(F)/H_1(\partial F)$. By taking a tubular neighborhood of these curves in F, we obtain a collection of $S^1\times I$ embedded in F. Using these embeddings we can attach round 2-handles $(B^2\times I)\times S^1$ along $(S^1\times I)\times S^1$ to the trivial cobordism $M_2'\times I$ and obtain a cobordism Ω between M_2 and M_2'.

Let $U=W_2'\cup_{M_2'}\Omega$ with boundary M_2. The $C_q\times C_\infty$-covering of W_2' extends uniquely to U. Note that Ω may also be viewed as the result of attaching round 1-handles to $M_2\times I$.

As in [Gi1], $\mathrm{Sign}(W_2')=\mathrm{Sign}(V_2)$. Since the intersection form on Ω is zero, we get $\mathrm{Sign}(U)=\mathrm{Sign}(W_2')=\mathrm{Sign}(V_2)=\sigma_L(-1)$. The $C_q\times C_\infty$-covering of Ω, restricted to each round 2-handle is q copies of $B^2\times I\times\mathbb{R}$ attached to the trivial cobordism $\widetilde{M}_\infty'\times I$ along q copies of $S^1\times I\times\mathbb{R}$. Using a Mayer-Vietoris sequence, one sees that the inclusion induces an isomorphism (which preserves the Hermitian form)

$$H_2^t(U;\mathbb{Q}(C_q)(t))\simeq H_2^t(W_2';\mathbb{Q}(C_q)(t)).$$

Thus, if $w(W_2')$ denotes the image of the intersection form on $H_2^t(W_2';\mathbb{Q}(C_q)(t))$ in $\mathcal{W}\ (\mathbb{Q}(C_q)(t))$, we get $\sigma_1(\tau(M_2,\chi^+))=\sigma_1(w(W_2'))-\sigma_L(-1)$.

If q is a prime power, we may apply Lemma 2 of [Gi1] and conclude that $H_i(\widetilde{W}_\infty';\mathbb{Q})$ is finite dimensional for all $i\neq 2$. Thus, $H_i^t(W_2';\mathbb{Q}(C_q)(t))$ is zero for all $i\neq 2$. Since the Euler characteristic of W_2' with coefficients in $\mathbb{Q}(C_q)(t)$ coincides with those with coefficients in $\mathbb{Q}$, we get $\dim H_2^t(W_2';\mathbb{Q}(C_q)(t))=\chi(W_2')=2\chi(W_0')=2(1-\chi(F))=2b_1(F)$. Thus $|\sigma_1(\tau(M_2,\chi^+)+\sigma_L(-1)|\leq 2b_1(F)$. Hence,

$$\begin{aligned}&|\sigma(L,\chi)+\sigma_L(-1)|\leq|\sigma(L,\chi)-\sigma_1(\tau(M_2,\chi)^+)|+|\sigma_1(\tau(M_2,\chi)^+)+\sigma_L(-1)|\\&\leq\eta(L,\chi)+\mu+2(2g+\mu-1)=\eta(L,\chi)+4g+3\mu-2 \text{ by Theorem 3.4.}\end{aligned}$$ □

5 Examples

Let $L=L_1\cup L_2$ be the link with two components of Figure 1 and S be the Seifert surface of L given by the picture. The squares with K denote two parallel copies with linking number 0 of an arc tied in the knot K. Note that L is actually a family of examples. Specific links are determined by the choice of two parameters: a knot K and a positive integer h. Since S has genus h, the slice genus of L is at most h.



Figure 1: The link L

One calculates that $\sigma_L(\lambda) = 1$, and $n_L(\lambda) = 0$ for all λ. Thus, the Murasugi-Tristram inequality says nothing about the slice genus of L. In fact, if K is a slice knot, then one can surger this surface to obtain a smooth cylinder in the 4-ball with boundary L. Thus there can be no arguments based solely on a Seifert pairing for L that would imply that the slice genus is non-zero.

Theorem 5.1 *If $\sigma_K(e^{2i\pi/3}) \geq 2h$ or $\sigma_K(e^{2i\pi/3}) \leq -2h-2$, then L has slice genus h.*

Proof Using [AK], a surgery presentation of N_2 as surgery on a framed link of $2h+1$ components can be obtained from the surface S (see Figure 2).

Let Q be the 3-manifold obtained from the link pictured in Figure 2. Here K' denotes K with the string orientation reversed. Since $RP(3)$ is obtained by surgery on the unknot framed 2, we get:

$$N_2 = RP(3) \#_h Q.$$

The linking matrix of the framed link of the surgery presentation of N_2 is

$\Lambda = [2] \bigoplus \oplus^h \begin{bmatrix} 0 & 3 \\ 3 & 0 \end{bmatrix}$. Λ is a presentation matrix of $(H_1(N_2)^*, \beta_L)$; we obtain

$$H_1(N_2)^* \simeq \mathbb{Z}_2 \bigoplus \oplus^{2h} \mathbb{Z}_3$$



Figure 2: Surgery presentation of Q

and β_L is given by the following matrix, with entries in $\mathbb{Q}/\mathbb{Z}$:

$$[1/2] \bigoplus \oplus^h \begin{bmatrix} 0 & 1/3 \\ 1/3 & 0 \end{bmatrix}.$$

By Theorem 4.1, if L bounds a surface of genus $h-1$ in B^4, then β_L must be decomposed as $\beta_1 \oplus \beta_2$ where:

1) β_1 has an even presentation matrix of rank $2h-1$, and signature 1 (all we really need here is that it has a rank $2h-1$ presentation.)

2) β_2 is metabolic and for all characters χ of prime power order in some metabolizer of β_2, the following inequality holds:

$$(*) \qquad |\sigma(L,\chi)+1| - \eta(L,\chi) \leq 4h.$$

As $\mathbb{Z}_2 \bigoplus \oplus^{2h}\mathbb{Z}_3$ does not have a rank $2h-1$ presentation, β_2 is non-trivial. As metabolic forms are defined on groups whose cardinality is a square, β_2 is defined on a group with no 2-torsion. Thus the metabolizer contains a non-trivial character of order three satisfying $\beta_L(\chi,\chi)=0$.

The first homology of Q is $\mathbb{Z}_3 \oplus \mathbb{Z}_3$, generated by, say, m_1 and m_2, positive meridians of these components. Each of these components is oriented counter-clockwise. We first work out $\sigma(Q,\chi)$ and $\eta(Q,\chi)$ for characters of order three. Let $\chi_{(a_1,a_2)}$ denote the character on $H_1(Q)$ sending m_j to $e^{\frac{2i\pi a_j}{3}}$, where the a_j take the values zero and ± 1.

We use Proposition 2.4 to compute $\sigma(Q,\chi_{(1,0)})$ and $\eta(Q,\chi_{(1,0)})$ assuming that K is trivial. For this, one may adapt the trick illustrated on a link with 2 twists between the components [Gi2, Fig (3.3), Remark (3.65b)]. In the case K is the unknot, we obtain

$$\sigma(Q,\chi_{(1,0)}) = 1 \quad \text{and} \quad \eta(Q,\chi_{(1,0)}) = 0.$$

It is not difficult to see that inserting the knots of the type K changes the result as follows (note that K and K' have the same Tristram-Levine signatures):

$$\sigma(Q,\chi_{(1,0)}) = 1 + 2\sigma_K(e^{2\pi i/3}) \quad \text{and} \quad \eta(Q,\chi_{(1,0)}) = 0.$$

These same values hold for the characters $\chi_{(-1,0)}$ and $\chi_{(0,\pm1)}$ by symmetry.

Using Proposition 2.4

$$\sigma(Q,\pm\chi_{(1,1)}) = -1 - 24/9 + 4\sigma_K(e^{2\pi i/3}), \qquad \eta(Q,\pm\chi_{(1,1)}) = 0$$

$$\sigma(Q,\pm\chi_{(1,-1)}) = 4 + 24/9 + 4\sigma_K(e^{2\pi i/3}) \quad \text{and} \quad \eta(Q,\pm\chi_{(1,-1)}) = 1.$$

One also has

$$\sigma(Q,\chi_{(0,0)}) = 0 \quad \text{and} \quad \eta(Q,\chi_{(0,0)}) = 0.$$

Any order three character on N_2 that is self annihilating under the linking form is given as the sum of the trivial character on $RP(3)$ and characters of type $\chi_{(0,0)}$, $\chi_{(\pm1,0)}$ and $\chi_{(0,\pm1)}$ on Q and characters of type $\pm\chi_{(1,1)} + \pm\chi_{(1,-1)}$ on $Q\#Q$. Using Proposition 2.5, one can calculate $\sigma(L,\chi)$ and $\eta(L,\chi)$ for all these characters χ. It is now a trivial matter to check that for every non-trivial character with $\beta(\chi,\chi) = 0$, the inequality (*) is not satisfied. □

References

[AK] Akbulut, S., Kirby, R., *Branched covers of surfaces in 4-manifolds*, Math. Ann. **252**, 111-131 (1980).

[AS] Atiyah, M. F., Singer, I. M., *The index of elliptic operators. III*, Ann. of Math. **(2) 87**, 546-604 (1968).

[CF] Conner, P. E., Floyd, E.E., *Differential Periodic Maps*, Ergebnisse der Mathematik und ihrer Grenzgebiete, **33**, Springer-Verlag, (1964).

[CG1] Casson, A. J., Gordon, C. Mc A., *Cobordism of classical knots*, Progr. Math., **62**, A La Recherche de la Topologie Perdue, Birkhauser, Boston, MA, 181–199 (1986).

[CG2] Casson, A. J., Gordon, C. Mc A., *On slice knots in dimension three*, Proc. Symp. in Pure Math. XXX, **2**, 39-53 (1978).

[Gi1] Gilmer, P. M., *On the slice genus of knots*, Invent. Math. **66**, 191-197 (1982).

[Gi2] Gilmer, P. M., *Configurations of surfaces in 4-manifolds*, Trans. Amer. Math. Soc. **264**, 353-380 (1981).

[Gi3] Gilmer, P. M., *Slice knots in S^3*, Quart. J. Math. Oxford **34**, 305-322 (1983).

[Gi4] Gilmer, P. M., *Classical knot and link concordance*, Comment. Math. Helv. **68**, 1-19 (1993).

[Gi5] Gilmer, P. M., *Signatures of singular branched covers*, Math. Ann. **295** (**4**), 643–659 (1993).

[GL] Gilmer, P. M., Livingston, C., *The Casson-Gordon invariant and link concordance*, Topology **31** , (**3**), 475–492 (1992).

[G] Gordon, C. McA., *Some aspects of classical knot theory*, Knot theory (Proc. Sem., Plans-sur-Bex, 1977), Lecture Notes in Math., **685**, Springer Verlag, Berlin, 1-60 (1978).

[K] Kirby, R. C., *The Topology of 4-manifolds*, Lecture Notes in Math **1374** Springer Verlag, Berlin (1989).

[Le] Levine, J., *Knot cobordism groups in codimension two*, Comment. Math. Helv. **44** 229–244 (1969).

[L] Lines D., *Cobordisme de noeuds fibrés et de leur monodromie*, Knots, braids and singularities (Plans-sur-Bex, 1982), 147–173, Monogr. Enseign. Math., **31**, Enseignement Math., Geneva, (1983).

[Li] Litherland, R. A., *Cobordism of satellite knots*, Four-Manifold Theory (Durham, N.H., 1982), Contemp. Math., **35**, Amer. Math. Soc., Providence, RI, 327–362 (1984).

[M] Murasugi, K., *On a certain numerical invariant of link types*, Trans. Amer. Math. Soc. **117** , 387–422 (1965).

[N] Naik, S. *Casson-Gordon invariants of genus one knots and concordance to reverses*, J. Knot Theory Ramifications **5**, 661–677 (1996).

[T] Tristram, A. G., *Some cobordism invariants for links*, Proc, Camb. Philos. Soc., **66**, 251-264 (1969).

[V] Viro, O. Ja. *Branched coverings of manifolds with boundary, and invariants of links. I*, Math. USSR-Izv. **7** 1239–1256 (1973).

Laboratoire I.R.M.A. Université Louis Pasteur
Strasbourg, France
and
Department of Mathematics, Louisiana State University
Baton Rouge, LA 70803, USA

Email: `vincent.florens@irma.u-strasbg.fr` and `gilmer@math.lsu.edu`