

УДК 512.555+517.982

A BECKENBACH–DRESHER TYPE INEQUALITY
IN UNIFORMLY COMPLETE f -ALGEBRAS

A. G. Kusraev

*To the memory of Gleb Akilov on the occasion
of the 90th anniversary of his birth*

A general form Beckenbach–Dresher inequality in uniformly complete f -algebras is given.

Mathematics Subject Classification (2000): 06F25, 46A40.

Key words: f -algebra, vector lattice, lattice homomorphism, positive operator.

An easy modification of the *continuous functional calculus* on unitary f -algebras as defined in [3] makes it possible to translate the Fenchel–Moreau duality to f -algebra setting and to produce some *envelope representations* results, see [8]. This machinery, often called *quasi-linearization* (see [2, 9]), yields the validity of some classical inequalities in every uniformly complete vector lattice [4, 5]. The aim of this note is to give general forms of Peetre–Persson and Beckenbach–Dresher inequalities in uniformly complete f -algebras.

The unexplained terms of use below can be found in [1] and [6].

1°. We need a slightly improved version of continuous functional calculus on uniformly complete f -algebras constructed in [3, Theorem 5.2].

Denote by $\mathcal{B}(\mathbb{R}_+^N)$ the f -algebra of continuous functions on \mathbb{R}_+^N with polynomial growth; i. e., $\varphi \in \mathcal{B}(\mathbb{R}_+^N)$ if and only if $\varphi \in C(\mathbb{R}_+^N)$ and there are $n \in \mathbb{N}$ and $M \in \mathbb{R}_+$ satisfying $|\varphi(\mathbf{t})| \leq M(\mathbf{1} + w(\mathbf{t}))^n$ ($\mathbf{t} \in \mathbb{R}_+^N$), where $\mathbf{t} := (t_1, \dots, t_N)$, $w(\mathbf{t}) := |t_1| + \dots + |t_N|$ and $\mathbf{1}$ is the function identically equal to 1 on \mathbb{R}_+^N . Denote by $\mathcal{B}_0(\mathbb{R}_+^N)$ the set of all functions in $\mathcal{B}(\mathbb{R}_+^N)$ vanishing at zero. Let $\mathcal{A}(\mathbb{R}_+^N)$ stands for the set of all $\varphi \in \mathcal{B}(\mathbb{R}_+^N)$ such that $\lim_{\alpha \downarrow 0} \alpha^{-1} \varphi(\alpha \mathbf{t})$ exists uniformly on bounded subsets of \mathbb{R}_+^N . Evidently, $\mathcal{A}(\mathbb{R}_+^N) \subset \mathcal{B}_0(\mathbb{R}_+^N)$. Finally, let $\mathcal{H}(\mathbb{R}_+^N)$ denotes the set of all continuous positively homogeneous functions on \mathbb{R}_+^N .

Lemma 1. *The sets $\mathcal{B}(\mathbb{R}_+^N)$, $\mathcal{B}_0(\mathbb{R}_+^N)$, and $\mathcal{A}(\mathbb{R}_+^N)$ are uniformly complete f -algebras with respect to pointwise operations and ordering. Any $\varphi \in \mathcal{A}(\mathbb{R}_+^N)$ admits a unique decomposition $\varphi = \varphi_1 + w\varphi_2$ with $\varphi_1 \in \mathcal{H}(\mathbb{R}_+^N)$ and $\varphi_2 \in \mathcal{B}_0(\mathbb{R}_+^N)$, i. e.*

$$\mathcal{A}(\mathbb{R}_+^N) = \mathcal{H}(\mathbb{R}_+^N) \oplus w\mathcal{B}_0(\mathbb{R}_+^N).$$

Moreover, $\varphi_1(\mathbf{t}) = \varphi'(0)\mathbf{t} := \lim_{\alpha \downarrow 0} \alpha^{-1} \varphi(\alpha \mathbf{t})$ for all $\mathbf{t} \in \mathbb{R}_+^N$.

◁ See [3, Lemma 4.8, Section 5]. ▷

2°. Consider an f -algebra E . Denote by $H(E)$ the set of all nonzero \mathbb{R} -valued lattice homomorphisms on E and by $H_m(E)$ the subset of $H(E)$ consisting of multiplicative functionals. We say that $\omega \in H(E)$ is *singular* if $\omega(xy) = 0$ for all $x, y \in E$. Let $H_s(E)$ denote the set of singular members of $H(E)$. Given a finite tuple $\mathbf{x} = (x_1, \dots, x_N) \in E^N$, denote by $\langle\langle \mathbf{x} \rangle\rangle := \langle\langle x_1, \dots, x_N \rangle\rangle$ the f -subalgebra of E generated by $\{x_1, \dots, x_N\}$.

DEFINITION. Let E be a uniformly complete f -algebra and $x_1, \dots, x_N \in E_+$. Take a continuous function $\varphi : \mathbb{R}_+^N \rightarrow \mathbb{R}$. Say that the element $\widehat{\varphi}(x_1, \dots, x_N)$ *exists* or is *well-defined* in E provided that there is $y \in E$ satisfying

$$\begin{aligned} \omega(y) &= \varphi(\omega(x_1), \dots, \omega(x_N)) \quad (\omega \in H_m(\langle\langle x_1, \dots, x_N, y \rangle\rangle)), \\ \omega(y) &= \varphi_1(\omega(x_1), \dots, \omega(x_N)) \quad (\omega \in H_s(\langle\langle x_1, \dots, x_N, y \rangle\rangle)), \end{aligned} \quad (1)$$

cp. [3, Remark 5.3 (ii)]. This is written down as $y = \widehat{\varphi}(x_1, \dots, x_N)$.

Lemma 2. *Assume that E is a uniformly complete f -algebra and $x_1, \dots, x_N \in E_+$, and $\mathbf{x} := (x_1, \dots, x_N)$. Then $\widehat{\mathbf{x}}(\varphi) := \widehat{\varphi}(x_1, \dots, x_N)$ exists for every $\varphi \in \mathcal{A}(\mathbb{R}_+^N)$, and the mapping $\widehat{\mathbf{x}} : \varphi \mapsto \widehat{\mathbf{x}}(\varphi) = \widehat{\varphi}(x_1, \dots, x_N)$ is the unique multiplicative lattice homomorphism from $\mathcal{A}(\mathbb{R}_+^N)$ to E such that $\widehat{dt}_j(x_1, \dots, x_N) = x_j$ for all $j := 1, \dots, N$. Moreover, $\widehat{\mathbf{x}}(\mathcal{A}(\mathbb{R}_+^N)) = \overline{\langle\langle x_1, \dots, x_N \rangle\rangle}$.*

◁ Take $\varphi \in \mathcal{A}(\mathbb{R}_+^N)$. In view of Lemma 1 $\varphi = \varphi_1 + w\varphi_2$ with $\varphi_1 \in \mathcal{H}(\mathbb{R}_+^N)$, $\varphi_2 \in \mathcal{B}_0(\mathbb{R}_+^N)$, and $w(\mathbf{t}) = |t_1| + \dots + |t_N|$. For $x \in E$ denote by $\dot{x} \in \text{Orth}(E)$ the multiplication operator $y \mapsto xy$ ($x \in E$). According to [5, Theorem 3.3] and [8, Theorem 2.10] we can define correctly $\widehat{\varphi}_1(x_1, \dots, x_N)$ in E and $\widehat{\varphi}_2(\dot{x}_1, \dots, \dot{x}_N)$ in $\text{Orth}(E)$, respectively. Now, it remains to put $\widehat{\varphi}(x_1, \dots, x_N) := \widehat{\varphi}_1(x_1, \dots, x_N) + \widehat{\varphi}_2(\dot{x}_1, \dots, \dot{x}_N)w(x_1, \dots, x_N)$ and check the soundness of this definition. Closer examination of the proof can be carried out as in the case of $\varphi \in \mathcal{A}(\mathbb{R}^N)$, see [3]. ▷

Lemma 3. *Assume that $\varphi \in \mathcal{A}(\mathbb{R}_+^N)$ is convex. Then for all $\mathbf{x} := (x_1, \dots, x_N) \in E^N$, $\mathbf{y} := (y_1, \dots, y_N) \in E^N$, and $\pi, \rho \in \text{Orth}(E)_+$ with $\pi + \rho = I_E$ we have $\widehat{\varphi}(\pi\mathbf{x} + \rho\mathbf{y}) \leq \pi\widehat{\varphi}(\mathbf{x}) + \rho\widehat{\varphi}(\mathbf{y})$, where $\pi\mathbf{x} := (\pi x_1, \dots, \pi x_N)$. The reverse inequality holds whenever φ is concave.*

◁ Let L be the order ideal generated by $\overline{\langle\langle x_1, \dots, y_N \rangle\rangle}$. Clearly, L is an f -subalgebra of E . If $\pi_0 := \pi|_L$ and $\rho_0 := \rho|_L$ then $\pi_0, \rho_0 \in \text{Orth}(L)$. For any $\omega \in H(L)$ there exists a unique $\widetilde{\omega} \in H_m(\text{Orth}(L))$ such that $\omega(\pi x) = \widetilde{\omega}(\pi)\omega(x)$ for all $x \in L$ and $\pi \in \text{Orth}(L)$, [3, Proposition 2.2 (i)]. If ω is nonsingular then $\alpha\omega$ is multiplicative for some $\alpha > 0$ [3, Corollary 2.5 (i)], and thus we may assume without loss of generality that $\omega \in H_m(L)$. By using (1), the convexity of φ , and the relation $\widetilde{\omega}(\pi) + \widetilde{\omega}(\rho) = 1$ we deduce

$$\begin{aligned} \omega(\widehat{\varphi}(\pi\mathbf{x} + \rho\mathbf{y})) &= \varphi(\widetilde{\omega}(\pi_0)\omega(\mathbf{x}) + \widetilde{\omega}(\rho_0)\omega(\mathbf{y})) \leq \widetilde{\omega}(\pi_0)\varphi(\omega(\mathbf{x})) + \widetilde{\omega}(\rho_0)\varphi(\omega(\mathbf{y})) \\ &= \widetilde{\omega}(\pi_0)\omega(\widehat{\varphi}(\mathbf{x})) + \widetilde{\omega}(\rho_0)\omega(\widehat{\varphi}(\mathbf{y})) = \omega(\pi\widehat{\varphi}(\mathbf{x}) + \rho\widehat{\varphi}(\mathbf{y})), \end{aligned}$$

where $\omega(\mathbf{x}) := (\omega(x_1), \dots, \omega(x_N))$. If ω is singular then by above definition we have $\omega(\widehat{\varphi}(\mathbf{x})) = \omega(\widehat{\varphi}_1(\mathbf{x}))$, $\omega(\widehat{\varphi}(\mathbf{y})) = \omega(\widehat{\varphi}_1(\mathbf{y}))$, and $\omega(\widehat{\varphi}(\pi\mathbf{x} + \rho\mathbf{y})) = \omega(\widehat{\varphi}_1(\pi\mathbf{x} + \rho\mathbf{y}))$. At the same time φ_1 is sublinear, since it coincides with the directional derivative of the convex function φ at zero, see Lemma 3. Thus, by replacing φ by φ_1 in the above arguments we again obtain $\omega(\widehat{\varphi}(\pi\mathbf{x} + \rho\mathbf{y})) \leq \omega(\pi\widehat{\varphi}(\mathbf{x}) + \rho\widehat{\varphi}(\mathbf{y}))$. It remains to observe that every $\omega_0 \in H(\langle\langle x_1, \dots, x_N \rangle\rangle)$ admits an extension to $\omega \in H(L)$ and thus $H(L)$ separates the points of $\langle\langle x_1, \dots, x_N \rangle\rangle$. ▷

Lemma 4. *If $\varphi \in \mathcal{A}(\mathbb{R}_+^N)$ is isotonic, then $\widehat{\varphi}$ is also isotonic, i. e. $\mathbf{x} \leq \mathbf{y}$ implies $\widehat{\varphi}(\mathbf{x}) \leq \widehat{\varphi}(\mathbf{y})$ for all $\mathbf{x}, \mathbf{y} \in E_+^N$. (The order in E^N is defined componentwise.)*

◁ Follows immediately from the above definition (1). ▷

3°. Everywhere below $(G, +)$ is a commutative semigroup, while E is a uniformly complete f -algebra and $f_1, \dots, f_N : G \rightarrow E_+$. Let $\mathcal{P}(M)$ stands for the power set of M . Assume that some set-valued map $\mathcal{F} : G \rightarrow \mathcal{P}(\text{Orth}(E)_+)$ meets the following three conditions:

- (i) π^{-1} exists in $\text{Orth}(E)$ for every $\pi \in \mathcal{F}(u)$,
- (ii) $\mathcal{F}(u) + \mathcal{F}(v) \subset \mathcal{F}(u+v) - \text{Orth}(E)_+$ for all $u, v \in G$, and
- (iii) the infimum (the supremum) of $\{\pi\widehat{\varphi}(\pi^{-1}\mathbf{f}(u)) : \pi \in \mathcal{F}(u)\}$ exists in E for each $u \in G$, where $\mathbf{f}(u) := (f_1(u), \dots, f_N(u)) \in E_+^N$ and $\pi^{-1}\mathbf{f}(u) := (\pi^{-1}f_1(u), \dots, \pi^{-1}f_N(u))$.

Lemma 5. *Given a function $\varphi : \mathcal{A}(\mathbb{R}_+^N)$ and a set-valued map $\mathcal{F} : G \rightarrow \mathcal{P}(\text{Orth}(E)_+)$ satisfying 3 (i–iii), we have the operator $g : G \rightarrow E$ ($h : G \rightarrow E$) well defined as*

$$g(u) := \inf_{\pi \in \mathcal{F}(u)} \{ \pi\widehat{\varphi}(\pi^{-1}\mathbf{f}(u)) \}, \quad \left(h(u) := \sup_{\pi \in \mathcal{F}(u)} \{ \pi\widehat{\varphi}(\pi^{-1}\mathbf{f}(u)) \} \right). \quad (2)$$

◁ By 3 (i) and Lemma 2 $\widehat{\varphi}(\pi^{-1}\mathbf{f}(u))$ exists in E and by 3 (iii) g and h are well defined. ▷

4°. Now we are able to state and prove our main result. A function $g : G \rightarrow F$ is said to be *subadditive* if $g(u+v) \leq g(u) + g(v)$ for all $u, v \in G$ and *superadditive* if the reversed inequality holds for all $u, v \in G$.

Theorem. *Suppose that the operators $g, h : G \rightarrow E$ are defined as in (2). Then:*

- (1) g is subadditive whenever f_1, \dots, f_N are subadditive and $\varphi \in \mathcal{A}(\mathbb{R}_+^N)$ is increasing and convex;
- (2) h is superadditive whenever f_1, \dots, f_N are superadditive and $\varphi \in \mathcal{A}(\mathbb{R}_+^N)$ is increasing and concave.

◁ We restrict ourselves to the subadditivity of g . The superadditivity of h can be proved in a similar way. Take $u, v \in G$ and let $\pi \in \mathcal{F}(u)$ and $\rho \in \mathcal{F}(v)$. By 3 (ii) we can choose $\sigma \in \mathcal{F}(u+v)$ with $\sigma \geq \pi + \rho$. In view of 3 (i) π, ρ , and σ are invertible. Taking subadditivity of $\mathbf{f} : G \rightarrow E^N$ and some elementary properties of orthomorphisms into account we have

$$\sigma^{-1}\mathbf{f}(u+v) \leq \sigma^{-1}(\mathbf{f}(u) + \mathbf{f}(v)) \leq \pi\sigma^{-1}(\pi^{-1}\mathbf{f}(u)) + \rho\sigma^{-1}(\rho^{-1}\mathbf{f}(v)).$$

Putting $\tau := \sigma - \pi - \rho$ and making use of Lemmas 3, 4 and 5 we deduce

$$\begin{aligned} g(u+v) &\leq \sigma\widehat{\varphi}(\sigma^{-1}\mathbf{f}(u+v)) \leq \sigma\widehat{\varphi}(\pi\sigma^{-1}(\pi^{-1}\mathbf{f}(u))) + \rho\sigma^{-1}(\rho^{-1}\mathbf{f}(v) + \tau\sigma^{-1}\mathbf{0}) \\ &\leq \pi\widehat{\varphi}(\pi^{-1}\mathbf{f}(u)) + \rho\widehat{\varphi}(\rho^{-1}\mathbf{f}(v)) + \sigma^{-1}\tau\widehat{\varphi}(\mathbf{0}) = \pi\widehat{\varphi}(\pi^{-1}\mathbf{f}(u)) + \rho\widehat{\varphi}(\rho^{-1}\mathbf{f}(v)). \end{aligned}$$

By taking infimum over $\pi \in \mathcal{F}(u)$ and $\rho \in \mathcal{F}(v)$ we come to the required inequality. ▷

REMARK 1. Suppose that the hypotheses of 3 (i–iii) are fulfilled for some fixed $u, v \in G$. Then the inequality $g(u+v) \leq g(u) + g(v)$ ($h(u+v) \geq h(u) + h(v)$) holds.

REMARK 2. An f -algebra E can be identified with $\text{Orth}(E)$ if and only if E has a unit element. Thus, above theorem remains true if E is a uniformly complete unitary f -algebra and the map $\mathcal{F} : G \rightarrow \mathcal{P}(E_+)$ satisfies the condition 3 (i–iii) with $\text{Orth}(E)$ replaced by E .

5°. For a single-valued map $\mathcal{F}(x) = \{f_0(x)\}$ ($x \in G$) with $f_0 : G \rightarrow \text{Orth}(E)_+$ we have the following particular case of the above Theorem, see [8].

Corollary 1. *Suppose that f_1, \dots, f_N are subadditive, $f_0 : G \rightarrow \text{Orth}(E)_+$ is superadditive, and $f_0(u)$ is invertible in $\text{Orth}(E)$ for every $u \in G$. Then, given an increasing continuous convex function $\varphi \in \mathcal{A}(\mathbb{R}_+^N)$, the Peetre–Persson inequality holds:*

$$f_0(u+v)\widehat{\varphi}\left(\frac{\mathbf{f}(u+v)}{f_0(u+v)}\right) \leq f_0(u)\widehat{\varphi}\left(\frac{\mathbf{f}(u)}{f_0(u)}\right) + f_0(v)\widehat{\varphi}\left(\frac{\mathbf{f}(v)}{f_0(v)}\right). \quad (3)$$

The reverse inequality holds in (3) whenever f_0, f_1, \dots, f_N are superadditive, and φ is an increasing concave function.

REMARK 3. The above theorem in the particular case of $E = \mathbb{R}$ was obtained by Persson [12, Theorems 1 and 2], while Corollary 2 covers the “single-valued case” by Peetre and Persson [11]. A short history of the Beckenbach–Dresher inequality is presented in [13]. Some instances of the inequality are also addressed in [9, 10].

6°. We need two more auxiliary facts. First of them is a generalized Minkowski inequality.

Lemma 6. *Let E and F be uniformly complete vector lattices, $f : E_+ \rightarrow F$ an increasing sublinear operator. If either $0 < \alpha \leq 1$ or $\alpha < 0$, then for all $x_1, \dots, x_N \in E$ we have*

$$f\left(\left(\sum_{i=1}^N |x_i|^\alpha\right)^{1/\alpha}\right) \leq \left(\sum_{i=1}^N f(|x_i|^\alpha)\right)^{1/\alpha}. \quad (4)$$

The reverse inequality holds if $f : E_+ \rightarrow F$ is superlinear and $\alpha \geq 1$.

◁ The function $\phi_\alpha(\mathbf{t}) = (t_1^\alpha + \dots + t_N^\alpha)^{1/\alpha}$ ($\mathbf{t} \in \mathbb{R}_+^N$) is superlinear if $0 < \alpha < 1$ and sublinear if $\alpha \geq 1$. In case $\alpha < 0$ we define $\phi_\alpha(\mathbf{t}) = 0$ whenever $t_1 \cdot \dots \cdot t_N = 0$ and then ϕ_α is superlinear on $\text{int}(\mathbb{R}_+^N)$. In all cases $\phi_\alpha \in \mathcal{H}(\mathbb{R}_+^N)$ and (4) follows from the generalized Jensen inequality in vector lattices, see [4, Theorem 5.2] and [7, Theorem 4.2]. ▷

Let A and B be uniformly complete unitary f -algebras, while $E \subset A$ is a vector sublattice. For every $x \in A_+$ and $0 < p \in \mathbb{R}$ the p -power x^p is well defined in A , see [3, Theorem 4.12]. If $x \in A_+$ is invertible and $p < 0$, then we can also define $x^p := (x^{-1})^{-p}$. It can be easily seen that $\omega(x^p) = \omega(x)^p$ for any $\omega \in H_m(A_0)$ with an f -subalgebra $A_0 \subset A$ containing x . Assume that $R : E \rightarrow B$ is a positive operator. Given $x \in A$ with $x^p \in E$, we define $R_p(x) := R(x^p)^{\frac{1}{p}}$. This definition is sound provided that x is invertible in A and $R(x^p)$ is invertible in B .

Lemma 7. *If $p \geq 1$ and $x_1, \dots, x_N \in A_+$ are such that $x_1^p, \dots, x_N^p \in E$ and $(x_1 + \dots + x_N)^p \in E$, then the inequality holds:*

$$R_p(x_1 + \dots + x_N) \leq R_p(x) + \dots + R_p(x_N). \quad (5)$$

The reversed inequality is true whenever $p \leq 1$, $p \neq 0$. (In case $p < 0$ the positive elements x_i and $R(x_i^p)$ are assumed to be invertible in A .)

◁ Denote $u_i := x_i^p$, $\alpha := 1/p$, and observe that $(u_1^\alpha + \dots + u_N^\alpha)^{\frac{1}{\alpha}} = \phi_\alpha(u_1, \dots, u_N)$ where $\phi_\alpha(u_1, \dots, u_N)$ is understood in the sense of homogeneous functional calculus. In particular, $(x_1 + \dots + x_N)^p = (u_1^\alpha + \dots + u_N^\alpha)^{\frac{1}{\alpha}} \in E$ for every $p \neq 0$. We need consider three cases. If $p \geq 0$ then by applying Lemma 6 to the right-hand side of the equality

$$R_p(x_1 + \dots + x_N) = R\left((u_1^\alpha + \dots + u_N^\alpha)^{\frac{1}{\alpha}}\right)^\alpha = (R(\phi_\alpha(u_1, \dots, u_N)))^\alpha$$

with $u_i^\alpha \in E$ replaced by x_i and making use of $R_p(x_i) = R(u_i)^\alpha$ ($i := 1, \dots, N$), we arrive immediately at the desired inequality (5). The same arguments involving the reversed version of (4) leads to the reversed inequality in (5) whenever $0 < p < 1$. Finally, in the case $p < 0$, again by Lemma 6, we have $R((u_1^\alpha + \dots + u_N^\alpha)^{\frac{1}{\alpha}}) \leq (R(u_1)^\alpha + \dots + R(u_N)^\alpha)^{\frac{1}{\alpha}}$ and rising both sides of this inequality to the α th power we get the reversed inequality (5). ▷

7°. Now, we can deduce a generalization of one more Beckenbach–Dresher type inequality due to Peetre and Persson [11].

Corollary 2. *Let $S : E \rightarrow F$ and $T : E \rightarrow \text{Orth}(F)$ be positive operators. Take $x_1, \dots, x_N \in A_+$ such that $x_i^\alpha, x_i^\beta, (\sum_{i=1}^N x_i)^\alpha, (\sum_{i=1}^N x_i)^\beta \in E$ ($i := 1, \dots, N$). If $p \geq 1$,*

$\beta \leq 1 \leq \alpha$, $\beta \neq 0$, and $T(x_i^\beta)$ are invertible in $\text{Orth}(F)$ whenever $\beta < 0$, then

$$\frac{\left(S\left(\left(\sum_{i=1}^N x_i\right)^\alpha\right)\right)^{p/\alpha}}{\left(T\left(\left(\sum_{i=1}^N x_i\right)^\beta\right)\right)^{(p-1)/\beta}} \leq \sum_{i=1}^N \frac{\left(S(x_i^\alpha)\right)^{p/\alpha}}{\left(T(x_i^\beta)\right)^{(p-1)/\beta}}. \quad (6)$$

◁ Put $G = E$, $f(x) := \mathbf{f}(x) := S(x^\alpha)^{1/\alpha}$, $f_0(x) := T(x^\beta)^{1/\beta}$, $N = 1$, and $\varphi(t) = t^p$ in Corollary 1. By Lemma 7 f is subadditive, f_0 is superadditive, and $f_0(x_i)$ is invertible in $\text{Orth}(F)$. Moreover, $\varphi \in \mathcal{A}(\mathbb{R}_+)$ is convex and increasing whenever $p \geq 1$. Now, the desired inequality is deduced by induction. ▷

REMARK 4. If $0 < p < 1$ then the concave function $\varphi(t) = t^p$ is not in $\mathcal{A}(\mathbb{R}_+)$ and we cannot guarantee the reversed inequality in (6). Nevertheless, in the case that F is a unitary f -algebra one can take $\varphi \in \mathcal{B}(\mathbb{R}_+^N)$ in Peetre–Persson’s inequality (3) and thus the reversed inequality is true in (6) whenever $0 < p \leq 1$, $\alpha, \beta \leq 1$, and $\alpha, \beta \neq 0$.

References

1. Aliprantis C. D., Burkinshaw O. Positive operators.—London etc.: Acad. Press Inc., 1985.—xvi+367 p.
2. Beckenbach E. F., Bellman R. Inequalities.—Berlin: Springer, 1983.
3. Buskes G., de Pagter B., van Rooij A. Functional calculus on Riesz spaces // Indag. Math.—1991.—Vol. 4, № 2.—P. 423–436.
4. Kusraev A. G. Homogeneous functional calculus on vector lattices.—Vladikavkaz: Southern Math. Inst. VSC RAS, 2008.—34 p.—(Preprint № 1).
5. Kusraev A. G. Functional calculus and Minkowski duality on vector lattices // Vladikavkaz Math. J.—2009.—Vol. 11, № 2.—P. 31–42.
6. Kusraev A. G., Kutateladze S. S. Subdifferentials: Theory and applications.—Dordrecht: Kluwer Academic Publ., 1995.—ix+398 p.
7. Kusraev A. G. Inequalities in vector lattices // Studies on Mathematical Analysis, Differential Equation, and their Applications / Eds. Korobeĭnik Yu. F., Kusraev A. G.—Vladikavkaz: SMI VSC RAS, 2010.—P. 82–96.—(Review of Science: The South of Russia).
8. Kusraev A. G., Kutateladze S. S. Envelopes and inequalities in vector lattices.—(to appear).
9. Mitrinović D. S., Pečarić J. E., Fink A. M. Classical and new inequalities in analysis.—Dordrecht: Kluwer, 1993.—xvii+740 p.
10. Pečarić J. E., Proschan F., Tong Y. L. Convex functions, partial orderings, and statistical application.—Boston a. o.: Academic Press, 1992.—xiii+469 p.
11. Peetre J., Persson L.-E. A general Beckenbach’s inequality with applications // Function Spaces, Differential Operators and Nonlinear Analysis. Pitman Res. Notes Math. Ser. 211.—1989.—P. 125–139.
12. Persson L.-E. Generalizations of some classical inequalities and their applications // Nonlinear Analysis, Function Spaces and Applications / Eds. Krbeć M., Kufner A., Opic B., Rákosník J.—Leipzig: Teubner, 1990.—P. 127–148.
13. Varošanec S. A generalized Beckenbach–Dresher inequality // Banach J. Math. Anal.—2010.—Vol. 4, № 1.—13–20.

ANATOLY G. KUSRAEV
 Southern Mathematical Institute
 Vladikavkaz Science Center of the RAS, Director
 RUSSIA, 362027, Vladikavkaz, Markus street, 22
 E-mail: kusraev@smath.ru

НЕРАВЕНСТВО ТИПА БЕККЕНБАХА — ДРЕШЕРА
В РАВНОМЕРНО ПОЛНЫХ f -АЛГЕБРАХ

А. Г. Кусраев

Установлено общее неравенство типа Беккенбаха — Дрешера в равномерно полных f -алгебрах.

Ключевые слова: f -алгебра, векторная решетка, решеточный гомоморфизм, положительный оператор.